Insect Pollination of Crops in Brazil

A Guide for Farmers, Gardeners, Politicians and Conservationists

DOI: 10.6094/UNIFR/151200

Insect Pollination of Crops in Brazil

A Guide for Farmers, Gardeners, Politicians and Conservationists

Prof Dr Alexandra-Maria Klein Prof Dr Breno M. Freitas Dr Isac G.A. Bomfim Dr Virginie Boreux Dr Felix Fornoff Dr Mikail O. Oliveira

"I have been an acerola grower for 13 years now. Right here in my plantation I see a lot of bees, they pollinate the plants, and if I have more bees to pollinate I think it will make it even better for the plants and yield because there are a lot of plants to pollinate. I've been working for a long time and I've seen a lot of bees, but I need to know which ones work for acerola and how I can promote their presence here. This pollination guide will be of much help."

Manoel Costa dos Santos

Acerola grower in Sítio Boa Vista dos Valentim, Maranguape, Brazil

"The book 'Insect Pollination of Crops in Brazil' has as main authors current protagonists in research on pollination globally, and presents as its main focus the situation of agricultural pollination in Brazil. This book also represents an important achievement, a partnership between various sectors of society, public and private, that enabled its production, seeking to facilitate horizontal knowledge about bees and their importance for agricultural crops, for the development of Brazil and food security."

Vera Lúcia Imperatriz Fonseca

Senior Professor at the University of Sao Paulo, Brazil

"This guide is an excellent resource for crop pollination researchers, students and agronomists interested in the finer details of crop pollination. This is a one-stop-shop that compiles information regarding pollination requirements, different varieties and flower visitor identity all in one place to optimize pollination in orchards and fields."

Romina Rader

Senior Lecturer in Community Ecology at the University of New England, Australia

"This book is a wonderful resource for Brazilian growers and agronomists working with pollinator-dependent crops, and for beekeepers wishing to promote the value of their stock for pollination services. Here you can easily find the pollination requirements of a crop, understand its floral biology and assess the relative importance of wild and managed pollinators."

Lynn Dicks

Professor at the University of Cambridge, United Kingdom

"The guide is an excellent tool not only for significant improvements in Brazilian agricultural production but also for students to have a holistic view of the pollinating processes, with integrated and very well illustrated information about plant species grown in Brazil, rich details about the biology and ecology of pollinating bees, as well as guidelines for maintaining healthier crops. It certainly improves our interest and understanding of such important insects as bees! "

Maiara Gonçalves

Visiting scholar at the University of Freiburg, Germany



The stingless bee Frieseomelitta longipes on a female açaí palm flower

 $\ensuremath{\mathbb{C}}$ 2020 Albert-Ludwigs University Freiburg, Nature Conservation and Landscape Ecology. All rights reserved.

DOI: 10.6094/UNIFR/151200

This book was edited using Scribus, an Open Source Desktop Publishing application.

Disclaimer

All information contained in this publication is for public use only. While care was taken to provide only correct information proved by scientific evidence at publication time, we do not claim completeness or correctness of all information displayed. Further expert knowledge is required depending on your circumstances.

Foreword

Farmers, gardeners and agronomists have detailed knowledge on how to maximise high quality crop production, especially in the context of external inputs including fertilization and pest management. However, detailed information on pollination is mainly available in the international scientific literature and often not accessible for local farmers or any person interested in growing crops while conserving bee diversity in Brazil.

Pollination is the first step in the reproduction of flowering plants. It is the transfer of pollen from the anthers (male reproduction organ) to the stigma (female reproduction organ) of flowers and therefore essential to develop seeds and to set fruits. Some staple food crops produce yield without insects or other animals and use wind for pollen transfer (e.g. rice, maize and other cereals). Other crops produce fruits or other plant organs we consume without pollination (e.g. banana or cassava). Nevertheless, the majority of crops depend to certain degrees on the transfer of pollen by animals. Only few of the major cultivated animal-pollinated crops around the globe are pollinated by birds and/or mammals such as durian in South-East Asia. Most other crops rely on insect pollinators..

Insect pollination can be provided by external inputs via the introduction of honeybee colonies or other managed pollinators during the crop blooming periods. Honeybee management for crop production is today established for some crop systems. The most famous example is the management of honeybees for almond production in California. Information how the naturally occurring pollinator species, especially wild bee species, contribute to overall crop production is largely missing, although they are crucial pollinators and can be more important than honeybees. The majority of wild bee species do not live in colonies, are therefore less dominant than honeybees and crop pollination increases when many species (biodiversity) share the pollination job. The diversity of wild bees is linked to the crop cultivation practices in the larger agricultural landscapes, in all places of the world. It is therefore crucial to know for each crop the main pollinator species to be able to promote and protect them in our farming systems or the surrounding landscape.

With our crop pollination guide for farmers, gardeners, agronomists, politicians and conservationists, we aim to transfer the current knowledge on crop pollination of the major crop species grown in Brazil. The overarching aims are to provide scientific knowledge of the dependency of a crop on insect pollination and to provide a list of relevant pollinator species for each crop. In detail, we show for each crop the growing regions in Brazil. We provide the pollination requirements, which we define as the plantmating system and the pollinators visiting the flowers. We also show common examples of the crops cultivated in Brazil. Our flower drawings show flower morphologies. Flowers and pollinators have co-evolved and therefore flower morphology is an important indicator for which pollinators can potentially access the nectar and pollen resources and most importantly which pollinators represent the fitting trait combination to successfully pollinate a flower. Besides providing a list of pollinator species for each crop listed in the scientific primary literature, we display information of pollinator groups in the second part of our crop pollination guide. This information provides the ecological knowledge to enhance the main pollinator groups on farmland, in farms, gardens and the wider landscape.

The idea of this crop pollination guide was developed at the University of Freiburg in collaboration with the Bayer Bee Care Centre of Bayer Crop Science in Germany.

Alexandra-M. Klein and Christian Maus initiated the guide and discussed the idea with Breno M. Freitas from the Departamento de Zootecnia at the Federal University of Ceará in Brazil and later with Martin Urban from Syngenta in Basel, Switzerland and with Juliana Jaramillo Salazar from the Bayer Bee Care Center. Virginie Boreux compiled a global crop pollination database at the University of Freiburg. Felix Fornoff of the University of Freiburg and Isac G. A. Bomfim and Mikail O. Oliveira from the Departamento de Zootecnia at the Federal University of Ceará checked the information and added information from additional publications from Brazil where necessary. They worked with Virginie Boreux on the layout and collected the photos. The University of Freiburg drew the flowers with help of a MSc student. Alexandra M. Klein and Breno M. Freitas wrote the remaining text and corrected the guide until final approval by all authors. Syngenta and Bayer Crop Sciences financed Virginie Boreux to extract information for our guide and financed the translation and print offs of this crop pollination guide. To make the guide accessible to farmers but also to students and the wider society, the English version is permanently stored at DOI:10.6094/UNIFR/151200 and the Portuguese version at DOI: 10.6094/UNIFR/151237. Pdfs are also available at the websites: https://www.nature.uni-freiburg.de/publications/Books, https://ppgzootecnia. ufc.br, https://beecare.baver.com/home and https://abelha.org.br/.

With this guide on insect pollinaton of crops in Brazil, we aim to increase the current knowledge of farmers and the wider society in Brazil. We hope to rise awareness of the importance of bees, show their beauty and fascinating diversity and to conserve and enhance pollinator diversity in crop-producing systems and landscapes.

Sincerely,

Alexandra Dein

Alexandra-M. Klein

&

AST'S

Breno M. Freitas

Contacts

Prof. Alexandra-Maria Klein

Nature Conservation and Landscape Ecology Faculty of Environment and Natural Resources University of Freiburg Tennenbacher Str. 4 79106 - Freiburg - Germany Email: alexandra.klein@nature.uni-freiburg.de Phone: + 49 761 203 67770

Prof. Breno M. Freitas

Setor de Abelhas Departamento de Zootecnia Centro de Ciências Agrárias Universidade Federal do Ceará CEP 60.356-970, Fortaleza - CE, Brazil Email: freitas@ufc.br Phone: + 55 85 3366 9220



Acknowledgements

We thank our MSc students of the University of Freiburg, Veronika Wenz for the amazing flower drawings and Tineke Materne for data cleaning. We kindly thank Dr. Maria Helena Pereira-Peixoto for the translation to Portuguese and MSc. Maiara Gonçalves for language corrections.

1 39×

÷.

The stingless bee Tetragonisca angustula collecting pollen of a strawberry flower

Content

Chapter 1: Introduction	
What is "pollination"?	4
What is a "plant-mating system"?	4
Level of dependence on pollinators	6
Main pollinator groups	7
Methodology	
Explanations	
Glossary	14
Chapter 2: Crop pollination	
Açaí palm/Açaí (<i>Euterpe oleracea</i>)	
Acerola/Acerola (Malpighia emarginata)	
Anatto/Urucum (Bixa orellana)	
Apple/Maçã (Malus domestica)	
Avocado/Abacate (Persea americana)	
Brazil nut/Castanha do Brasil (Bertholletia excelsa)	
Brazilian grapetree/Jabuticaba (Plinia cauliflora)	
Canola/Canola (Brassica napus)	
Cashew/Caju (Anacardium occidentale)	
Castor bean/Mamona (Ricinus communis)	
Citrus, Lime, Orange/Frutas cítricas (Citrus spp.)	
Cocoa/Cacau (Theobroma cacao)	
Coconut/Coco (Cocos nucifera)	
Coffee/Café (Coffea arabica)	
Common bean/Feijão (Phaseolus vulgaris)	
Cotton/Algodão (Gossypium hirsutum)	
Cowpea/Feijão-caupi (Vigna unguiculata)	
Cupuassu/Cupuaçu (Theobroma grandiflorum)	
Guava/Goiaba (<i>Psidium guajava</i>)	
Kiwi/Kiwi (<i>Actinidia deliciosa</i>)	
Lychee/Lichia (Litchi chinensis)	
Mango/Manga (<i>Mangifera indica</i>)	
Melon/Melão (<i>Cucumis melo</i>)	64
Oil palm/Dendê (<i>Elaeis guineensis</i>)	
Okra/Quiabo (Abelmoschus esculentus)	
Papaya/Mamão (<i>Carica papaya</i>)	
Passion fruit/Maracujá (Passiflora edulis)	
Peach/Pêssego (Prunus persica)	74

Pear/Pera (Pyrus communis)	76
Persimmon/Caqui (Diospyros kaki)	78
Plum/Ameixa (Prunus domestica)	80
Pumpkin/Abóbora (Cucurbita maxima)	
Rambutan/Rambuta (Nephelium lappaceum)	
Sesame/Gergelim (Sesamum indicum)	
Soursop/Graviola (Annona muricata)	
Soybean/ Soja (Glycine max)	90
Strawberry/Morango (Fragaria × ananassa)	92
Sunflower/Girassol (Helianthus annuus)	
Sweet pepper/Pimentão (Capsicum annuum)	96
Tomato/Tomate (Solanum lycopersicum)	98
Watermelon/Melancia (Citrullus lanatus)	100
Zucchini/Abobrinha (Cucurbita pepo)	102
Chapter 3: Comprehensive crop list	105
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators	105 109
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators Western honeybee (<i>Atis mellifera</i>)	105 109
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators Western honeybee (<i>Apis mellifera</i>) Bumblebees (<i>Bombus</i> spp.)	105 109 110 112
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators Western honeybee (<i>Apis mellifera</i>) Bumblebees (<i>Bombus</i> spp.) Stingless bees (Meliponini)	105 109 110 112 112
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators Western honeybee (<i>Apis mellifera</i>) Bumblebees (<i>Bombus</i> spp.) Stingless bees (Meliponini) Wild solitary bees	105 109 110 112 112 114
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators Western honeybee (<i>Apis mellifera</i>) Bumblebees (<i>Bombus</i> spp.) Stingless bees (Meliponini) Wild solitary bees Carpenter bees (<i>Xylocopa</i> spp.)	105 109 110 112 112 114 116
Chapter 3: Comprehensive crop list. Chapter 4: Managing pollinators. Western honeybee (<i>Apis mellifera</i>) Bumblebees (<i>Bombus</i> spp.). Stingless bees (Meliponini). Wild solitary bees. Carpenter bees (<i>Xylocopa</i> spp.). Oil-collectig bees (<i>Centris</i> spp.).	105 109 110 112 112 114 116 117
Chapter 3: Comprehensive crop list. Chapter 4: Managing pollinators. Western honeybee (<i>Apis mellifera</i>). Bumblebees (<i>Bombus</i> spp.). Stingless bees (Meliponini). Wild solitary bees. Carpenter bees (<i>Xylocopa</i> spp.). Oil-collectig bees (<i>Centris</i> spp.). Appendix, additional lists for crops and pollinators.	105 109 110 112 112 114 116 117 119
Chapter 3: Comprehensive crop list Chapter 4: Managing pollinators Western honeybee (<i>Apis mellifera</i>) Bumblebees (<i>Bombus</i> spp.) Stingless bees (Meliponini) Wild solitary bees Carpenter bees (<i>Xylocopa</i> spp.) Oil-collectig bees (<i>Centris</i> spp.) Appendix, additional lists for crops and pollinators Scientific literature	105 109 110 112 112 114 116 117 119 123



Introduction

Worldwide, most plants produce a fruit or a seed only after successful pollination. Pollination, the process of pollen transfer within or between fertile flowers depends on the plant species under consideration and is most commonly achieved through wind or insects. In about 75% of our global food crops pollination by insects (rarely by birds and mammals) can benefit crop production (Klein et al. 2007) and pollinators are important for optimal reproduction of the majority of wild plants in almost all places of the world (Ollerton et al. 2011). The world annual value of pollinators is estimated between US\$ 235-577 billons (IPBES 2016), while in Brazil it is estimated US\$ 42-43 billion (Giannini et al. 2015, BPBES 2019). Although many different insect species act as pollinators for specific plants, both managed and wild bees are the main contributors to crop pollination. The managed Western honeybee *Apis mellifera* is often favoured by farmers, as their high number of individuals per colony and long flight and foraging distances lead to potential pollination of large numbers of flowers and spacious crop areas that can be covered by managed pollination. They also provide honey, an important income source for many people including farmers worldwide, and are therefore important domestic animals. High levels of colony losses in some regions of the world over the past decades have increased concerns that scarcity of honeybees might negatively affect food production. While honeybees are the main managed pollinator, wild insects, mainly wild bees have been shown to contribute to the pollination of many crops (Garibaldi et al. 2013, Kleijn et al. 2015). While some crops, for example cocoa, are only pollinated by wild insects (Frimpong et al. 2009), wild bees may always provide some "insurance" to crop production, for example when honeybee colonies are weak or weather conditions are unfavourable for honeybee flight (Winfree et al. 2007, Brittain et al. 2013, Ellis et al. 2017). Pollination by wild bees is also shown to lead to better fruit quality compared to the pollination by honeybees (Klatt et al. 2014, Brittain et al. 2014, MacInnis & Forrest 2019). As of today, wild bee populations of Brazil have not been evaluated, but for some countries, especially in central Europe, insect abundance including pollinating species are discussed to be in severe decline (Potts et al. 2010 & 2016, Hallmann et al. 2018, IPBES 2019, Habel et al. 2019).

Wild bees include feral honeybees (Apis mellifera) but usually refer to all bee species excluding Apis mellifera. The ecology of wild bees is highly diverse. Some are specialized on a specific plant species; some are highly generalistic visiting a broad range of different flowers or even alternative food resources like fruits. Some species build their nests in dead wood, some in hollow sticks, some mine their nests into the ground, some create it with collected plant materials and others are cuckoo bees parasitizing other bee species (Westrich 1989, Michener 2000). Most species are solitary, where one female creates its own nests and cares for its own offspring, but depending on the species a variety of semisocial to highly eusocial behaviours exist (Michener 2000). Most bees collect nectar to cover their energy demands and collect pollen as the protein source to feed their offspring. The body size determines the flight distance between foraging and nesting areas; small bee species travel only up to 200m, while large bees cover distances of few kilometres (Zurbuchen et al. 2010). Hence, the wild bee ecology requires a nesting site and a foraging ground with flower resources within a limited area. While it is important to maintain suitable flowering plant species, including flowers of many crop species for wild bee species, protection of the breeding and nesting sites and structures is crucial for wild bee reproduction and sustainable populations.

In crop production areas, many different factors can reduce wild bee diversity (number of

species and number of individuals). Strong effects can be expected if flowers or nesting sites are limited. Flowers might be reduced by intensive land-use practices, including transformation to non-natural habitat, removal of wild plants or contamination of nectar and pollen. Application of certain pesticides, especially insecticides but also fungicides can also directly harm bees (IPBES 2019), especially when the application is not done in compliance with the safety measures stipulated on the product label. The conversion of non-crop habitat, potentially suitable for nesting wild bees, like the removal of dead wood, turning over of soils, and conversion to urban sealed areas can reduce or eliminate pollinator populations (Klein et al. 2018, Kleijn et al. 2018). For producers of crops that depend on insect pollination it is crucial to maintain a reliable pollinator community which should ideally constitute of as many species as possible, as each additional species increases the reliability of crop pollination even with variable seasonality and weather conditions (Brittain et al. 2013, Ellis et al. 2017). Without the knowledge of the ecology of wild bees, no directed conservation measures can be applied.

Flowers are as variable in morphology and phenology as are their pollinators. The knowledge on which pollinators are important for which flower is crucial for a management towards optimal pollination, including wild pollinators. Therefore, extensive information on the plant-mating system and the pollination system (which animal groups or species are visiting and pollinating the crop flowers) are needed for the assessment of factors potentially affecting pollinators and pollination.

This crop pollination guide of Brazil is presenting the following information:

- i) the main crop species grown in Brazil and their pollination requirements
- ii) the main pollinating insect groups and species found in Brazil
- iii) the management practices that can be used to protect and enhance pollinator populations.

What is "pollination"?

Pollination is the transfer of pollen grains from the male parts (anthers) of a flower to the female parts (stigmas) of a flower, leading to the fertilisation of the ovule or ovules of that flower (Fig 1). When compatible and viable pollen grains touch the stigma, they germinate on the stigma and grow a pollen tube down through the style into the ovary. There, the pollen tube ends and transfers the male genome. This leads to the fertilisation of the ovules containing the female genome. Pollen grains can be deposited on the stigma through biotic (insects, birds, mammals, etc.) or abiotic (wind, water, gravity) agents and pollen tube germination depends on the plant-mating system.

What is a "plant-mating system"?

The plant-mating system describes how a plant reproduces, from pollination to successful fertilisation of the ovule. In most crop species, pollination is required to set fruits and seeds, although in some cases asexual reproduction occurs. Some plants can produce seedless fruits without the transfer of pollen (parthenocarpy e.g. in banana or some pear or cucumber varieties). Other plants can produce fruits with seeds without the transfer of pollen (agamospermy e.g. in some apple varieties). Crops with these two asexual mating systems can however still benefit from the transfer of pollen (Delaplane et al. 2013). For instance production in parthenocarpic cucumber varieties, such as Hokushin, Yoshinari and Soudai, increases up to 40% (for Yoshinari) when bees are visiting flowers (Nicodemo et al. 2013).



Figure 1: Diagram of a flower showing the different parts involved in pollination and fertilisation of the flower.

All other mating forms reproduce only by successful pollination. Some flowers produce with their own pollen (autogamy, see Fig. 2a) but when pollinators have to transfer the pollen within the flower, the plant is called a self-pollinating species. When the pollen within the flower is transferred without pollinators, the plant is called an auto-pollinating species. Some plants need pollen of another flower of the same plant individual for successful pollination (geitonogamy, Fig. 2b). Geitonogamy is also a form of self pollination. Other crops need the pollen of a flower of a different plant individual (xenogamy, Fig. 2c), which is known as cross pollination. Many plants have a mixed-mating system and can either reproduce with pollen of the same or of other plant individuals, although pollen of different plant individuals often increases fruit and seed production (Delaplane et al. 2000).

Self pollination

Cross pollination



Figure 2: Main plant-mating systems and their mechanisms. a) autogamy, b) geitonogamy, c) xenogamy

The mating system usually differs between crop varieties. Some produce mainly or strictly with autogamy whereas others produce mainly or strictly with xenogamy. For some crops such as apple, cherry and pear, the compatible pollen has to come from a different variety. For example, xenogamy happens when pollen from a particular variety (e.g. Fuji) fertilises the ovules of a flower from another compatible variety (e.g. Gala). Similarly, autogamy happens in apple when pollen of one variety (e.g. Fuji) is fertilising the ovules of the same variety (in that case, Fuji). For apple, varieties are mostly self incompatible and require pollen from an apple tree of a different variety. Each variety is compatible with a subset of other varieties, e.g. Gala is compatible with Fuji and Eva with Princesa. Unfortunately, up to date information of the mating system is hardly available at the variety level for most Brazilian crops. We therefore provide the most common mating system for a crop if no detailed information of different varieties are available.



Figure 3: Yield increase (mean and standard error) through presence of pollinators compared to pollinator exclusion (only wind and self pollination). Yield measures were averaged per variety, crop and study and can include values of e.g. seed number, fruit weight and fruit set. The analysis is based on all publications evaluating yield changes published before 2018, including studies outside of Brazil. The number of studies analysed is given within the bars.

When a plant can partly produce with pollen of the same plant (self pollination) the plant is also termed as partially self fertile or partly self-pollinated. Examples of partially selffertile species include coffee and cucumber. Coffee (*Coffea arabica*) does not require cross pollination (xenogamy) to set fruits, but its production can be increased when bees, likely transporting pollen from different coffee plants, are visiting flowers (Klein et al. 2013). Another example was given above with an increased cucumber production when bees cross-pollinated primarily partheocarpic cucumber varieties (Nicodemo et al. 2013). Therefore, it is usually impossible to assign a crop to only one mating system. In our guide, we indicate the mating systems described in the scientific literature.

Level of dependence on pollinators

The level of pollinator dependency presents the probability of how many flowers turn into fruits when an optimal number of pollinators were available or when pollen was experimentally transferred via hand-cross pollination. The pollinator dependency is given in four categories (essential, high, modest, little) following the mean percent fruit set calculated from the scientific literature according to Klein et al. (2007). Some examples are shown in Fig. 3 which summarizes the results of yield measures obtained from many studies on the comparison of flowers with pollinator exclosure to flowers with insect pollinators present.

Main pollinator groups

There is a substantial diversity of pollinators across the globe. Even though mammals, such as bats or squirrels, pollinate some crop species (e.g. agave, *Agave* spp.; pitaya, *Hylocereus* spp.; *Mucuna macrocarpa*), most crops across the globe are pollinated by insects. Among insect pollinators, honeybees, stingless bees, wild bees and flies (especially hoverflies) are the main and most common pollinators. All crops presented in this guide are pollinated by insects.

In Brazil, and for the crop species presented in this guide, bees were by far the most commonly observed pollinators. However, this is not always the case. For example, certain beetle species are the best pollinators of oil palm (*Elaeis guineensis*) and certain midges pollinate cocoa (*Theobroma cacao*). For this guide, we used a database of scientific literature to compile lists of pollinator identities and extract information on the species-specific pollination success (Table 1 and 2, see methods for information on the database).

Table 1: List of the 20 most common pollinator species visiting crop flowers in Brazil. Occurrence represents the number of scientific papers in which a pollinator species was observed visiting crop flowers. Table 2: List of the 20 most common bee genera visiting crop flowers in Brazil. Occurrence represents the number of scientific papers in which a bee genus was observed visiting crop flowers

Species	Occurrence	Genus	Occurrence
Apis mellifera	49	Centris	75
Trigona spinipes	24	Xylocopa	54
Xylocopa frontalis	13	Apis	52
Bombus morio	11	Trigona	51
Exomalopsis analis	11	Bombus	30
Eulaema nigrita	10	Exomalopsis	30
Xylocopa grisescens	9	Eulaema	25
Centris tarsata	9	Epicharis	24
Centris flavifrons	8	Augochloropsis	21
Centris aenea	8	Augochlora	20
Xylocopa cearensis	6	Melipona	19
Paratrigona lineata	6	Plebeia	11
Trigona fuscipennis	6	Frieseomelitta	10
Centris fuscata	6	Pseudaugochlora	10
Oxaea flavescens	5	Trigonisca	10
Tetragonisca angustula	5	Lasioglossum	9
Eulaema cingulata	5	Oxaea	8
Bombus atratus	5	Paratrigona	8
Melipona quadrifasciata	5	Partamona	8
Centris sponsa	4	Ceratina	7

Honeybees

Managed Western honeybee

(Apis mellifera, Linnaeus, 1758)

The Western honeybee is a highly eusocial species native to Europe, Africa and the Middle East. It is now present in all continents, except Antarctica. Due to its generalistic feeding behaviour, large colonies usually comprising 40,000 -60,000 bee workers but sometimes reaching up to 80,000, its ease of breeding and managing in manufactured hives and its adaptability to different environment, the Western honeybee is



Honeybee (Apis mellifera) on an avocado flower

the bee species most commonly managed for pollination of agricultural crops. In Brazil, the use of honeybees as managed pollinators is still limited and done in large scales mainly in melon and watermelon fields in the northeastern region and apple and peach orchards in the southern part of the country. However, the introduction of honeybee colonies for pollination of canola and soybean has increased recently after researchers demonstrated considerable yield gains (Rosa et al. 2011, Milfont et al. 2013).

Africanized honeybee

The term 'Africanized honeybee' refers to the polyhybrid bee produced from the accidental and uncontrolled crossing of four European sub-species of *Apis mellifera* (*A. m. mellifera*, *A.m. ligustica*, *A.m. carnica*, *A. m. caucasica*) introduced to Brazil in the 1800's and an African race (*A. m. scutellata*) brought to the country in 1956. However, genetic studies suggest that this miscegenation was not even throughout the country and while in southern parts it is evident, colonies in other regions are genetically close to *A.m. scutellata* leading some scientists to simply use the term Africanized honeybees to refer to African honeybees in the Americas. These bees have been naturalized, are now dominant in the



whole country and are usually kept for the production of honey, pollen, propolis, wax and also for crop pollination.

Across the world, 11 species of honeybees (Apis spp.) exist. In this guide we summarize literature that does not always discriminate between Africanized honeybees and Western honevbees. both subspecies of Apis mellifera. To account for this uncertainty and differentiate any kind of subspecies originating from mellifera Apis from other honeybees we use the term "honeybee".

Honeybee (Apis mellifera) on a pear flower

Social wild bees

Many different social and eusocial wild bees occur and act as important pollinators. These include bumblebees (Bombus spp.) as well as stingless bees (e.g. Melipona spp.). In Brazil, their contribution to crop pollination comes from feral colonies nesting around cultivated areas. Only recently, a few stingless bee species are being studied aiming for their use as managed pollinators in crops. However, the small colony size and slow build up of the population associated with



Stingless bees (Scaptotrigona sp.) on rambutam flowers

slow colony reproduction makes it difficult to keep them for crop pollination at a larger scale. Unlike other bumblebee species around the world, Brazilian bumblebees are extremely defensive and attack fiercely everyone approaching their nests. Due to this highly defensive behaviour, their colonies are usually exterminated by farmers when naturally occurring on their farms. Scientists and bee breeders avoid using them in experiments to prove their value as crop pollinators. The import of foreign bumblebee species is forbidden by law in Brazil.

Solitary wild bees

This group covers a substantial array of wild bees, which are very different in habits and morphology. It ranges from the large carpenter bees (Apidae) to the small sweat bees (Halictidae). Despite having solitary nesting behaviour, many species are gregarious and nest side by side building up large natural populations becoming important pollinators for

wild and cultivated plants. Some species of solitary bees are already bred in captivity and used for pollination purpose in some countries around the world. In Brazil, despite the existence of a great number of solitary bee species proven important pollinators of crops, there is no systematic breeding and use of any species as pollinator. The predominantly tropical climate of Brazil allows bees to be active all-year round. and the high diversity of solitary bees seems promising for the systematic management of some species as crop pollinators.



Solitary bee (Augochlora sp.) on a tomato flower



Solitary wasp on a cotton flower

Wasps

Wasps comprise a group of insects closely related to bees and like bees, there are solitary and social species. However, unlike bee larvae that feed on nectar and pollen, wasp larvae are carnivorous and feed on spiders, caterpillars or other insects hunted by the adult wasps. Despite this feeding behaviour of their brood, adult wasps usually feed on nectar and pollen, hence visiting and sometimes pollinating flowers. Some plants are even strongly dependent on wasp pollination. For example in the well-known

pollination coevolution involving fig wasps (Agonidae) and wild figs (*Ficus* spp.), fruits are only produced after wasp pollination. In general wasps are extremely important to agriculture because, although there are no crops in this guide entirely dependent on wasps for pollination, they contribute to the pollination of a variety of crops such as raspberries, and also prey on important agricultural pests such as caterpillars, aphids and cicadas. Solitary wasps contributing to pollination and pest control can be supported by the same practices listed for solitary bees in the chapter "Managing pollinators".



Biting midge on a cocoa flower

Flies

Flies belong to an ancient insect group characterized by having only one pair of fully developed wings. So far 160,000 species were named but a multitude of undescribed species is expected to exist worldwide (Borkent et al. 2018). Many fly species are valuable pollinators of flowers especially of the families Bombylidae, Tachinidae and Syrphidae. Flies show a variety of forms and sizes and include for example short- to long-tongued species. They pollinate a variety of wild and cultivated plants and can be important for increasing yield in crops such as mango (*Mangifera indica*) and are essential for cocoa (*Theobroma cacao*) yield and seed production of some vegetables.



Curculionid beetles on oil plam flowers

Beetles

Beetles are the most species-rich insect order worldwide, but in the context of crop production commonly seen as crop pests rather than pollinators. Indeed, many of them are relevant pests but others are important pollinators of plants in the tropics, particularly in Brazil. Besides many crop species where they can contribute to increase yield, some crops of local, national and worldwide importance such as custard apple (*Annona reticulata*), soursop (*Annona muricata*), and palm oil (*Elaeis guineensis*) are totally dependent on beetle species to set and produce fruits.

Methodology

Crop selection

Major crops of Brazil for which pollinators are important were selected for this guide, based on the authors' expert knowledge. With this, our aim was to cover most of locally and generally important crops of Brazil. However, Brazil is home to many more known crop species and others are still not described or valuated for cultivation. Even though, this guide is not complete it covers a wide range of the major crops in Brazil. For completeness and as an outlook, we composed a list of all generally known crop species of Brazil, including our estimate of pollinator attractiveness and insect pollination dependency at page 105-107.

Data on crops and pollinators

Data concerning crops grown in Brazil, their plant-mating system, varieties, pollinators as well as information concerning fruit set and seed set were extracted from a database on crop pollination collected between 2012-2018 at the University of Freiburg, Germany. When data were not available for Brazil, data from other countries in South America were used from the database. If no such data were available in the database, the crops were excluded from the guide. Bee pollinators observed in South America but not Brazil were checked against Moure's database (http://moure.cria.org.br/credits) to find if their range includes Brazil as well. If so, they were included in the pollinator list of the crop, if not they were removed. Species of midges were similarly checked using the publication of Santarém and Felippe-Bauer (2017) and species of ants the publication of Ulysséa et al. (2011). The database at University of Freiburg was built using scientific publications on crop pollination. Publications were identified and gathered using the search engines ISI Web of Knowledge (http://apps.webofknowledge.com/) and Google Scholar (http://scholar.google.com/). For each crop, the following search strings were used on both websites: i) scientific name of the species AND pollin*, ii) scientific name of the species AND pollen, iii) scientific name of the species AND nectar iv) common name of the species AND pollin^{*}, v) common name of the species AND pollen, vi) common name of the species AND nectar. Data on crops (varieties studied, breeding system, flowering and harvesting time), pollinators (name, forage target, abundance) and their impact on production were extracted from the scientific publications.

Disclaimer crop attractiveness score

Any classification of attractiveness has been made based on expert judgement and using the best scientific knowledge available. This classification does not represent an absolute truth because attractiveness is depending on many variable factors such as variety, manners of cultivation, environment, etc. For these reasons, the authors cannot assume any liability for decisions made using this classification.

References

- Borkent A., Brown B.V., Adler P.H., et al. (2018) Remarkable fly (Diptera) diversity in a patch of Costa Rican cloud forest: Why inventory is a vital science. Zootaxa 4402: 53-90
- BPBES (2019) Brazilian platform on biodiversity and ecosystem services. In: BPBES. https://www.bpbes.net.br/en/press/. Accessed 14 Jun 2019
- Brittain C., Kremen C., Garber A., Klein A.M. (2014) Pollination and plant resources change the nutritional quality of almonds for human health. PLOS ONE 9:e90082

Brittain C., Williams N., Kremen C., Klein A.M. (2013) Synergistic effects of non-Apis bees and honey bees

for pollination services. Proc R Soc B 280: 20122767

- Milfont de O.M., Rocha E.E.M., Lima A.O.N., Freitas B.M. (2013) Higher soybean production using honeybee and wild pollinators, a sustainable alternative to pesticides and autopollination. Environmental Chemistry Letters 11: 335-341
- Delaplane K.S., Mayer D.R., Mayer D.F. (2000) Crop pollination by bees. Cabi Publishing, Wallingford, UK
- Ellis C.R., Feltham H., Park K., et al. (2017) Seasonal complementary in pollinators of soft-fruit crops. Basic and Applied Ecology 19: 45-55
- Frimpong E.A., Gordon I., Kwapong P.K., Gemmill-Herren B. (2009) Dynamics of cocca pollination: tools and applications for surveying and monitoring cocca pollinators. International Journal of Tropical Insect Science 29: 62-69
- Garibaldi L.A., Steffan-Dewenter I., Winfree R., et al. (2013) Wild pollinators enhance fruit set of crops regardless of honey bee abundance. Science 339: 1608-1611
- Giannini T.C., Cordeiro G.D., Freitas B.M., et al. (2015) The dependence of crops for pollinators and the economic value of pollination in Brazil. Journal Economic Entomology 108: 849-857
- Habel J.C., Samways M.J., Schmitt T. (2019) Mitigating the precipitous decline of terrestrial European insects: Requirements for a new strategy. Biodiversity Conservation 28: 1343-1360
- Hallmann C.A., Sorg M., Jongejans E., et al. (2017) More than 75 percent decline over 27 years in total flying insect biomass in protected areas. PLOS ONE 12: e0185809
- IPBES (2019) Global assessment of biodiversity and ecosystem services, Brondizio E. S. Settele J. Díaz S. and Ngo H. T. (editors). IPBES Secretariat, Bonn, Germany
- Klatt B.K., Holzschuh A., Westphal C., et al. (2014) Bee pollination improves crop quality, shelf life and commercial value. Proceedings of the Royal Society B: Biological Sciences 281: 20132440
- Kleijn D., Biesmeijer K., Dupont Y.L., et al. (2018) Bee conservation: Inclusive solutions. Science 360:389-390
- Kleijn D, et al. (2015) Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. Nature Communications 6: 7414
- Klein A.M., Steffan-Dewenter I., Tscharntke T. (2003) Bee pollination and fruit set of *Coffea arabica* and *C. canephora* (Rubiaceae). American Journal of Botany 90: 153-157
- Klein A.M., Boreux V., Fornoff F., et al. (2018) Relevance of wild and managed bees for human well-being. Current Opinion in Insect Science 26: 82-88
- Klein A.M., Vaissière B.E., Cane J.H., et al. (2007) Importance of pollinators in changing landscapes for world crops. Proceedings of the Royal Society B: Biological Sciences 274: 303-313
- MacInnis G, Forrest JRK (2019) Pollination by wild bees yields larger strawberries than pollination by honey bees. Journal of Applied Ecology 56: 824-832
- Michener C.D. (2000) The Bees of the World. JHU Press. Baltimore and London, USA and UK
- Nicodemo D., Malheiros E.B., Jong D.D., Couto R.H.N. (2013) Enhanced production of parthenocarpic cucumbers pollinated with stingless bees and Africanized honey bees in greenhouses. Semina: Ciências Agrárias 34: 3625-3634
- Ollerton J., Winfree R., Tarrant S. (2011) How many flowering plants are pollinated by animals? Oikos 120: 321-326
- Potts S.G., Biesmeijer J.C., Kremen C., et al. (2010) Global pollinator declines: trends, impacts and drivers. Trends in Ecology & Evolution 25: 345-353
- Potts S.G., Imperatriz-Fonseca V., Ngo H.T., et al. (2016) Safeguarding pollinators and their values to human well-being. Nature 540: 220-229
- Rosa A. de S., Blochtein B., Lima D.K. (2011) Honey bee contribution to canola pollination in Southern Brazil. Scientia Agricola 68: 255-259
- Santarém M.C.A., Felippe-Bauer M.L. (2017) Brazilian species of biting midges. Rio de Janeiro, 67p., Brazil
- Ulysséa M.A., Cereto C.E., Rosumek F.B., et al. (2011) Updated list of ant species (Hymenoptera, Formicidae) recorded in Santa Catarina State, southern Brazil, with a discussion of research advances and priorities. Revista Brasileira de Entomologia 55: 603-611

Westrich P. (1989) Die Wildbienen Baden-Württembergs. E. Ulmer, Stuttgart, Germany

- Winfree R., Williams N.M., Dushoff J., Kremen C. (2007) Native bees provide insurance against ongoing honey bee losses. Ecology Letters 10: 1105-1113
- Zurbuchen A., Landert L., Klaiber J., et al. (2010) Maximum foraging ranges in solitary bees: only few individuals have the capability to cover long foraging distances. Biological Conservation 143: 669-676

Explanations

Attractiveness score: To give an indication of the attractiveness of each crop, we used a scale ranging from 1 to 5, as follows:

- 1 Crop has a low or no attractiveness for animal pollinators
- 2 Crop has a low to medium attractiveness for animal pollinators
- **3** Crop has a **medium** attractiveness for animal pollinators
- 4 Crop has medium to high attractiveness for animal pollinators
- 5 Crop is **highly** attractive to animal pollinators
- NA degree of attractiveness is unknown

Cultivation: The type of cultivation (open fields, shaded, agroforests etc.) which might be relevant from the pollinators` perspective.

Distribution map: Distribution maps were drawn based on information available at floradobrasil (http://floradobrasil.jbrj.gov.br/) and completed using the authors' knowledge.

Flower description: Hand drawn lateral cross section of flowers showing reproductive organs based on templates, available copyright free at the repository plantillustrations.org (see http://plantillustrations.org, under creative commons licence). Further information describing the flower morphology and organs is given in the text.

Flowering and harvest periods: If available, flowering times were taken from the data base and checked by Brazilian authors of this guide. Many crops are grown at different times of the year throughout Brazil. Therefore, this information is only a rough approximation and may not be applicable in all regions.

Growing areas: List of Brazilian states (abbreviated, see list of abbreviations) where the crop is grown.

Pollinator dependency: Indicating the dependency of a crop on pollinating insects for optimal fruit and seed production. Based on experiments comparing seed and fruit production with and without animal pollinators the following categories were assigned:

Essential - pollinators are essential for most varieties (production reduction by \geq 90% without pollinators)

High - high production increase with insect pollinators (40 - \leq 90% reduction without pollinators)

Modest - modest production increase with insect pollinators (10 - 40% reduction without pollinators)

Little - little production increase with insect pollinators (>0 - <10% reduction without pollinators)

Main varieties grown in Brazil: Only the most commonly cultivated and established varieties are listed.

Other information: Information relating to fruit set or seed set. When available taken

from Brazilian or other South American papers of the database or any additional information according to the knowledge and experiences of the authors.

Plant-mating system: Description of the distribution of mating units in a plant population (see glossary for detailed descriptions).

Pollination requirements: More detailed information on the plant-mating system together with the flower morphology and physiology to better understand the pollination requirements.

References: List of references providing information mostly on the pollinator species but also on additional information of the crop.

Status: Native/cultivated (i.e. introduced)/naturalised.

Table of pollinators: List of flower-visiting insect species found in the scientific studies of the database. Common names were entered if available; otherwise, the common name of the genus was used. We describe the sociality of the pollinators in terms of "eusocial", "social" or "solitary".

Glossary

Agamospermy: Flower sets fruits and seeds without the transfer of pollen/pollination

Agroforest: Area used for the production of one to several types of understory crop and one to several crop trees and/or bushes

Androdioecious: Some plant individuals with male and female reproduction organs on the same plant individual and other individuals with male flowers

Androecious: Plant individuals produce only male flowers

Andromonoecious: Bisexual and male flowers on the same plant individual

Anemophilous: Wind pollinated

Anther: Head of the stamen producing the pollen

Auto pollination: Form of self pollination in which the flower transfers pollen from its anthers to its stigma without the aid of a pollinating agent (pollinator, wind, etc.)

Autogamy: Flower sets fruits and seeds with pollen of it's own stamens

Bract: A modified leave usually associated to flowers, it serves to protect the flower and sometimes to attract pollinators

Buzz pollination: Usually flowers with porous anthers require bees to buzz (sonicate) the flowers to loosen the pollen. This behaviour called "buzz pollination" is specific to certain species of bees, such as *Bombus* spp. The insect vibrates its flight muscles at high frequency when visiting a flower, releasing the pollen enclosed in the anthers. Some flowers, e.g. tomato flowers, are pollinated by buzz pollination

Carpel: The female organ of a flower consisting of four major parts: stigma, style, ovary, ovules. When several carpels in a flower are connate, the whole structure is termed pistil

Cauliflorous: A plant with the flowers at the woody stem or trunk

Cleistogamous: Pollination occurring in unopened flowers. Virtually all plants that produce cleistogamous flowers also produce flowers that open (chasmogamous flowers)

Cross pollination: The transfer of pollen from the male reproductive organ (anther) of

one plant individual to the female reproductive organ (stigma) of another plant individual. This mechanism requires abiotic (e.g. wind, water) or biotic (animals) agents and is termed xenogamy

Dichogamous (dichogamy): Bisexual flower, but female function and male function are separated in time

Dioecious: Individual flowers are either male or female, but only one sex per plant individual occurs. Therefore, the presence of male and female plants is required to set fruits and seeds

Distylous (distyly): Species with two morphologically different flower types that are self-incompatible, but cross-compatible

Entomophilous: Plant requires insect pollination for optimal fruit set

Eusocial: see Social/Sociality

Field: Area used for the production of fruits from forbs, herbs or grasses

Filament: Stalk attached to the flower and holding the anther

Flower visitor: An animal that touches a flower. Even though not all flower visitors contribute to pollination, many flower visitors carry pollen at least accidentally and often act as pollinators. Although we can often not distinguish between pollinators and flower visitors from the scientific data we refer to all flower visitors as potential pollinators as the chance that a flower-visiting species act as a pollinator is high

Foraging resources (for pollinators): Many insect pollinators (mainly bees) usually use two types of food resources, pollen and nectar, which they collect from flowers

Geitonogamy: Flower sets fruits and seeds after pollination with pollen of another flower of the same plant individual. This needs a pollination agent such as a biotic pollinator or wind

Gynodioecious: Some plant individuals with male and female reproductive organs on the same individual and other individuals with female flowers only

Gynoecious: Plant individuals produce only female flowers

Half-inferior ovary: Ovary embedded or surrounded by the receptacle. This ovary position is also termed subinferior or half superior

Hermaphrodite: Flowers have both male and female reproductive organs

Home garden: Area used for the production of a variety of crops used for subsistence

Inferior ovary: Ovary lies below the attachment of other floral parts

Inflorescence: Cluster of flowers arranged on a stem that is composed of a main branch or an arrangement of branches (compound inflorescences)

Mellitophilous: Plant requires bee pollination for optimal fruit set

Mixed-mating system: A combination of any of the different main mating systems: agamospermy, parthenocarpy, autogamy, geitonogamy and xenogamy. We refer to a mixed-mating system when cross and self pollination occurs. Otherwise, we specify the mix of the mating system for specific crops

Monoecious: Individual flowers are either male or female but on the same plant individual

Nesting resources (for pollinators): Insect pollinators, in particular bees, nest in a

variety of mediums, including sandy soil (e.g. ground-nesting bees such as *Centris* spp.) and hollow twigs and branches (e.g. cavity-nesting bees such as *Tetrapedia* spp.)

Orchard: Area used for the production of fruits of trees and bushes

Ovary: The swollen location at the base of the pistil in a flower that houses the ovules containing the female gamete

Ovules: The flower eggs located inside the ovary. Successfully pollinated they will grow into seeds

Panicle: Inflorescence with many branches

Parthenocarpy: Flower sets seedless fruits without the transfer of pollen/pollination

Pedicel: Stem that attaches a single flower to the inflorescence

Perianth: Petals and sepals together when it is possible to distinguish between the two. Otherwise, it is termed perigonium

Perigonium: The non-reproductive part of the flower, and structure that forms an envelope surrounding the sexual organs

Petals: Modified leaves surrounding the reproductive parts in a flower. Petals are often brightly coloured to attract pollinators. Taking all petals in a flower together is termed corolla

Pistil: The pistils of a flower are considered to be composed of carpels. The gynoecium may consist of one or more pistils

Pistillode: Sterile usually rudimentary (reduced) pistil

Plantation: Area used for the production of usually one type of cash crop (plant (-part) sold to the market)

Plant-mating system: The plant-mating system describes the distribution of mating units in a plant population. In flowering plants it refers to self pollination (self fertilization), to cross pollination (outcrossing) and to reproduction without mating. In this guide we refer to the following forms of mating systems: agamospermy, parthenocarpy, autogamy (self pollination and auto pollination), geitonogamy, xenogamy and a mix of these mating systems termed mixed-mating system. We explain these seven forms of mating systems separately

Pollinizer: Plant (sometimes a specific cultivar) that provides abundant, compatible, and viable pollen

Pollination: The transfer of pollen grains from the male reproductive organs (anthers) to the female reproductive organs (stigmas)

Pollinator: An insect (or another animal), which interacts with a flower, usually by landing or climbing on it, and contributes to pollination. The animal transports pollen on its body, which touches the female reproductive organ of the flower (stigma). If this happens, the flower-visiting insect is usually classified as a pollinator. In only few studies on flower visiting insects, further analyses are carried out in how far flower visitors are indeed pollinating, for example by looking at the pollen tube growth in the style, or checking the fruit set of the visited flower after several days

Polygamous: Hermaphrodite, male and female flowers on the same individual

Protandrous, protandry: Anthers mature before the stigmas

Protogynous hermaphroditism: A flower changes from female to hermaphrodite

Protogynous, protogyny: Stigmas mature before the anthers

Radial symmetrical flowers: Flowers can be divided into three or more identical sectors, which are related to each other by rotation around the flower centre

Receptacle: The part of a flower stalk where the parts of the flower are attached

Self-compatible plants: A species which can set fruits or seeds by self-pollinating its stigmas with its own pollen. Self-compatible plants do not require cross pollination for successful production, but can benefit from it

Self-incompatible plants: A species that can set fruits only when pollen comes from different plant individuals from the same species. Self-incompatible plants require wind or animal pollination, and do not set fruits by self pollination. Self incompatibility is usually associated to cross pollination

Self pollination: The transfer of pollen from the male reproductive organ (anther) of one flower to the female reproductive organ (stigma) in the same flower (autogamy) or to the stigmas of another flower of the same plant (geitonogamy). This can be done passively (auto pollination) or via biotic or abiotic pollinating agents

Sepals: Modified leaves usually surrounding the petals. They are usually green and function to protect the reproductive flower organs in a bud. Taking all petals in a flower together is termed calyx

Social/sociality: While most bee species are solitary, honeybees, bumblebees and stingless bees live in complex societies with clear tasks for queens, workers and males. These three bee groups are also termed as eusocial, which is the most evolved form of sociality expressing features like the simultaneous existence of different generations in one colony and the existence of non-reproductive castes which are doing brood care. Social organisation can also vary within some bee species. For example, some species of sweat bees (Halictidae) express solitary or social behaviour in different environments or at different live stages. Another type of social system can emerge when several females of the same generation join to initiate a new colony. Such behaviours in forming social organisations are termed semi-social

Solitary: Most bee species live solitary, meaning that each female builds its own nest. There is no division of labor and no different castes. A female bee searches for a nesting place and starts to build brood cells for example with leaves in a reed internode, collect pollen as larval food and lay an egg in each brood cell. After closing a nest the female leaves their progeny to their own devices

Stamen: Male organ of the flower consisting of the filament and anther. It is also termed androecium

Staminode: Sterile usually rudimentary stamen

Stigma and style: The female part of a flower. The stigma is the top part of the style, where deposited pollen grains can grow into pollen tubes when they are compatible with the flower

Stigma: The sticky end of the style responsible to catch pollen and the place in the flower where pollen germination starts

Style: A thin tube-like structure holding the stigma. It is connected with the ovary at its base

Superior ovary: Ovary is attached to the receptacle above the attachment of other flower parts

Tepals: Modified leaves surrounding the reproductive parts of a flower and comprises petals and sepals. The term is only used when petals and sepals cannot be easily separated. Taking all tepals in a flower together is termed perigonium

Wind pollination: Reproduction with airborne pollen transferred by wind

Xenogamy: Flower sets fruits and seeds with pollen of another flower of another plant individual. This needs a pollination agent such as a pollinator or wind

Zygomorph: Flower symmetry is bilateral

States of Brazil

State	abbreviation	
Acre	AC	3
Alagoas	AL	RR
Amapá	AP	S AP
Amazonas	AM	
Bahia	BA	
Ceará	CE	AM
Distrito Federal	DF	PA MA CE RN
Espírito Santo	ES	E PR
Goiás	GO	PI PI PI
Maranhão	MA	AC
Mato Grosso	МТ	RO BA SE
Mato Grosso do Sul	MS	MT
Minas Gerais	MG	DE
Pará	PA	
Paraíba	PB	
Paraná	PR	Mu
Pernambuco	PE	MS
Piauí	PI	SP
Rio de Janeiro	RJ	RJ
Rio Grande do Norte	ŔŇ	PR
Rio Grande do Sul	RS	SC
Rondônia	RO	
Roraima	RR	RS 🛃
Santa Catarina	SC	
São Paulo	SP	
Sergipe	SE	
Tocantins	TO	

Crop pollination

Açaí palm/Açaí Euterpe oleracea Mart.

Status: Native

Arecacea

Growing areas: AP, PA, TO, MA, GO

Cultivation: Semi shade in light woodlands/ agroforestry systems or no shade in open plantations

Attractiveness score: 5

Pollinator dependency: High

Plant-mating system: Mixed-mating system (xenogamy and geitonogamy)

Pollination requirements

Highly dependent on insect pollination, especially beetles, honeybees and stingless bees. Wind contributes little to pollination.

Flowering and harvest periods

Flowering: January to May Harvest: September to December

Main varieties grown in Brazil

Açaí-roxo, Açaí chumbinho, Açaí tinga, Açaí-açu, Açaí-branco, Açaí-espada, Açaísangue-de-boi

Some varieties have a dark skin and purple juice while others have a dark-green skin and greenish juice

Other information

Açaí palm is pollinated by a diverse range of insects, including curculionid beetles, generalist bees, beetles, flies and wasps. It was shown that pollinator diversity increases fruit set. The exclosure of pollinators visiting only male or female flowers, for example ants, can increase pollination and fruit set. An extensive list of pollinators is given in Campbell et al. (2018).









Flower description

Monoecious palm species with female and male flowers on large inflorescences of the same plant individual. Asynchrony of male and female flowers can occur. Flower symmetry is radially symmetrical. \bigcirc flowers are purple to light brown with sepals and petals broadly triangular, with a trilocular ovary with three stigmas. \bigcirc flowers are purplish with sepals triangular to ovate and petals ovate purple to purplered. Male flowers bear six free stamens arranged on a short receptacle.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bees	Melipona fasciculata	Eusocial
Stingless bees	Melipona flavolineata	Eusocial
Stingless bees	Frieseomelitta longipes	Eusocial
Solitary bee	Anthophila sp.	Solitary
Other bees	Apidae	Eusocial/Social/Solitary
Flies	Diptera	Solitary
Wasps	Apocrita	Eusocial/Social/Solitary
Beetles	Coleoptera	Solitary
Ants	Formicidae	Eusocial

Further information: A recent study of Campbell et al. (2018) demonstrates that (1) the palm is pollinated by more than 100 different types of insects, (2) fruit production is 25% higher in areas with high diversity of pollinators; (3) extensively used mixed forest stands support insect diversity pollinating açaí palm and (4) bees pollinating açaí palm were more dependent on the mixed forest stands than other pollinating insects. It is therefore important to conserve extensively used mixed forest stands adjacent to the açaí palm production areas or to manage açaí palm in diverse agroforestry systems with low pesticide inputs.



The stingless bee *Frieseomelitta longipes* on a female açaí palm flower

References

Campbell (2018); Lamarão (2018); Nascimento (2008); Oliveira do (2000); Venturieri (2008)

Acerola/Acerola *Malpighia emarginata* DC.

Status: Cultivated, originates from the northern Neotropics

Growing areas: AM, PA, BA, GO, ES, MG, RJ, SP, PR, PE, PB, PI, CE

Cultivation: Shrubs in open plantations

Pollinator dependency: High

Cultivation: Shrubs in oper Attractiveness score: 3 Pollinator dependency: H Plant-mating system: (geitonogamy) and cross p occur Pollination requirements The flower can potentially array of insect flower of collecting bee species, especies pollination Self (geitonogamy) and cross pollination (xenogamy)

The flower can potentially be pollinated by an array of insect flower visitors, but only oilcollecting bee species, especially those of the genus Centris, find it very attractive and visit



the flowers regularly.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

C. F. Rehnborg, F. Haley Red Jumbo, Hawaiian Queen, J. H. Beaumont, Manoa Sweet, Maunawili, Tropical Ruby

Other information

Acerola pollination is highly dependent on oil-collecting bees. Despite high levels of natural pollination, only an average of 30 % flowers set fruits. As many oil-collecting bee species prefer to use pre-existing holes in dead trees and other species nest in the soil. The presence of dead trees and soil with open vegetation in the vicinity of acerola cultivation is beneficial.

Bees visiting acerola also visit many other wild plants during acerola bloom. Hence, the presence of these plants may benefit acerola pollination. A study from the savannah of Central Brazil highlighted the potential to cultivate acerola in savannah areas when natural ecosystems that provide nesting and flowering resources, promoting bee diversity, are available.





Flower description

Flowers are hermaphrodite and radially symmetrical with five pale to deep pink fringed petals. Each inflorescence with three to five flowers. \mathcal{Q} organ with three styles emerging from a fused ovary. \mathcal{J} organ with ten stamen, all with fertile anthers.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Oil-collecting bee	Centris varia	Solitary
Oil-collecting bee	Centris aenea	Solitary
Oil-collecting bee	Centris tarsata	Solitary
Oil-collecting bee	Centris analis	Solitary
Oil-collecting bee	Centris flavifrons	Solitary
Oil-collecting bee	Centris nitens	Solitary
Oil-collecting bee	Centris spilopoda	Solitary
Oil-collecting bee	Centris fuscata	Solitary
Oil-collecting bee	Centris sponsa	Solitary
Oil-collecting bee	Centris vittata	Solitary
Oil-collecting bee	Centris bicolor	Solitary
Oil-collecting bee	Centris inermis	Solitary
Oil-collecting bee	Centris scopipes	Solitary
Oil-collecting bee	Centris longimana	Solitary
Oil-collecting bee	Centris mocsaryi	Solitary
Oil-collecting bee	Centris denudans	Solitary
Oil-collecting bee	Centris trigonoides	Solitary
Oil-collecting bee	Epicharis affinis	Solitary
Oil-collecting bee	Epicharis albofasciata	Solitary
Oil-collecting bee	Epicharis analis	Solitary
Oil-collecting bee	Epicharis bicolor	Solitary
Oil-collecting bee	Epicharis cockerelli	Solitary
Carpenter bee	Xylocopa cearensis	Solitary

References

Calgaro (2012); Freitas et al. (1999); Guedes et al. (2011); Magalhães & Freitas (2013); Oliveira et al. (2013); Vilhena et al. (2012)
Anatto/Urucum Bixa orellana L.

Status: Native (pan tropical)

Growing areas: All states of Brazil

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: High

Plant-mating system: Primarily cross pollination (xenogamy), small degree of self pollination (autogamy, geitonogamy)

Pollination requirements

Annatto is mainly buzz pollinated by carpenter bees, bumblebees and stingless bees. Self pollination leads to low fruit set, confirming that cross pollination (xenogamy) is the main reproductive strategy.





Flowering and harvest periods

Flowering: year round Harvest: year round

Main varieties grown in Brazil

Bico-de-Pato, BR-36, BR-37, Peruana Pará, Peruana Paulista, Piave Vermelha

Other information

Large bees such as carpenter bees, orchid bees and bumblebees but also small bees such as stingless bees (especially Melipona sp.) perform buzz pollination, vibrating the anthers and are therefore excellent pollinators, increasing fruit set. Honeybees, soft-wing flower beetles and ants are frequent flower visitors but contribute only little to pollination as they do not buzz pollinate and rarely touch the elongated stigma. Fruits produced by self pollination were shown to be lighter with less seeds originating than fruits from cross pollination. We assume that a combination of different pollinator species able to buzz pollinate and to transfer large amounts of pollen is the best strategy to pollinate annatto.





Annato trees have hermaphrodite flowers and flowers are radially symmetrical with four to five sepals and four to seven petals that often tend to curve up. Flowers are coloured pinkish, whitish or purplish. \bigcirc flower organ composed of one long pistil with bristly one-celled, superior ovary and the style thickened upwards with a short, two-lobed stigma. \bigcirc flower organ with numerous violet stamens and anthers.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Bumblebee	Bombus morio	Social
Bumblebee	Bombus sp.	Social
Orchid bee	Centris sp.	Solitary
Orchid bee	Euglossa sp.	Solitary
Orchid bee	Eulaema nigrita	Solitary
Stingless bee	Melipona subnitida	Eusocial
Stingless bee	Schwarziana quadripunctata	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Tetragonisca sp.	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Carpenter bee	Xylocopa frontalis	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Beetle	Ästylus sp.	Solitary
Ants	Formicidae	Eusocial

The stingless bee *Melipona* subnitida on an anatto flower

References

Caro et al. (2017); Castro (2009); Costa et al. (2008); Fabriet al. (2008); Rocha & Polatto (2017)

Apple/Maçã Malus domestica Borkh.

Status: Cultivated, originates from Central Asia

Growing areas: BA, MG, SP, PR, RS, SC

Cultivation: Open plantations/orchards

Attractiveness score: 4

Pollinator dependency: High

Plant-mating system: Most varieties with primarily cross pollination (xenogamy) but small degree of self pollination (autogamy, geitonogamy) and also parthenocarpy occurs

Pollination requirements

Generally, apple flowers need cross pollen from compatible varieties, acting as pollinizers, to set fruits. Pollen is transferred by insects and not wind

because apple pollen is too heavy to become airborne. Flowers provide both nectar and

pollen to insect populations and are attractive for many insects, especially bees.

Flowering and harvest periods

Flowering: August to October, both flowering and harvest depend on the variety and state where it is cultivated. Flowering last for 15 days, flowers open for three to five days

Harvest: December to January

Main varieties grown in Brazil

Anna, Brasil, Catarina, Condessa, Eva, Fuji, Gala, Golden Delicious, Granny Smith

Other information

Apis mellifera is the main pollinator but flowers are also visited by many solitary bee and bumblebees. Pollinator species effectiveness of different bee species varies among apple varieties due to variation in flower morphology. For example, Apis mellifera often collect nectar from the side of the stigma of large flowers reducing the transfer of pollen. Small insects also often fail to touch the stigma to transfer pollen. Optimal pollination increases seed number to ten seeds and fruit calcium content, important for storage quality. Orchards with few pollinators risk a low quality harvest.









Apple are deciduous trees with hermaphrodite flowers and flowers are radially symmetrical with five petals and sepals. Petal colour changes from white to deep pink with successful pollination. The central flower of the inflorescence is termed king flower and opens first and can develop a larger fruit. \bigcirc organ composed of one carpel with five stigmas with styles united at the base. Each of the five ovaries usually carries two seeds. \bigcirc organ with about 20 stamens in three whorls (10 + 5 + 5) with yellow anthers.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Bumblebees Solitary bees Solitary bees

Scientific name	
Apis mellifera	
Schwarziana quadripunctata	
Bombus sp.	
Augochloropsis sp.	
Dialictus sp.	

Sociality Eusocial Eusocial Social Solitary Solitary



Wasp on an apple flower



Honeybee (Apis mellifera) on an apple flower

References

Benedek & Nyeki (1997); Blitzer et al. (2016); Boyle & Philogène (1983); DeGrandi-Hoffman et al. (1985); Free (1962); Free (1966a); Free (1966b); Free & Spencer-Booth (1964); Freitas (1995); Goodell & Thomson (1997); Hem & Mattu (2014); Kendall & Solomon (1973); Kuhn & Ambrose (1984); Lewis & Smith (1969); Mayer (1984); Nunes-Silva et al. (2016); Park et al. (2016); Salomé (2014); Schneider et al. (2002); Smith & Lewis (1972); Stern et al. (2001); Storhaug (2014); Torchio (1985); Viana et al. (2015); Vicens & Bosh (2000)

Avocado/Abacate Persea americana Mill.

Status: Naturalized, originates from Mexico

Growing areas: AL, BA, CE, PB, PE, RN, SE, ES, MG, RJ, SP, PR, RS, SC

Cultivation: Mainly in open plantations

Attractiveness score: 4

Pollinator dependency: High

Plant-mating system: Cross pollination (xenogamy) always enhances fruit production. Self pollination is technically not possible as stigmas are not receptive at day times when pollen is released

Pollination requirements

Flowers provide nectar and pollen and are attractive to bees, flies and even bats are sometimes visiting flowers. All varieties have flowers with a protogynous dichogamy





pollination mechanism with two flowering types of different varieties: flowers open as female in the morning of the first day and close in late morning or early afternoon. The next day they open as male in the afternoon. Alternatively, they open as female in the afternoon of the first day, close in late afternoon, and reopen as male the next morning.

Flowering and harvest periods

Flowering: year round in the overall country, but few weeks in each region

Harvest: year round in the overall country, but few weeks in each region

Main varieties grown in Brazil

Antilhano, Bertanha, Breda, Emor, Fortuna, Fuerte, Geada, Guatemalense, Hass, Herculano, Linda, Margarida, Ouro Verde, Paulista, Pollock, Princesa, Quintal, Simmonds Solano Wagner, Waldin, Westin

Other information

At least two cultivars of different dichogamy variations should be planted less than 100 m apart from each other to ensure optimal pollen transfer and pollination success. Flowers contain nectaries with additional sugary rewards for pollinators.



Flowers of the perennial tree are protogynous hermaphrodite. Each flower is radially symmetrical and inconspicuous, small (5-10mm) and greenish-yellow with two perianth whorls with six tepals altogether. \bigcirc flower organ is hairy and consists of one unicarpellate ovary and the stylus carries a slightly lobed stigma \eth flower organ composed of 12 stamens in four whorls.

Flower visitors and pollinators

Common name	Scientific name	Sociality	
Honeybee	Apis mellifera	Eusocial	
Stingless bee	Trigona fulviventris	Eusocial	
Bumblebee	Bombus sp.	Social	
Other bees	Exomalopsis sp.	Solitary	71
Blow fly	Chrysomya megacephala	Solitary	
Wasp	Polistes canadensis	Social	

Table 3: Avocado flower opening sequence for flowering types A and B. For example Hass is a variety of flower type A, Forte a variety of flower type B they can be planted together to ensure optimal cross pollination when honeybees or other bees are available.

Flower type/	Day 1	Day 1	Day 2	Day 2
variety	Morning	Afternoon	Morning	Afternoon
Α	female	closed	closed	male
В	closed	female	male	closed



Avocado fruit is a mass flowering crop, producing millions of flowers, most of which are shed without producing fruits. Because of the many flowers, the crop is attractive for social insects and the flower-visiting community usually is by honeybees, dominated which are effective avocado pollinators for plantations.

References

Clark (1923); Davenport (1986); Ish-Am et al. (1998, 1999); Ish-Am & Eisikowitch (1993, 1998); Papademetriou (1976); Perez-Balam et al. (2012); Read et al. (2017); Vithanage (1990)

Brazil nut/Castanha do Brasil *Bertholletia excelsa* Bonpl.

Status: Native

Growing areas: AC, AM, AP, PA, RO, RR, MT **Cultivation:** Open plantations

Cultivation. Open plantation

Attractiveness score: 4

Pollinator dependency: Essential

Plant-mating system: Predominantly cross pollination (xenogamy)

Pollination requirements

Where cultivated, brazil nut is a mellitophilous cross-pollinated crop. It is considered selfincompatible, but some fruits result from geitonogamy.







Flowering and harvest periods

Flowering: September to December Harvest: January to April

Main varieties grown in Brazil

Abufari, 609, 606

Other information

The brazil nut is a plant species still under the process of cultivation. Most of the nuts are collected from wild trees and much less from planted tree individuals. Only large bees, especially orchid bees, are strong enough to access the flower and pollinate. These bees are usually solitary wild bees nesting in natural habitats, therefore is the proximity to rainforest important to ensure the presence of these pollinators in adequate numbers and diversity of species for sufficient pollination.

Brazil nut is an example showing how important it is to conserve rainforest in Brazil as we are currently not able to manage the large wild bee species pollinating brazil nut flowers. Most of the large bees depend on resources of intact rainforest. In a nutshell "No rainforest, no brazil nut".





Flowers are hermaphrodite and bilaterally symmetrical with a globous appearance and six yellow-cream petals over a thick receptacle. The androecium includes a hood of sterile, nectar producing appendages that cover a ring of fertile stamens beneath. \mathcal{Q} organ with inferior ovary and short styles. \mathcal{J} organ with short and numerous stamens.

Flower visitors and pollinators

Common name	Scientific name	Sociality	
Honeybee	Apis mellifera	Eusocial	
Stingless bee	Frieseomelitta longipes	Eusocial	
Stingless bee	Melipona lateralis	Eusocial	
Bumblebee	Bombus brevivillus	Social	
Bumblebee	Bombus transversalis	Social	
Oil-collecting bee	Epicharis conica	Solitary	
Oil-collecting bee	Ēpicharis umbraculata	Solitary	
Oil-collecting bee	Ēpicharis zonata	Solitary	
Oil-collecting bee	Ĉentris americana	Solitary	
Oil-collecting bee	Centris carrikeri	Solitary	
Oil-collecting bee	Centris ferruginea	Solitary	
Oil-collecting bee	Centris denudans	Solitary	
Orchid bee	Eulaema meriana	Solitary	
Orchid bee	Eulaema mocsaryi	Solitary	
Orchid bee	Eulaema cingulata	Solitary	
Orchid bee	Eufrisea purpurata	Solitary	
Orchid bee	Eufrisea flaviventris	Solitary	
Carpenter bee	Xylocopa frontalis	Solitary	
Leafcutter bee	<i>Megachile</i> sp.	Solitary	

The orchid bee *Eulaema meriana* collecting pollen and nectar from a brazil nut flower

References

Cavalcante (2008); Cavalcante et al. (2012); Cavalcante et al. (2018); Maués et al. (2015); Ministério da Agricultura, Pecuária e Abastecimento (2014); Nelson et al. (1985); Prance (1976); Santos & Absy (2010); Santos & Absy (2012)

Brazilian grapetree/Jabuticaba Plinia cauliflora (Mart.) Kausel

Status: Native

Growing areas: PR, SC, MG, RJ, ES, BA, PE, SP, PΒ

Cultivation: Open plantations

Plant-mating system: Mixed-mating system (autogamy, geitonogamy, xenogamy)

Pollinator dependency: Little

(autogamy, geitonogamy, xe Attractiveness score: 3 Pollinator dependency: Li Pollination requirements It has been reported that se It has been reported that solitary trees bear poorly compared to trees planted in groups, which indicates that cross pollination may enhance productivity. Pollinator exclusion







experiments showed only slight increases in fruit set of cross-pollinated compared to self-pollinated flowers.

Flowering and harvest periods

Flowering: July to August

Harvest: September to November

Main varieties grown in Brazil

Branca, Mineira, Paulista, Ponhema, Rajada, Roxa, Rujada, Sabará, São Paulo

Other information

Flowers of the slow-growing evergreen tree are visited mainly by bees, especially honeybees. Other insects such as flies and wasps also visit the flowers. The crop can produce commercially without pollinators, but flowers are visited by many different beneficial insects. Therefore, if applied, spraying of pesticides, especially insecticides, has to be avoided when the crop is in bloom. When the brazilian grapetree is planted in a garden it can be a valuable pollen and nectar source for insects at a certain time. But also gardeners should avoid pesticides at least when the tree is in bloom. As the fruits are consumed by wild birds this crop can promote wildlife in private gardens or public parks.





The cauliflorous flowers are hermaphrodite and radially symmetrical with small, white petals. \bigcirc flower organ consists of a single compound pistil with a single style and stigma. \bigcirc flower organ consists of numerous stamens.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Stingless bee Other bees Wasps Flies

- Scientific name Apis mellifera Tetragonisca angustula Trigona spinipes Chloralictus sp. Apocrita Diptera
- Sociality Eusocial Eusocial Solitary Solitary/Social Solitary



A solitary bee on a brazilian grapetree flower

References

Gobato et al. (2018); Gressler et al. (2006); Malerbo et al. (1991); Malerbo-Souza et al. (2004); Morton (1987)

Canola/Canola Brassica napus L.

Status: Cultivated, originates from Europe

Growing areas: RS, PR, MG, SC, MT

Cultivation: Open fields

Attractiveness score: 5

Pollinator dependency: Modest

Plant-mating system: Dependent on the variety mostly self pollination (autogamy, geitonogamy) but cross pollination (xenogamy) can occur

Pollination requirements

Canola is considered a predominantly selfpollinated and self-compatible plant. Pollen transport by insects (mainly bees) or wind can lead to higher seed set, seed quality and yield. Although

canola flowers remain open for 12 to 96 hours, the period in which they can be fertilized





varies only from 4 to 24 hours.

Flowering and harvest periods

Flowering: July to September

Harvest: August to November

Main varieties grown in Brazil

CTC-4, Hyola 401, Hyola 411, Hyola 420, Hyola 43, Hyola 432, Hyola 433, Hyola 60, Hyola 61, Hyola 76, PFB-2

Other information

Autogamous self pollination through gravity is most likely during the onset of the anthesis, when the stigma is below the interiorly facing anthers and hence pollen falls easily on the stigma. Once the stigma exceeds the anthers self pollination (geitonogamy) and cross pollination through insects and wind is facilitated. The introduction of honeybee colonies to canola plantation can, depending on the canola variety, increase crop yield. Moreover, wild social and solitary bees have been shown to work in a complementary way increasing the pollination rates and productivity of canola seeds. As canola is pollinated by many wild bee species but fields provide no nesting resources, farmers should provide set aside patches with flowering herbs adjacent to and within large fields.





Flowers are hermaphrodite and radially symmetrical with four yellow petals alternating with four sepals in the typical crosswise arrangement. \bigcirc organ with one single carpel and a superior ovary. \eth organ with two lateral stamens with short filaments and 4 long stamens.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Plebeia emerina	Eusocial
Stingless bee	Plebeia droryana	Eusocial
Stingless bee	Plebeia nigriceps	Eusocial
Stingless bee	Mourella caerulea	Eusocial
Stingless bee	Nannotrigona testaceicornis	Eusocial
Stingless bee	Scaptotrigona bipunctata	Eusocial
Stingless bee	Schwarziana quadripunctata	Eusocial
Stingless bee	Tetragonisca fiebrigi	Eusocial
Bumblebee	Bombus sp.	Social
Bumblebee	Bombus pauloensis	Social
Carpenter bee	Xylocopa sp.	Solitary
Leafcutter bee	Megachile sp.	Solitary
Mining bee	Callonychium petuniae	Solitary
Small carpenter bees	Ceratina sp.	Solitary
Sweat bee	Lasioglossum phaedrum	Solitary/Social
Sweat bee	Thectochlora alaris	Solitary/Social
Sweat bee	Dialictus pabulator	Solitary/Social
Sweat bee	Dialictus sp.	Solitary/Social
List continues at p. 120	1	

References

Abrol & Shankar (2012); Adegas & Nogueira Couto (1992); Arthur et al. (2010); Benedek & Prenner (1972); Blochtein et al. (2015); Bommarco et al. (2012); Brunel et al. (1994); Chambó et al. (2014); Chifflet et al. (2011); Eisikowitch (1981); Free & Ferguson (1980, 1983); Free & Nuttall (1968); Hayter & Cresswell (2006); Holzschuh et al. (2011); Hoyle et al. (2007); Jauker et al. (2012); Jenkinson & Glynne Jones (1953); Kamel et al. (2015); Kevan & Eisikowitch (1990); Koltowski (2001a,b & 2002, 2005, 2007); Langridge & Goodman (1982); Marsaro-Jr. (2017); Mesquida et al. (1988); Mesquida & Renard (1979, 1981); Mesquida et al. (1988) List continues at p. 119

Cashew/Caju Anacardium occidentale L.

Status: Native

Growing areas: AC, AM, AP, PA, RR, TO, AL, BA, CE, MA, PB, PE, PI, RN, SE, DF, GO, MS,

Cultivation: Open plantations

Pollinator dependency: High

Self pollination (geitonogamy) and cross pollination (xenogamy)

Las: AC, 1 L, MA, PB, PE, LAT, ES, MG, RJ, SP **Cultivation:** Open plant **Attractiveness score:** 4 **Pollinator dependency: Plant-mating system** (geitonogamy) and cross pc **Pollination requirements** Insects transporting pollen of either male or effective poll: role in Insects transporting pollen from the large stamen of either male or hermaphrodite flowers are effective pollinators, whereas wind plays a minor

role in cashew pollination. Pollen from hermaphrodite flower staminoids are unfertile or







show very low germination and therefore pollination rates.

Flowering and harvest periods

Flowering: June to October

Harvest: August to December

Main varieties grown in Brazil

Dwarf varieties (BRS 189, BRS 226, BRS 253, BRS 265, CCP 06, CCP 09, CCP 1001, CCP 76, Embrapa 50, Embrapa 51), Giant variety (BRS 274)

Other information

Self pollination produces low yield while cross pollination results in high fruit set and harvest. Apis mellifera is the most common pollinator in cashew orchards but the presence of wild bees increases yield.

Oil-collecting bees of the genus Centris are pollinators of cashew flowers. good However, they only occur in areas where oilflower plants are present because they need their oil for nest building and cell provisioning. Cashew growers can attract them to their orchards intercropping cashew with an oil-flower species such as acerola (Malphigia emarginata) (Freitas & Pereira, 2004).





Flowers are male or hermaphrodite and appear radially symmetrical with five backwards bent, white or pinkish petals. Male flowers are more numerous. \mathcal{Q} organ with a hairless ovoid ovary and a long lateral style. \mathcal{J} organ with a corona of stamens surrounding the ovary, which is rudimentary in male flowers. One stamen is exerted and nine are short and inserted.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Stingless bee	Melipona subnitida	Eusocial
Stingless bee	Scaptotrigona sp.	Eusocial
Stingless bee	Trigona sp.	Eusocial
Bumblebee	Bombus sp.	Social
Oil-collecting bee	Centris analis	Solitary
Oil-collecting bee	Centris flavifrons	Solitary
Oil-collecting bee	Centris fuscata	Solitary
Oil-collecting bee	Centris sponsa	Solitary
Oil-collecting bee	Centris tarsata	Solitary
Cleptoparasitic bee	Coelioxys sp.	Solitary
Orchid bee	Euglossa sp.	Solitary
Leafcutter bee	Megachile sp.	Solitary
Carpenter bee	Xylocopa grisescens	Solitary
Carpenter bee	Xylocopa cearensis	Solitary
Other bee	Exomalopsis sp.	Solitary
Wasp	Polistes sp.	Social
Ant	Camponotus sp.	Social

Further information: The management of honeybee colonies in cashew plantations during blossom increases flower visits and optimizes fruit and nuts harvest. Additionally cashew provides abundant resources for honeybees allowing income through the harvest of valuable bee products like honey, beeswax and propolis.

References

Bhattacharya (2004); Eradasappa & Mohana (2016); Freitas (2018); Freitas et al. (2002, 2014a,b); Freitas & Paxton (1996, 1998); Freitas & Pereira (2004); Heard et al. (1990); Holanda Neto et al. (2002)

Castor bean/Mamona Ricinus communis L.

Status: Naturalized, originates from Eastern Africa to India

Growing areas: BA, CE, MG, PR, PE, PI, RN, SP

Cultivation: Open fields

Attractiveness score: 2

Pollinator dependency: Modest

Plant-mating system: Cross- and self pollination (mixed-mating system), male flowers are below female flowers hence self pollination (geitonogamy) and cross pollination (xenogamy) occurs, of which the first was shown to increase reproductive success.

Pollination requirements

Castor flowers are cross-pollinated mainly by wind, whereas bees contribute to self





promote pollination. Honevbees geitonogamy in two ways: i) by actively transporting pollen from male flowers to female flowers of the same inflorescence when they search for nectar at extrafloral nectaries located at the base of each pedicel; ii) passively by activating the explosive dehiscence of pollen from the anthers during foraging activities. This increases the airborne amount of pollen mostly pollinating the stigmas of female flowers on the same or neighbouring inflorescence.

Flowering and harvest periods

Flowering period: December to July

Harvest period: February to Octber

Main varieties grown in Brazil

Al Guarany 2002, BRS 149 Nordestina, BRS 188 Sertaneja (Paraguaçu), BRS Energia, IAC 2028, IAC 226, IAC 80, IAC Guarani

Other information

The introduction of honeybee colonies in castor bean plantations increases crop fruit set per inflorescence and seed oil content.





Flowers are monoecious, with radially symmetrical male and female flowers on one plant with three to five sepals. Ovoid male flowers yellowish-green with creamy stamens. \mathcal{Q} flower with superior ovary covered by soft spines and a short style with three prominent red, deeply two-parted stigmatic branches. \mathcal{J} flower with many yellowish stamens, branched and tightly packed together in a perianth. Premature flowers are green and ovoid.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Scaptotrigona bipunctata	Eusocial



Stingless bees (Scaptotrigona sp.) on castor bean flowers



Honeybee (Apis mellifera) on castor bean flowers

References

Giannini et al. (2015); Rizzardo (2007); Rizzardo et al. (2012)

Citrus, Lime, Orange/Frutas cítricas Citrus spp. L.

Status: Naturalized, originate from Southeast Asia. **Growing areas:** BA, DF, GO, MS, MG, SP, PR, RS, SC

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: Little

Plant-mating system:

Citrus is an aggregated group of more than ten species grown in Brazil with many varieties and including all mating systems ranging from agamospermy and parthenocarpy (seedless mandarins) to all forms of self- and cross pollination

Pollination requirements

Many varieties of grapefruit, lemon, lime and oranges are agamosperm or parthenocarpic



and do not require pollination to set fruits. However some varieties, especially pomelo and mandarins, require or benefit from entomophilous cross pollination to set fruits or to improve yield, juice content or sweetness. In general citrus plants are attractive to pollinators.

Flowering and harvest periods

Flowering: September to November

Harvest: February to June

Main varieties grown in Brazil

Oranges: Baianinha (Piralima), Charmute de Brotas, Hamlin, João Nunes, Natal, Pera Rio, Pineapple, Rubi, Seleta, Valência, Valência Americana1, Valência Folha Murcha, Westin. List continues at p. 119

Other information

In some orange varieties, like 'Pera Rio', fruit set increases when pollinators have access to the flowers and flowers open to pollinators even bear more fruits than handcross-pollinated flowers. Pollination also increases seed numbers, depending on how the fruits will be used, this can decrease their market value. In the same variety yield increases up to 30% with honeybee visitation.





Flowers are mainly hermaphrodite and radially symmetrical. The five petals are white, thick and leathery. \bigcirc flower organ consists of a superior ovary, one single cylindrical style and a big, round stigma. \eth flower organ with numerous stamens with yellow quadrilobate anthers and a short, rudimentary pistil.

Flower visitors and pollinators

Common name Honeybee Stingless bee Stingless bee Stingless bee Carpenter bee Carpenter bee Small carpenter bee Oil-collecting bee Sweat bee Masked bee Leafcutter bee

Scientific name Apis mellifera Melipona quadrifasciata Melipona scutellaris Tetragonisca angustula Frieseomelitta sp. Xylocopa grisescens Xylocopa suspecta Ceratina sp. Centris analis Centris fuscata Augochlora sp. Hylaeus sp. Megachile sp.

Sociality Eusocial Eusocial Eusocial Eusocial Solitary Solitary



Honeybee (Apis mellifera) on a lemon flower

References

Grajales-Conesa et al. (2013); Malerbo-Souza et al. (2004); Ribeiro et al. (2017); Toledo et al. (2013)

Cocoa/Cacau Theobroma cacao L.

Status: Native

Growing areas: AC, AM, AP, PA, RO, BA, MA, RR

Cultivation: Agroforest

Attractiveness score: 4

Pollinator dependency: Essential

Plant-mating system: mostly cross pollination (xenogamy)

Pollination requirements

Most varieties of cocoa are self-incompatible and self pollination produces no fruit set. Naturally, midges (Diptera) are the most important pollinators. Hand-cross pollination produces very high fruit set and fruits can be five times heavier



than open-pollinated flowers, which suggests a pollination deficit in cultivation areas.



Flowering and harvest periods

Flowering: October to May Harvest: April to August

Main varieties grown in Brazil

Criollo, Forastero, Trinitário

Other information

Many insects live on and visit the flowers of cocoa, but most do not contribute to pollination. Usually natural fruit set is only 0,3% (3 in every 1,000 flowers). Small-sized stingless bees were tested as potential cocoa pollinators, but despite the evidence that they visit the flowers their efficiency as pollinators is still to be proven. When hand pollination is considered, it should be avoided to set the maximum number of fruits in a given year because this weakens the tree, which may die or become too weak for a good yield in the following year. The midges main pollinating insects are (Ceratopogonidae, also known as biting midges or gnats) which need moisture to reproduce. Moist microhabitats can be promoted by leaving decaying organic materials in the agroforestry systems. Insect control of nuisance biting midges should be reconsidered as their removal might lead to pollination failure or a low quality harvest.





Flowers are hermaphrodite, cauliflorous and radially symmetrical with five triangular, whitish or reddish sepals and five whitish-yellow petals with dark purple bands, fused into a cup-like structure at the base. \bigcirc organ with superior ovary and a single style terminating in five sticky lobes. \bigcirc organ with five fertile stamens fused, each with two anthers, alternating with five staminodes. The two whorls together form a tube.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Biting midge	Forcipomyia nana	Solitary
Biting midge	Culicoides fluviatilis	Solitary
Biting midge	Forcipomyia jipajapae	Solitary
Biting midge	Atrichopogon fusculus	Solitary
Biting midge	Culicoides diabolicus	Solitary
Biting midge	Culicoides glabellus	Solitary
Biting midge	Culicoides hylas	Solitary
Biting midge	Culicoides paraensis	Solitary
Biting midge	Culicoides pusillus	Solitary
Biting midge	Fittkauhelea amazonica	Solitary
Biting midge	Forcipomyia argenteola	Solitary
Biting midge	Forcipomyia blantoni	Solitary
Biting midge	Forcipomyia cinctipes	Solitary
Biting midge	Forcipomyia genualis	Solitary
Biting midge	Forcipomyia quatei	Solitary
Biting midge	Forcipomyia spatulifera	Solitary
Biting midge	Forcipomyia squamitibia	Solitary
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Plebeia cf. flavocincta	Eusocial



Biting midge on a cocoa flower

References

Adjaloo & Oduro (2013); Chumacero de Schawe et al. (2016); Erickson et al. (1988); Frimpong et al. (2009); Groeneveld et al. (2010); Lemos (2014); Lopes et al. (2011); Santarém & Felippe-Bauer (2016); Silva et al. (2011); Soria (1981); Young (1982); Young (1983)

Coconut/Coco Cocos nucifera L.

Status: Naturalized in Brazil. Pan-tropical but likely originated in South Asia and Melanesia

Growing areas: PA, AL, BA, CE, MA, PB, PE, PI, RN, SE, ES, RJ, SP

Cultivation: Mainly in open plantations or single trees in gardens or along streets or in agroforests

Attractiveness score: 4

Mixed-mating system

Pollinator dependency: Modest
Plant-mating system: Mixed (geitono- and xenogamy)
Pollination requirements
Dioescious flowers are present inflorescence and male flowers female flowers. Wind pollination of the second sec Dioescious flowers are present in the same inflorescence and male flowers mature before female flowers. Wind pollination can facilitate self- and cross pollination. Flowers are also







attractive for insects and especially in windless areas bees facilitate self- and cross pollination and this was shown to increase production.

Flowering and harvest periods

Flowering: year round

Harvest: year round

Main varieties grown in Brazil

Giant cultivars: Gigante-da-Costa-Oeste, Gigante-da-Malásia, Gigante-da-Praia-do-Forte, Gigante-de-Renell, Gigante-do-Anão-Oeste-Africano; dwarf cultivars: Malásia, Anão-amarelo amarelo da de Gramame, Anão-verde-de-Jequi, Anãoverde-do-Brasil, Anão-vermelho da Malásia, Anão-vermelho de Camarões. Anãovermelho de Gramame; hybrid cultivars: PB 111, PB 121, PB 131, PB 141

Other information

Male flowers are highly attractive as pollen source and female flowers as nectar source for honeybees. Hand- cross pollination increases fruit set over self pollination and insect-mediated pollination increases yield over wind pollination. Wasps visit the female flowers for nectar but carry little amounts of pollen. Studies from outside of Brazil reported weevils as important pollinators.







Coconut trees are monoecious with radially symmetrical female and male flowers occurring in the same inflorescences. Flowers are small, light yellow and the perianth usually consists of two whorls of three rudimentary petals and sepals. Male flowers are more numerous than female flowers. \mathcal{Q} flower with three pistils. \mathcal{J} flower with six stamens consisting of two whorls with three stamens each.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Stingless bee Stingless bees Sweat bee Scientific name Apis mellifera Melipona quadrifasciata Oxytrigona tataira Trigona sp. Augochlora sp.

Sociality Eusocial Eusocial Eusocial Eusocial Solitary/Social



Honeybee (Apis mellifera) on coconut flowers



Stingless bee on coconut flowers

References

Benassi et al. (2013); Conceição et al. (2004); De Castro M.S. (2002); Free et al. (1975); Hedström (1986); Melendez-Ramirez et al. (2004); Regi & Josephrajkumar (2013)

Coffee/Café Coffea arabica L.

Status: Cultivated, originates from highlands of Ethiopia

Growing areas: AC, AL, BA, CE, DF, ES, GO, MG, MS, PB, PE, PR, RJ, RO, RS, SC, SE, SP

Cultivation: Open plantations, agroforests

Attractiveness score: 4

Pollinator dependency: Modest

Rubiaceae Plant-mating system: Self pollination (autogamy, geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Coffea arabica is self-compatible and flowers can auto pollinate under certain circumstances. Wind pollination occurs, but self- and cross pollination mediated by bees significantly increases fruit set.



Flower: August to October Harvest: April to May

Main varieties grown in Brazil

Acaiá, Acauã, Catuaí, Mundo Novo

Other information

Coffea arabica is the coffee species most cultivated in Brazil (70% of crops). Another coffee species, C. canephora, known as Robusta, is also cultivated in the country (30%). This species is self-incompatible and depends on cross pollination by bees and wind.

For a long time it was thought that, only Robusta coffee needs bees for optimal pollination. Then several studies showed that common varieties of Arabica coffee in different places of the world, including South America produces more fruits when bees are visiting the coffee flowers. It was also shown that social bees contribute effectively to coffee production but many of the wild social bees depend on resources of rainforest. Therefore, coffee intact production can be promoted by the vicinity of intact rainforest. D.W Roubik mentioned in a personal communication that the taste of coffee can even be better when bees are pollinating the coffee flowers.









Flowers are hermaphrodite and radially symmetrical with five sepals and five white petals forming a star-shaped corolla tube with a sweet scent. \bigcirc organ with inferior ovaries and two united unilocular carpels. The slender style terminates in short, two-parted stigmas. \eth organ with five stamens inserted in the corolla tube and anthers on long, slender filaments.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Bumblebee	Bombus brevivillus	Social
Bumblebee	Bombus morio	Social
Stingless bee	Cephalotrigona capitata	Eusocial
Stingless bee	Geotrigona subterranea	Eusocial
Stingless bee	Nannotrigona testaceicornis	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Trigona amalthea	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Oil-collecting bee	Centris aenea	Solitary
Oil-collecting bee	Centris decolorata	Solitary
Oil-collecting bee	Centris flavifrons	Solitary
Oil-collecting bee	Centris tarsata	Solitary
Small carpenter bee	Ceratina chloris	Solitary
Other bee	Exomalopsis iridipennis	Solitary
Carpenter bee	Xylocopa grisescens	Solitary
Stingless bee	Paratrigona sp.	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Stingless bee	Trigona sp.	Eusocial
Sweat bee	Augochlora sp.	Solitary/Social
Sweat bee	Augochloropsis sp.	Solitary/Social
Oil-collecting bee	Centris sp.	Solitary
Small carpenter bee	Ceratina sp.	Solitary
Carpenter bee	Xylocopa sp.	Solitary

References

Badano & Vergara (2011); Brokaw (2013); Hipolito et al. (2018); Hutchinson (2012); Klein et al. (2003 a,b); Mesquita et al. (2016); Philpott et al. (2006); Roubik (2002); Tarno et al. (2018); Veddeler et al. (2008); Vergara & Badano (2009)

Common bean/Feijão Phaseolus vulgaris L.

Status: Cultivated, originated from Mesoamerica

Growing areas: All states of Brazil

Cultivation: Open plantations and gardens

Attractiveness score: 4

Pollinator dependency: Little

Plant-mating system: Most varieties with primarily self pollination (autogamy) and a small degree of cross pollination (xenogamy)

Pollination requirements

Flowers of cultivated forms are monoecious and self-pollinated. Cross pollination by insects, particularly large-bodied bumblebees and carpenter bees increases yield and seed quality.



Flowering and harvest periods

Flowering: Year round Harvest: 80 to 100 days after planting

Main varieties grown in Brazil

BGF13013, BRS Agreste, BRS Ametista, BRS Ártico, BRS Campeiro, BRS Embaixador, BRS Estilo, BRS FC402, BRS Notável, BRS Pitanga, BRS Pontal, BRS Realce, BRS Requinte BRS Vereda, BRSMG Madreperola, BRSMG União, Jalo Precoce, Pérola

Other information

Insect pollination usually increases the number of seeds per pod. The number of pods depends on each variety and mostly on its auto pollination rate.

The morphology of the flower fits better to large bees with long tongues than to small bees with short tongues. Therefore, farmers should promote large bees with long tongues to optimize the pollination of the common bean. These can be carpenter bees, orchid bees, leafcutter bees or long-horned bees. Leafcutter bees can be promoted by the use of bee houses, see picture on page 113.





Flowers are hermaphrodite and bilaterally symmetrical. The tubular calyx is green and hairy. The corolla is white or pinkish white, with five petals, the posterior petal is outermost, the two lateral petals with long wings and the two anterior petals form a united keel. \bigcirc organ with one carpel. The ovary is superior, elongated, green, unilocular with numerous ovules. The style is long, bent at its base and terminates with the stigma. \eth organ with nine stamens, enclosed within the keel and fused into a tube, one posterior stamen is free.

Flower visitors and pollinators

Common name

Honeybee Bumblebee Carpenter bee Long-horned bee Leafcutter bee

Scientific name Apis mellifera Bombus atratus Xylocopa sp. Thygater analis Megachile sp.

Sociality Eusocial Social Solitary Solitary Solitary



The carpenter bee *Xylocopa grisescens* on a common bean flower

References

Carpentieri-Pípolo et al. (2001); Free (1966); Ibarra-Perez et al. (1999); Shree et al. (1991)

Cotton/Algodão Gossypium hirsutum L.

Status: Naturalized, originates from northern Neotropics

Growing areas: AL, AM, BA, CE, ES, GO, MA, MG, MS, MT, PB, PE, PI, PR, RJ, RN, SE, SP, TO

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: Modest

Plant-mating system: Commonly self pollination (autogamy), but a mixed-mating system with geitonogamy and xenogamy occurs

Pollination requirements

The hermaphrodite flowers of cultivated varieties are usually self-pollinating, additional cross pollination by bees increases cotton production. Bumblebees and carpenter bees are







considerably more effective pollinators, though less frequent in flowers than other bee species.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

BRS 187, BRS 336, BRS 368RF, BRS Topázio, BRS Verde, IMA6501B2RF, IMA7201B2RF, IMA7501WS, TMG 11 WS, TMG 41 WS, TMG 42 WS, TMG 43 WS, TMG 81 WS, TMG 82 WS

Other information

Productivity of some varieties increases by approximately 20% when pollinated by bees. Pollination by honeybees resulted in more cotton bolls than passive self pollination. The cotton bolls of open-pollinated flowers were 1.2 times heavier than cotton bolls forming after wind pollination. The cotton bolls of hand cross-pollinated flowers were 1.3 times heavier than cotton bolls forming after passive self pollination.





Flowers are hermaphrodite and radially symmetrical with three epicalyx bracts, five fused sepals and five free white petals. \bigcirc organ with three to five carpels fused into a superior ovary. The undivided style terminates in three to five stigmas. \bigcirc organ with 13 or more stamens fused into a tube around the pistil.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Paratrigona lineata	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Geotrigona mombuca	Eusocial
Stingless bee	Partamona cupira	Eusocial
Stingless bee	Partamona mulata	Eusocial
Stingless bee	Melipona quinquefasciata	Eusocial
Stingless bee	Tetragona clavipes	Eusocial
Stingless bee	Trigona hyalinata	Eusocial
Stingless bee	Frieseomelitta doederleini	Eusocial
Stingless bee	Trigona fuscipennis	Eusocial
Stingless bee	Frieseomelitta varia	Eusocial
Stingless bee	Trigona recursa	Eusocial
Stingless bee	Schwarziana quadripunctata	Eusocial
Stingless bee	Trigonisca sp.	Eusocial
Bumblebee	Bombus morio	Social
Bumblebee	Bombus atratus	Social
Carpenter bee	Xylocopa suspecta	Solitary
Carpenter bee	Xylocopa hirsutissima	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Small carpenter bee	Čeratina gossypii	Solitary
Small carpenter bee	Ceratina sp.	Solitary
Long-horned bee	Melissoptila cnecomala	Solitary
Long-horned bee	Melissoptila pubescens	Solitary
Long-horned bee	Florilegus festivus	Solitary
Oil-collecting bee	Centris scopipes	Solitary
Oil-collecting bee	Centris collaris	Solitary
Oil-collecting bee	Epicharis bicolor	Solitary
List continues at p. 119		-

References

Bozbek et al. (2008); Cusser et al. (2016); Eisikowitch & Loper (1984); FAO (2018); Heuberger et al. (2010); McGregor (1959); Moffett (1977); Moffett et al. (1975); Moffett et al. (1980); Pires et al. (2014, 2015); Rhodes (2002); Waller et al. (1985a,b)

Cowpea/Feijão-caupi Vigna unguiculata (L.) Walp.

Status: Cultivated, was first domesticated in Ghana

Growing areas: CE, PB, PI, PE, RN, MS, MT, MG, RJ, SP, PR, AL, SE, BA

Cultivation: Open fields

Attractiveness score: 3

Pollinator dependency: Little

Plant-mating system: Cleistogamous, self pollination (autogamy) and cross pollination (xenogamy)

Pollination requirements

Cowpea flowers produce nectar, which is attractive for insects, especially bees. However, flowers are usually auto-pollinated, before they open. Insect flower visitors increase pollen



STATION OF THE STATE

deposition, especially on flowers that failed to self pollinate and therefore increase seed set per pod and number of pods.

Flowering and harvest periods

Flowering: Year round

Harvest: 60 to 90 days after planting

Main varieties grown in Brazil

Arigozinho, BR 17 - Gurgueia, BR3 -Tracuateua, Branco de praia, BRS Amapá, BRS Aracê, BRS Cauamé, BRS Guariba, BRS Itaim, BRS Juruá, BRS Marataoã, BRS Mazagão, BRS Milênio BRS Novaera, BRS Pajeú, BRS Paraguacu, BRS Rouxinol, Potengi, BRS BRS Tumucumaque, BRS Urubuquara, BRS Corujinha, Xiquexique, Manteiguinha, Manteiguinha roxo, Mudubim de Rama, Preto de praia, Quarentão, Roxinho de praia

Other information

Seed number, dry weight of seed/pod, pod length and the percentage of normal form seed/pod decreases with insect exclusion from flowers. Insect exclusion decreases fruit production rate from 62% to 48% and the proportion of healthy seeds from 98% to 76%.





Flowers are hermaphrodite and zygomorphic with five free sepals and petals, the upper petal is outermost, the two side petals have long wings and the two lower petals form a united keel. \bigcirc organ with multiple carpels fused and an insertion in the terminal style. \bigcirc organ with one out of nine stamen longer than the others and with free anthers.

Flower visitors and pollinators

Common name

Honevbee
Stingless bee
Stingless bee
D 111
Bumblebee
Carpenter bee
Carpenter bee
Leafcutter bee
Beetle

Scientific name Apis mellifera Melipona subnitida Trigona fuscipennis Bombus brevivillus Xylocopa cearensis Xylocopa grisescens Megachile sp. Lagria villosa

Sociality Eusocial Eusocial Social Solitary Solitary Solitary Solitary



Stingless bees (Trigona fuscipennis) on a cowpea flower

References

Araújo (2012); Asiwe (2009); D'Andrea et al. (2007); Fohouo et al. (2009); Ige et al. (2011); Sheahan (2012); Venter (1996)

Cupuassu/Cupuaçu *Theobroma grandiflorum* (Willd. ex Spreng.) K.Schum.

Status: Native

Growing areas: AC, AM, PA, RO, BA Cultivation: Shade (Tropical forest) Attractiveness score: 2

Pollinator dependency: High

Plant-mating system: Cross pollination (xenogamy)

Pollination requirements

Self-incompatible and requires cross pollination by insects. The main effective pollinators are bees that move between trees, but ants and aphids are also considered to be pollinators.



Flower: July to September

Harvest: October to June

Main varieties grown in Brazil

Cupuaçu-mamau, Cupuaçu-mamorana, Cupuaçu-redondo

Other information

Hand pollination resulted in a much higher fruit set than open pollination, indicating that lack of effective pollination is a reason for low yields. Experimental pollination using compatible pollen grains have shown that a flower which receives 60 compatible pollen grains has 20% probability of setting fruit; a flower which receives more than 400 pollen grains always sets fruit. However, fewer than 2% of naturally-pollinated Cupuassu flowers have more than 50 pollen grains on their stigmas. Cupuassu varieties with many flowers, blooming for a long time per tree, might increase the attractiveness of the tree for pollinators and with this the chance that many and diverse pollinating available. In general, the species are pollination system of cupuassu is not well studied, but the more insect species visit the flowers the higher the fruit set should be. Therefore, conserving this beneficial insects should be valuable in commercial cupuassu production systems.











Flowers are hermaphrodite and radially symmetrical with a perianth of five thick fleshy, boat-shaped, whitish green and tomentose sepals and five thick, whitish petals terminating in purple tongues. \mathcal{Q} organ with five carpels fused into a superior, tomentose ovary, with a yellowish style. \mathcal{J} organ with ten stamens including five infertile, long, lanceolate and purple stamen and five fertile stamen with short three parted anthers.

Flower visitors and pollinators

Common name

- Stingless bee Stingless bee Stingless bee Stingless bee Ant Beetle Beetle
- Scientific name Plebeia minima Tetragonisca angustula Trigona pallens Aparatrigona impunctata Wasmannia sp. Baris sp. Chrysomelidae
- Sociality Eusocial Eusocial Eusocial Eusocial Solitary Solitary



Ant on a cupuassu flower



Stingless bee (Trigona sp.) in a cupuassu flower

References

Alves et al. 2003; Calzavara et al. (1984); Falcao & Lleras, E (1983); Maués et al. (2000); Venturieri (1993); Venturieri (1994); Venturieri (2011); Venturieri & Ribeiro Filho (1995)

Guava/Goiaba Psidium guajava L.

Status: Native

Growing areas: AC, AM, AL, BA, CE, MA, PE, PI, SE, MS, MT, ES, MG, RJ, SP, PR, RS, SC, PB, RN

Cultivation: Open plantations

Attractiveness score: 3

Pollinator dependency: Modest

Plant-mating system: Self pollination (autogamy, geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Self pollination by insects or wind is possible, and isolated trees often set satisfactory fruit numbers without cross pollination. Cross pollination is frequently aided by bees and other pollen carrying insects.







Flowering and harvest periods

Flowering: September to November Harvest: December to March

Main varieties grown in Brazil

Branca de Valinhos, Cortibel, IPA B-22, Kumagai, Ogawa, Paluma, Pedra Branca, Pedro Sato, Pentecostes, Pirassununga Vermelha, Rica, Ruby Supreme, Sassaoka, Tailandesa

Other information

The exclusion of pollinators other than wind reduces fruit set. Cross pollination increases yield. The presence of biotic pollinators like Apis mellifera and other bees promotes cross pollination in guava orchards and maximizes crop productivity. Guava trees are commonly planted in private gardens. Here they can be pollinated by hand with a brush, preferably with pollen of another guava tree but as self pollination leads to fruit set, the transport of pollen within a tree can also increase the production. As guava is also consumed by wildlife, the trees contribute to conserve biodiversity in private gardens and public parks. When planting guava trees it is recommended to plant at least two trees to optimize pollination.





Flowers are in cauliflorous inflorescences, hermaphrodite and radially symmetrical with four to six sepals and four to six white petals. \bigcirc organ with an inferior ovary, three to five locular and one long style that terminates in capitate stigma. \bigcirc organ with many stamens in many circles around the stigma and ellipsoid anthers.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Melipona mandacaia	Eusocial
Stingless bee	Partamona seridoensis	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Frieseomelitta doederleini	Eusocial
Stingless bee	Partamona cupira	Eusocial
Stingless bee	Tetragona dorsalis	Eusocial
Stingless bee	Trigona amalthea	Eusocial
Stingless bee	Trigona sp.	Eusocial
Oil-collecting bee	Centris aenea	Solitary
Carpenter bee	Xylocopa cearensis	Solitary
Carpenter bee	Хуlocopa sp.	Solitary
Other bee	Ēxomalopsis analis	Solitary

References

Alves & Freitas (2006, 2007); Freitas & Alves (2008); Hamilton, & Seagrave-Smith (1959); Hedström (1988); Siqueira et al. (2012); Tchuenguem Fohouo et al. (2007); Viana (2008)

Kiwifruit/Kiwi *Actinidia deliciosa* (A.Chev.) C.F.Liang & A.R.Ferguson

Status: Cultivated **Growing areas:** SP, SC, PR, RS **Cultivation:** Open plantations

Attractiveness score: 4

Pollinator dependency: Essential

Plant-mating system: Cross pollination (xenogamy), only few varieties are self fertile

Pollination requirements

Kiwifruit is dioecious and requires pollen transportation by wind or insects between male and female plant individuals. Although wind pollination is common, it is insufficient for a successful fruit yield. Entomophilous





pollination is essential to ensure high yields. With increasing bee visitation rates pollen deposition on stigmas, seed numbers and fruit weight increases.

Flowering and harvest periods

Flower: November to December

Harvest: January to February

Main varieties grown in Brazil

Abbott, Allison, Bruno, Hayward, Monty, Tomuri

Other information

Wind pollination results in 15% fruit set, while hand cross pollination reaches over 95% fruit set and open pollination 92% fruit set, in one study. Open pollination increases fruit weight on average by 25% compared to wind pollination. While syrphids and solitary bees are not known to pollinate kiwifruit, Apis mellifera is an effective kiwifruit pollinator. In New Zealand, but also Argentina, commercial kiwi production applies pollen spraying with different forms of suspensors. Pollen spraying in these countries includes products available to maintain the viability of kiwi pollen in a water-based suspension. With this suspension the spraying of pollen can be done under various weather conditions.





Flowers are dioecious and radially symmetrical with six white petals. \bigcirc flower with many carpels fused forming the style which terminates into radially organized stigmas. A wreath of anthers sheding sterile pollen surrounds the ovary. \bigcirc flower with many long stamens in a wreath and with no style.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Bumblebee	Bombus sp.	Social
Solitary bee	Leioproctus sp.	Solitary

References

Anonymous (1984); Blanchet et al. (1991); Clinch (1984); Clinch & Heath (1985); Clinch & Houten A.T. (1985); Costa et al. (1993); Donovan & Read (1990); Gonzalez et al. (1998); Goodwin et al. (1997); Howpage et al. (2001); Jay & Jay (1983); MacFarlane & Ferguson (1984); Manino et al. (1996); Miñarro & Twizell (2015); Palmer-Jones & Clinch (1974); Palmer-Jones & Clinch PG. (1975); Palmer-Jones & Clinch (1976); Sharma et al. (2013); Simonetto & Grellmann (1998); Testolin et al. (1991); Vassiere et al. (1996)
Lychee/Lichia Litchi chinensis Sonn.

Status: Cultivated, native to Southeast Asia **Growing areas:** SP, PR, MG, GO

Cultivation: Open plantations and gardens

Attractiveness score: 4

Pollinator dependency: Little

Plant-mating system: Cross pollination (xenogamy) and self pollination (geitonogamy)

Pollination requirements

Lychee flowers are monoecious with protandrous hermaphroditic dichogamy. Male and female flowering stages overlap within one plant individual, therefore, self pollination occurs, but cross pollination is much more common. Fruit set

requires the transfer of pollen from male flowers to the stigma of female flowers. Lychee







pollinated by bees, flies, ants and wasps.

flowers are entomophilous and usually

Flowering and harvest periods

Flower: August to September

Harvest: November to January

Main varieties grown in Brazil

Bengal, Brewster, Groff, Sweet Clift

Other information

Wind pollination resulted in 23, hand cross pollination in 45 and open pollination in 29 fruits/panicle, suggesting а pollination deficit in commercial plantations. Honeybees constitute 78% of the lycheepollinating insects. Fruit and seed weight is higher in cross-pollinated fruits than in selfpollinated fruits. Lychee can produce commercially without pollinators, but different beneficial insects frequently visit flowers. Therefore, if pesticides, especially insecticides, are applied their spraying during bloom needs to be avoided. The addition of honeybees to lychee plantations increased fruit set. Honeybees are managed for lychee production in India, Madagascar and other places of the world. It is recommended to use 5 to 10 honeybee colonies per ha. Lychee honey is known for a delicate scent reminiscent of roses.





Lychee inflorescences have many male and female flowers on one panicle. Flowers are monoecious and radially symmetrical without petals and with four to five tomentose sepals which open early. \mathcal{Q} flower with a short, well developed pistil. Ovary is long with two to four carpels, two lobes of the sticky stigma open down into a vertical cleft. Some flowers with five to eight short stamen and anthers containing little pollen. \mathcal{J} flowers with six to eight stamen that are much longer than the sepals and with hairy filaments. The anthers are elliptical and glabrous (bald, no hair). The rudimentary pistil is lacking both stigma and style.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Stingless bee Sweat bee Ladybird Lady beetle

Scientific name

- Apis mellifera Tetragonisca angustula Trigona sp. Melipona sp. Halictus sp. Coccinella septempunctata Harmonia axyridis
- Sociality Eusocial Eusocial Eusocial Solitary/Social Solitary Solitary



Stingless bee (*Tetragonisca angustula*) approaching a lychee flower



Solitary bee (*Halictus* sp.) on lychee flowers

References

Abrol (2006); Ali et al. (2013); Batten (1986); Batten & McConchie (1995); Mandal et al. (2016); Matos (2012); Morton (1987); Poonam et al. (2010); Rai et al. (2017); Kuman (2014); Singh et al. (2017); Somnuk & Suavansri (2005); Srivastava et al. (2017); Stern & Gazit (1996)

Mango/Manga *Mangifera indica* L.

Status: Cultivated, originates from Southeast Asia Growing areas: All states of Brazil Cultivation: Open plantations Attractiveness score: 4

Pollinator dependency: High Plant-mating system: Diu dichogamy limits autogamous s increases cross pollination (xe varieties are self-incompatible pollination in few varieties lik and self pollination in 'Keitt'. Pollination requirements Mango fruit set is generally lo number of fertile flowers (only Plant-mating system: Diurnal protogynous dichogamy limits autogamous self pollination and increases cross pollination (xenogamy), common varieties are self-incompatible and require cross pollination in few varieties like 'Tommy Atkins'

Mango fruit set is generally low compared to the number of fertile flowers (only 1-5% of flowers set fruit). Pollination of various insect

groups occurs. Especially highly mobile and abundant species such as honeybees and flies are expected to be efficient pollinators. Wind is under debate due to the short viability of pollen (12h) and stickiness of pollen especially in the beginning of anthesis. As mango varieties range from self-sterile to self-pollinating and fruit set is generally very low, further research and variety-specific information is needed.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

Bourbon, Coitê, Espada, Haden, Keitt, Kent, Ourinho, Palmer, Rosa, Tommy, Van dyke

Other information

Pollinator contribution to mango fruit set is estimated at about 50% of total fruit set production. In Brazil, native flies and honeybees have been documented as valuable pollinators of mango trees.









Flowers are organized in panicles bearing 500 to 10.000 flowers including both male and hermaphrodite, radially symmetrical flowers with usually five pubescent sepals, usually five greenish yellow or pale cream short petals, and a prominent disk, functioning as nectary, between the petals and stamens. Colour of sepals and petals changes with age from bright to dark. \bigcirc organ with spherical ovary and a style that emerges from one side. \eth organ with usually one longer fertile stamen and four shorter staminodes born from the disk, \eth flowers without gynoecium.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Small carpenter bee Carpenter bee Sweat bee Housefly Blow fly Blow fly Butterfly Ladybird Scientific name Apis mellifera Melipona sp. Ceratina sp. Xylocopa sp. Halictus sp. Musca domestica Chrysomya megacephala Papilio polytes Coccinella septempunctata Sociality Eusocial Eusocial Solitary Solitary Solitary Solitary Solitary Solitary Solitary



Fly on mango flowers

References

Amin et al. (2015); Bally (2006); Carvalheiro et al. (2010); Fajardo et al. (2008); Gajendra (1989); Huda et al. (2015); Kiill (2008); Kumari et al. (2014); Ramírez & Davenport (2016); Singh (1984); Sung et al. (2006); Tayeng & Gogoi (2016)

Melon/Melão Cucumis melo L.

Status: Cultivated, originates from Southwest Asia

Growing areas: AL, BA, CE, MA, PB, PE, PI, RN, SE, DF, GO, MS, MT, ES, MG, RJ, SP, PR, RS, SC

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: Essential

Plant-mating system: Usually self-pollinated (geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Melon plants are self-pollinated, but insect pollination is essential for pollen transfer from anthers to stigmas, within the hermaphrodite flowers or between male and hermaphrodite or





female flowers.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

Yellow (Natal, Goldex, Dali), Cantaloupe (Caribbean Gold, Rangers, Pitayo), Charentais (Magrite, Magistro), Galia (DRG 3228, Eldoor, Gladial), Piel del Sapo (Grand Prix, Ricura, Finura), Honey Dew (Dino, Orange Country

Other information

Many bee species show potential to pollinate Cucumis melo and to influence its productivity, but Apis mellifera is the Brazilian managed species used in plantations. Pollination by honeybees can increase yield per hectare and improve the size, quality and marketability of the fruit. In a recent study it was shown that different Brazilian melon varieties (agronomic types) (Cantaloupe, Charentais, Galia, Piel de sapo, Yellow) have different volatile organic compounds (VOCs). Some are attractive and some repellent for honeybees. Also, content varies from male nectar to hermaphrodite flowers inducing honeybees to favour the latter ones. It is therefore suggested to breed new varieties with VOC and nectar content attractive for bees and other pollinating insects in Brazil.





Flowers are monoecious or hermaphrodite, commonly male and hermaphrodite or female at the same plant. Hermaphrodite and female flowers are single, male flowers smaller and in clusters at the same plant, both are radially symmetrical with five yellow petals. \bigcirc organ with a three-lobed stigma on a short style and an inferior ellipsoid ovary. \bigcirc organ with five stamens and \bigcirc flowers without ovary.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona fulviventris	Eusocial
Stingless bee	Trigona pallens	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Bumblebee	Bombus sp.	Social
Long horned bee	Peponapis sp.	Solitary
Long-horned bee	Melissodes sp.	Solitary
Small carpenter bee	Ceratina sp.	Solitary
Leafcutter bee	Megachile sp.	Solitary
Sweat bee	Augochlora sp.	Solitary/Social
Sweat bee	Pseudaugochloropsis sp.	Solitary/Social
Sweat bee	Agapostemon sp.	Solitary/Social
Sweat bee	Halictus sp.	Solitary/Social
Other bee	Exomalopsis sp.	Solitary

References

Bomfim et al. (2016); Bomfim et al. (2019); Fernandes et al. (2019); Gomez et al. (2016); Goodell & Thomson (2007); Grewal & Sidhu (1978); Kaziev & Seidova (1965); Kiill et al. (2014, 2015, 2016); Lemasson (1987); Mann (1954); McGregor et al. (1965); McGregor & Todd (1952); Meléndez-Ramirez et al. (2002); Mouzin et al. (1980); Nerson (2009); Ribeiro et al. (2015); Stanghellini et al. (2002); Taylor (1955); Tschoeke et al. (2015); Williams (1987); Wilson et al. (2016)

Oil palm/Dendê Elaeis guineensis Jacq.

Status: Naturalized, originates from West and Southwest Africa

Growing areas: BA, PE, RN, PA, AM

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: High

Plant-mating system: Cross pollination (xenogamy)

Pollination requirements

Elaeis guineensis requires cross pollination through pollen transport from a neighbouring tree by wind or pollinators to set fruit. Beetles (Coleoptera) are the most numerous, diverse and commonly

associated pollinators of oil palm and therefore most valuable for crop production.



Flowering and harvest periods

Flowering: Year round Harvest: Year round

Main varieties grown in Brazil

Dura, Psifera, Tenera

Other information

The weevil genus *Elaeidobius* is a highly effective pollinator and specified on oil palm species. Pollination of oil palm by Elaeidobius kamerunicus is well studied and known to increase fruit set. Pollination of Elaeidobius sp. weevils is therefore essential for high production. The application of broad spectrum insecticides to control insect pest species also kills the pollinating weevils. Therefore, if insecticides are applied their application should be suspended during oil palm blooming and the weeks before blooming. As the weevils are living in the plantations all year around, the necessity of insecticides needs careful consideration. Moreover, honeybees frequently visit oil palm flowers, to avoid honey contamination, also pesticides should not be applied during the main flowering period in commercial plantations. A sustainable solution might be the use of natural enemies to control insect pests.



Flowers are monoecious with radially symmetrical male and zygomorph female flowers with three sepals and petals. \mathcal{D} flower with sterile stamens and a three-celled ovary with three spreading stigmas. \mathcal{D} flower with six stamens and a small, sterile pistil.

Flower visitors and pollinators

Common name

- Beetle Beetle Honeybee Stingless bee Stingless bee Leafcutter bee Carpenter bee Other bee
- Scientific name Elaeidobius kamerunicus Elaeidobius subvittatus Apis mellifera Trigona fulviventris Trigona sp. Megachile sp. Xylocopa sp. Lithurgus sp.
- Sociality Solitary Eusocial Eusocial Eusocial Solitary Solitary



Curculionid beetles on oil plam flowers



Honeybees (Apis melliferd) on oil palm flowers

References

Barcelos et al. (2015); Meléndez & Ponce (2016); Mayfield (2005); Moura et al (2008); Siregar et al. (2016)

Okra/Quiabo Abelmoschus esculentus (L.) Moench

Status: Cultivated

Growing areas: MG, RJ, SP

Cultivation: Open fields

Attractiveness score: 3

Pollinator dependency: Modest

Plant-mating system: Primarily self pollination (autogamy and geitonogamy) and a small degree of cross pollination (xenogamy)

Pollination requirements

Okra flowers are hermaphrodite and auto pollination is common, but cross pollination by insects also occurs. Auto pollination leads to 100% fruit set (pods). However, insect pollination

increases the number of seeds per pod, the seed weight and pod length, thus increases





yield. Flowering and harvest periods

Flower: Year round

Harvest: Year round

Main varieties grown in Brazil

Alecrim, Amarelinho, Campinas 1 (IAC-4075), Campinas 2 (IAC-4076), Chifre-deveado, Green Velvet, Santa Cruz 47, White Velvet

Other information

Despite okra flowers auto pollinate to set fruits, insect pollinators, especially bees, contribute to increase yield as they transfer more pollen grains to the stigma and promote self and cross pollination. Hence, although the number of fruits is usually not increased, the fruits become larger, heavier and bear more seeds after insect pollination. To increase yields at the small scale of a private garden, okra can be hand pollinated. Okra flowers open for one day only. Therefore, bud development needs to be observed carefully. For hand pollination pollen from the same or different plant individuals can be transferred using a slightly oily or wet (distilled water) brush or cotton swap. In hot weather and moist soil the fruit will develop in a few days.





Flowers are hermaphrodite and radially symmetrical with eight to 12 bracts, five sepals and five free petals forming a white to cream coloured Hibiscus-like corolla with a dark to pinkish centre. \bigcirc organ with superior ovary with five to ten fused styles and a dark red stigma. \bigcirc organ with many stamens fused into a tube around the pistil.

Flower visitors and pollinators

Common name

- Honeybee Bumblebee Leafcutter bee Carpenter bee
- Scientific name Apis mellifera Bombus sp. Megachile sp. Xylocopa sp.
- Sociality Eusocial Social Solitary Solitary



Honeybee (Apis mellifera) on an okra flower

References

Azo'o et al. (2011); Ghzawi et al. (2003); Ige & Eludire (2014); Malerbo-Souza et al. (2001); Njoya et al. (2005); Purewal & Randhawa (1947); Shalaby (1998); Singh et al. (2017)

Papaya/Mamão Carica papaya L.

Status: Naturalized, originates from the northern Neotropics

Growing areas: All states of Brazil

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: Little

Plant-mating system: Self pollination (geitonogamy) but cross pollination (xenogamy) can also occur

Pollination requirements

Commercially grown papaya flowers are usually hermaphrodite highly dimorphic, but also trees with only male or female flowers occur. Papaya is facultative self-pollinating with a low cross





pollination rate by pollinators.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

Formosa (Tainung n° 1, Tainung n° 2'), Mamão Comum, Mamão Papaia, Solo (Sunrise Solo, Improved Sunrise Solo cv 72/12')

Other information

Trees within and between varieties can be male, female or hermaphrodite. However commercially grown trees are mostly hermaphrodites. Papaya does not have typical bee-pollinated flowers, but they are visited by bees. The primary pollinators, around the world, are hawkmoths and some skipper butterflies. but for Brazil information about pollinator species is scarce. Cross pollination of pistillate flowers by insects produces fruit of high quality. Even though several pest insects can be problematic in commercial papaya production, the use of insecticides at least during flowering should be avoided to protect the nocturnal moth and diurnal bee flower visitors.





Flowers can be dioecious, male and female on different plant individuals, or hermaphrodite, but allways radially symmetrical with five petals. \mathcal{Q} flower with large functional pistil, ovoid ovary and no stamens. \mathcal{J} flower with ten stamens in two rows, with a pistillode. Hermaphrodite flowers with a combination of both functional \mathcal{J} and \mathcal{Q} organs.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Hawk moth	Erinnyis ello	Solitary
Butterfly	Papilio polytes	Solitary
Butterfly	Phoebis sennae	Solitary
Beetle	Acanthinus aequinoctialis	Solitary
Ant	Monomorium floricola	Eusocial
Stingless bee	Trigona sp.	Eusocial
Carpenter bee	Xylocopa sp.	Solitary
Hoverfly	Mesograpta sp.	Solitary
Ant	Brachymyrmex	Eusocial
Ant	Pheidole sp.	Eusocial
Ant	Solenopsis sp.	Eusocial
Thrips	Frankliniella sp.	Solitary

References

Allan (1963); Avila Jr. et al. (2012); Damasceno Jr. et al. (2009); Dey et al. (2016); Garrett (1995); Marin-Acosta (1969); Martins & Johnson (2009); Morrisen (1995)

Passion fruit/Maracujá Passiflora edulis Sims

Status: Native

Growing areas: AM, PA, TO, AL, BA, CE, MA, PB, PE, PI, RN, SE, DF, GO, MS, MT, ES, MG, RJ, SP, PR, RS, SC

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: Essential

pollination Plant-mating system: cross

Pollination requirements

Passiflora flowers are self-incompatible and show protandrous dichogamy, by releasing pollen before the stigma is receptive. The flowers of P. edulis have characteristics adapted to suit pollination by large bees, carpenter bees (Xylocopa spp.)

are considered the most effective pollinators of commercial passion fruit due to their morphological and behavioural characteristics.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

BRS Gigante Amarelo, BRS Ouro Vermelho, BRS Rubi do Cerrado, BRS Sol do Cerrado, FB 200, Golden Star, Marília

Other information

An important feature of the passion fruit flower is the androgynophore, an extension of the flower receptacle which elevates the stigma and stamens high above the petals. This requires large bodied insects, such as carpenter bees, large orchid bees or bumblebees, to touch the reproductive organs while consuming nectar. The absence of carpenter bees implies low production or producers to hand pollinate the flowers, which increases production costs. Alternatively, the protection of dead wood and old trees adjacent to the field can naturally attract carpenter bees by providing nesting structures.









Flower is hermaphrodite and radially symmetrical with five ovate, white petals and sepals and whorls of coronal appendages between the perianth and stamens. \bigcirc organ with three carpels with three styles and stigmas and a superior ovary. \bigcirc organ with five stamens.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Frieseomelitta varia	Eusocial
Stingless bee	Paratrigona lineata	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Trigona hyalinata	Eusocial
Bumblebee	Bombus pauloensis	Social
Bumblebee	Bombus morio	Social
Carpenter bee	Xylocopa suspecta	Solitary
Carpenter bee	Xylocopa cearensis	Solitary
Carpenter bee	Xylocopa hirsutissima	Solitary
Carpenter bee	Xylocopa grisescens	Solitary
Carpenter bee	Xylocopa brasilianorum	Solitary
Carpenter bee	Xylocopa frontalis	Solitary
Carpenter bee	Xylocopa ordinaria	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Oil-collecting bee	Centris scopipes	Solitary
Oil-collecting bee	Centris longimana	Solitary
Oil-collecting bee	Epicharis flava	Solitary
Oil-collecting bee	Épicharis bicolor	Solitary
Oil-collecting bee	Ĉentris denudans	Solitary
Oil-collecting bee	Centris lutea	Solitary
Oil-collecting bee	Centris flavifrons	Solitary
Oil-collecting bee	Centris sponsa	Solitary
Oil-collecting bee	Centris dorsata	Solitary
Oil-collecting bee	Centris similis	Solitary
Orchid bee	Eulaema nigrita	Solitary
List continues at p. 120	_	-

References

Baran et al. (2017); Bezerra et al. (2019); Freitas, & Oliveira Filho (2001, 2003); Freitas et al. (2009); Gaglianone (2008); Hoffmann et al. (2000); Kiill (2008); Oliveira (2008); Silva & Freitas (2018); Silveira et al. (2012); Siqueira et al. (2009); Yamamoto et al. (2012)

Peach/Péssego Prunus persica (L.) Batsch

Status: Cultivated, originates from Northwest China

Growning areas: BA, DF, ES, MG, RJ, SP, PR, RS, SC

Cultivation: Open plantations

Attractiveness score: 5

Pollinator dependency: High

Plant-mating system: Self pollination (autogamy, geitonogamy) in most varieties

Pollination requirements

Most varieties of peach are pollinated by honeybees and wild bees, transporting pollen within and between flowers.

Flowering and harvest periods

Flower: July to September Harvest: November to December

Main varieties grown in Brazil

Ágata, Ametista, Barbosa, BR-2, BR3, BR-6, BRS Rubimel, Capdebosq, Cerrito, Charme, Chimarrita, Chinoca, Chiripá, Chirua, Chula, Coral, Coral, Della Nona, Diamante, El dorado, Eldorado, Esmeralda, Farrapos, FLA (13-72), Flordaprince, Flordasun, Granada, Jade, Leonense, Maciel, Maciel, Magno, Marfim, Marli, Marli, Morro Redondo, Ônix, Pampeano, Pearl, Pilcha, Planalto, Precocinho, Premier, Riograndense, Safira, Sentinela, Sentinela, Sinuelo, Sulina, Vanguarda, Vila Nova

Other information

There is little information about peach pollination in Brazil. Depending on the varieties autogamous self pollination and wind pollination results in 7 to 35% fruit set. However, also depending on the variety pollination by *Apis mellifera* may increase fruit set up to 66%. Several wild bee species are visiting peach flowers and may complement honeybee pollination in space and time. Therefore, flower resources should be available in peach orchards or their surroundings. Bee houses can also be established close to the orchards to promote beneficial solitary bees and wasps.











Flowers are hermaphrodite and radially symmetrical with five hairy sepals and five pink petals. \bigcirc organ with one free carpel, composed of a long style terminating in a round stigma. \bigcirc organ with about 20 long stamens in three whorls.

Flower visitors and pollinators

Common name Honeybee Stinglees bee Stinglees bee Bumblebee Carpenter bee Ladybird

Scientific name Apis mellifera Melipona sp. Trigona spinipes Trigona sp. Bombus sp. Xylocopa sp. Coccinella septempunctata

Sociality Eusocial Eusocial Eusocial Social Solitary Solitary



Carpenter bee *(Xylocopa* sp.) on a peach flower



Stingless bee (Trigona sp.) on a peach flower



Honeybee (*Apis mellifera*) on a peach flower

References

Free (1993); Gariglio et al. (2009); Mayer et al. (2017); Mota & Nogueira-Couto (2002); Raj & Mattu (2014); Weinbaum et al. (1989); Zhang et al. (2015)

Pear/Pera Pyrus communis L.

Status: Cultivated, originates from mediterranean Europe and West Asia

Growing areas: BA, MG, SP, PR, RS, SC **Cultivation:** Open plantations

Attractiveness score: 3

Pollinator dependency: High

Plant-mating system: Primarily cross pollination (xenogamy), but self pollination and parthenocarpy can occur in some varieties

Pollination requirements

Most pear cultivars are self-incompatible and must be pollinated with pollen from a different cultivar. Therefore, pollination requires flowering times of different varieties to overlap and the presence of



insect pollinators.

Flowering and harvest periods

Flowering: August to September

Harvest: November to December

Main varieties grown in Brazil

Asiática, D'água, Europeia, Seleta (IAC 16-28), Triunfo (IAC 16-34), Williams, D'anjou, Ercolini, Rocha Portuguesa

Other information

Wind pollination usually results in no or little fruit set, therefore the activity of insects, mainly bees, in orchards is essential for high yields. As different wild bee species frequently visit pear flowers they should be promoted in the surrounding or in the orchards complement honeybee to pollination. Care should be taken not to create habitat in places where pesticides are applied. In general for orchard crops, bee houses are an option to promote and conserve beneficial solitary bees and wasps and hence to conserve biodiversity.





Flowers are hermaphrodite and radially symmetrical with five sepals and five white petals. \bigcirc organ with five carpels and an inferior ovary. \eth organ with 10-30 stamens with dark red anthers.

Flower visitors and pollinators

Common name

Honeybee Bumblebee Stingless bee Carpenter bee Ladybird

Scientific name Apis mellifera Bombus sp. Scaptotrigona sp. Xylocopa sp. Coccinella septempunctata

Sociality Eusocial Social Eusocial Solitary Solitary



Stringless bee (Scaptotrigona sp.) on a pear flower



Honeybee (Apis mellifera) on a pear flower

References

Free (1993); Hsieh et al. (2002); Jacquemart et al. (2006); Langridge & Jenkins. (1972); Lewis & Smith (1969); Maccagnani et al. (2003); Mayer & Lunden (1997); Monzon et al. (2004); Onarska et al. (2005); Quinet et al. (2016); Raj & Mattu (2014); Stern et al. (2004); Van den Eijnde (1996)

Persimmon/Caqui Diospyros kaki L.F.

Status: Cultivated, originates from China Growing areas: AL, BA, CE, MA, PB, PE, PI, RN, SE, DF, GO, MS, MT, ES, MG, RJ, SP Cultivation: Open plantations

Attractiveness score: 3

Pollinator dependency: Little

Plant-mating system: Self pollination (autogamy, geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Persimmon plants are self-pollinating, but entomophilous pollination can improve fruit set and yield. Persimmon trees may bear parthenocarpic fruits and depending on the cultivar, the plants may be monoecious, dioecious



STATION OF THE STATE

or hermaphrodites.

Flowering and harvest periods

Flowering: September to December

Harvest: February to June

Main varieties grown in Brazil

Costata, Fuyu, Hachiya, Kaoru, Mikado, Okira, Pomelo, Rama Forte, Regina, Rubi, Taubaté

Other information

The 'Giombo' cultivar produces fruits parthenocarpically, these are seedless fruits which form without pollination. By contrast, 'Fuyu' persimmon is visited the bv pollinators and produces fruits containing seed. In general, persimmon trees are variable and sometimes only bear female or only male flowers. Wild persimmon trees seem to be valuable pollinizers commercial varieties and a low degree in wind pollination is described in some papers. Therefore, although we classified little dependent persimmon as on pollinators promoting and conserving bees in landscapes with persimmon production should stabilize persimmon production. Trees in gardens can also be pollinated by hand with a brush to secure high production.





Flowers are hermaphrodite or more commonly female and male flowers separate. Flowers are radially symmetrical with a cream coloured four to five-lobed calyx tube. \bigcirc flowers with eight cream coloured carpels, four cellular ovary and eight infertile stamens. \bigcirc flowers with 16-24 pink tinged stamens in two rows on the petals.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Oil-collecting bee Carpenter bee Scientific name Apis mellifera Trigona sp. Centris sp. Xylocopa sp.

Sociality Eusocial Eusocial Solitary Solitary



Carpenter bee (*Xylocopa* sp.) on a persimmon flower



Stingless bee (Trigona sp.) on a persimmon flower



Honeybee (*Apis mellifera*) on a persimmon flower

References

Agustí. (2010); Campos et al. (2015); Chauhan et al. (2017); Free (1993); Giannini et al. (2015); Gould (1940); Hodgson (1938); Hodgson (1939); Martins &; Pereira (1989); McGregor (1976); Neuwald et al. (2009); Popenoe (1924); Ryerson (1927); Silva et al. (2016); Tessmer et al. (2014)

Plum/Ameixa Prunus domestica L.

Status: Cultivated

Growing areas: RS, SC, SP, PR

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: High

Plant-mating system: Most varieties are cross-pollinated (xenogamy)

Pollination requirements

The flowers of most cultivars self are pollen incompatible and require from an appropriate pollinizer cultivar and insect activity for cross pollination. Wind is not a good pollen vector for *Prunus* spp.



Flowering and harvest periods

Flowering: August to September Harvest: November to December

Main varieties grown in Brazil

Americana, Bluefree, D'Agen, Damson, D'Ente 707, Européia e Selvagem, Japonesa, Magnific, Ornamental, President, Stanley, Sugar

Other information

Honeybees can improve fruit set and yield. Cross pollination with other cultivars results in the highest initial and final fruit set, revealing the importance of pollinizer trees and pollinating insects for the fruit set of plum. Various wild social and solitary bee species visit plum flowers in Brazil (own observations). As these bees likelv complement pollination by honeybees they should be promoted and conserved in landscapes with plum orchards. Bee houses can be installed within the fields and natural habitat for nesting sites and flower resources providing bee forage before and after plum blooming can be created or conserved.







Flowers are hermaphrodite and radially symmetrical with five pubescent sepals and five white petals. \bigcirc organ with one free carpel and a long style terminating in a round stigma. \bigcirc organ with about 20 long stamens and yellow anthers.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Melipona sp.	Eusocial
Stingless bee	Trigona sp.	Eusocial
Bumblebee	Bombus sp.	Social
Ladybird	Coccinella [*] septempunctata	Solitary

References

Benedek & Nyeki (1996); Dordević et al. (2016); Free (1993); Hassan et al. (2007); Jun & Chung (2007); Mattu (2014); Raj & Mattu (2014); Sapir et al. (2007); Wadhwa & Sihag (2015)

Pumpkin/Abóbora Cucurbita maxima Duchesne

Status: Cultivated, originates from South American species Cucurbita andreana

Growing areas: All states of Brazil

Cultivation: Open fields

Attractiveness score: 4

Pollinator dependency: Essential

Plant-mating system: Self pollination (geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Cucurbita maxima is a monoecious plant, and pollen must be transferred from staminate male to pistillate female flowers to set fruit, either within the same or between different plants. Wind does

not contribute to pollination, only insects, such as honeybees transfer pollen. Increased





, such as honeybees transfer pollen. Increased pollen deposition on the stigma may optimize pumpkin fruit size, weight and seed number.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

1st Seca, Brasileira, Delicious, Exposição, Faxon's Brazilian, Golden Hubbard, Hokkaido, Hubbard, Italiana, Japonesa, Kabocha, Libanesa, Majestade, Mammoth Gold, Menina, Mini Jack, Moranga, Paulista, Tetsukabuto Chikara

Other information

Pumpkin set fruits with an insufficient amount of pollen placed on the stigmas, but these fruits are small, lighter in weight and with fewer seeds. Introduction of honeybee colonies can ensure sufficient bee visits to maximize pollen deposition and fruit size. On average, up to 16 honeybee visits per female flower increases fruit set, but excessive visitation can lead to the removal of pollen grains already deposited on the stigmas. Insect pollinator activity before about 9 a.m. is required to set fruit, likely due to ceasing viability of the pollen grains.



Flowers are monoecious with more male than female flowers on the same plant. Flowers are radially symmetrical with five free sepals with white hair and five yellow to orange petals that form a tubular corolla. \bigcirc flower with inferior ovoid unilocular ovary with short thick style and a three to five lobed stigma. \bigcirc flower with three filaments joining above the receptacle to form a column of stamens.

Flower visitors and pollinators

Common name

Honeybee Long horned bee Bumblebee Carpenter bee Sweat bee Sweat bee Stingless bee Beetle

Scientific name Apis mellifera Peponapis sp. Bombus sp. Xylocopa sp. Lasioglossum sp. Agapostemon sp. Trigona spinipes Diabrotica speciosa

Sociality Eusocial Solitary Social Solitary Solitary Eusocial Solitary



Singless bees (*Trigona* sp.) visiting a male pumpkin flower

References

Amarante & Macedo (2000); Fronk & Slater (1956); Matsumoto & Yamazaki (2013); Nicodemo et al. (2009); Pfister et al. (2017); Ramos et al. (2010); Robinson & Decker-Walters (1997); Shuler et al. (2005); Walters & Taylor (2006)

Rambutan/Rambutã Nephelium lappaceum L.

Status: Cultivated, originates from Southeast Asia

Growing areas: PA, BA, SP

Cultivation: Trees in open plantations

Attractiveness score: 3

Pollinator dependency: High

Breeding system: Cross pollination (xenogamy) with pollen transport between hermaphrodite and male trees is common (androdioecious), self pollination through insects (geitonogamy) may be possible Pollination requirements Rambutan is a cross-pollinated crop and depends on insects for pollination and fruit set. The flowers are highly attractive to many insects. However, the

are highly attractive to many insects. However, the hermaphrodite flowers are not known

to shed viable pollen, and cross pollination by male plants is thought essential for efficient production.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

Jit Lee, Rongrien, R134, R156, R162, R167, R170, R191, R193

Other information

Fruit mass can significantly increase and number of mature fruits can increase nine times in trees caged with pollinators and trees with open pollination compared to trees without pollinators. As rambutan is very attractive for social bees natural habitats adjacent to the plantations should be conserved to promote for example stingless bees. Honeybees could also be managed in rambutan plantations. Honey from rambutan flowers is well known as a traditional medicine, which can accelerate healing of oral mucosa wounds. It might therefore be of interest for some beekeepers to move their hives into the plantations during blooming.









Flowers are dioecious, either hermaphrodite or male flowers and radially symmetrical with four to seven nearly free, whitish sepals and yellowish or greenish petals that can be absent or reduced. \bigcirc organ with 2 carpels. \eth organ in hermaphrodite flowers with small stamens and not opening anthers and in male flowers with well developed stamens, but not opening stigma.

Flower visitors and pollinators

Common name Honeybee Stingless bee

Stingless bee

Scientific name Apis mellifera Trigona sp. Scaptotrigona sp. Sociality Eusocial Eusocial Eusocial



Singless bees (Scaptotrigona sp.) on rambutam flowers

References

Andrade (2012); Leão (2014); Lim (1984); Muhamed & Kurien (2018); Rincón-Rabanales et al. (2015); Sacramento & Andrade (2014)

Sesame/Gergelim Sesamum indicum L.

Status: Cultivated, originates from Southeast Asia Growing areas: MA, PI, CE, RN, PB, PE, AL, SE, BA, GO, MT

Cultivation: Open fields

Attractiveness score: 4

Pollinator dependency: Modest

Plant-mating system: Self pollination (auto- and geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Flowers of sesame auto pollinate around the time of anthesis, but its stigma remains receptive to receive auto or cross pollen from wind and insect pollinators, mainly bees.



Flowering: January to March Harvest: April to May

Main varieties grown in Brazil

BRS Seda, CNPA G2, CNPA G3, CNPA G4, IAC- Ouro, IAC-China, IAC-GUATEMALA

Other information

Although cross pollination by insects reduces seed set and weight per fruit, it increases fruit set and yields per plant. Sesame can be very beneficial to rotate in a cropping system. In rotation it can break the cycle of other crop pests such as nematodes and fungi and provide attractive food resources for many pollinating insect species. It can also be grown together with crops heavily depending on pollinators. For example a favorable crop rotation could be sesame with cotton. As cotton is usually treated with insecticides, in rotation with cotton, pollinator populations might recover during the sesame rotation time.







Flowers are hermaphrodite and bilaterally symmetrical with two bracts at the base, five petals, a calyx with oblonged lobes, five sepals, a bell–shaped white to violet corolla, with a purple spotted yellow throat. \bigcirc organ with a greyish hairy superior ovary and a style with a two-lobed stigma. \eth organ with four stamens and one staminod between the two upper shorter stamens.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Carpenter bee	Xylocopa grisescens	Solitary
Carpenter bee	Xylocopa carbonaria	Solitary
Carpenter bee	Xylocopa conf. suspecta	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Other bee	Melissoptila uncicornis	Solitary
Other bee	Dicranthidium arenarium	Solitary
Other bee	<i>Tetrapedia</i> sp.	Solitary
Other bee	Arhysoceble sp.	Solitary
Other bee	Anthidium sp.	Solitary
Leafcutter bee	Megachile sp.	Solitary
Sweat bee	Lasioglossum sp.	Solitary/Social
Wasp	Brachygastra lecheguana	Social
Southern green shield bug	Nezara viridula	Solitary

References

Andrade (2008); Andrade, et al. (2014); Ashri (2007); Porto et al. (2013); Kamel et al. (2013); Mahfouz et al. (2012); Mahmoud (2012); Napoletano (2008); Ngongolo et al. (2015); Pashte & Shylesha (2013); Porto (2013); Sarker (2004)

Soursop/Graviola Annona muricata L.

Status: Cultivated, originates from the northern Neotropics

Growing areas: AC, AM, AP, PA, RO, RR, TO, AL, BA, CE, MA, PB, PE, PI, SE, DF, GO, MS, MT, ES, MG, RJ, SP, RN

Cultivation: Open plantations, trees in gardens

Attractiveness score: 2

Pollinator dependency: Essential

Plant-mating system: Predominantly cross pollination (xenogamy), protogynous dichogamy reduces possibility of self pollination

Pollination requirements

The flowers are naturally cross-pollinated by beetles which transport pollen from flowers in the male phase to flowers in the female phase.

Flowering and harvest periods

Flowering: September to January Harvest: April to July

Main varieties grown in Brazil

Blanca, Boriundas, Crioula, Lisa, Morada

Other information

Beetles are attracted to the flowers for sheltering, mating and feeding on the nutritious tissues of the petals. During this activities they occasionally transport pollen and pollinate the flowers. Beetles of the genus Cyclocephala are the main pollinators of soursop. After leaving the flowers, female beetles often excavated holes in the soil to lay eggs. Larvae can be found between the leaf litter and the first layer of the soil. Therefore, no herbicides should be applied in soursop production to increase the availability of the natural pollinating beetles. However, due to the lack of reliable natural pollinators in agricultural settings, hand pollination is used to improve fruit set and guarantee increased production and better fruit quality than usually archived by natural pollination. Several videos on hand pollination are available for free in the internet.









Flowers are hermaphrodite and radially symmetrical with three tiny, broad triangular, green sepals and six thick fleshy, yellowish-green petals arranged in two whorls. The outer petals are larger and pointy ovate, the inner petals are thinner, smaller and rounded. Q organ with one ovule in each of numerous carpels with outstanding white, narrow pistil shaped stamens, with sticky stigmas. δ organ with numerous shield-shaped stamens and parallel anthers opening alongside.

Flower visitors and pollinators

Common name Beetle

Beetle Beetle Beetle Beetle

Scientific name Carpophilos domidiahus Cyclocephala gravis Cyclocephala vestita Cyclocephala hyrsuta Cyclocephala picipes

Sociality Solitary Solitary Solitary Solitary Solitary

Ants collect honeydew produced by scale insects at the flower stalk of soursop. The ants are unlikely to pollinate soursop but might protect the flower from other herbivores



References

Aguiar et al. (2000); Freitas (2012); Jalikop & Kumar (2007); Maia et al. (2012); Silva & Souza (1999); Vinay et al. (2017); Worrel et al. (1994)

Soybean/ Soja *Glycine max* (L.) Merr.

Status: Cultivated, originates from East Asia

Growing areas: MT, MS, GO, DF, RS, PR, MG, BA, TO, MA, PI, CE

Cultivation: Open fields

Attractiveness score: 4

Pollinator dependency: Modest

varieties Plant-mating system: Most are cleistogamous self-pollinating (autogamy), geitonogamy and cross pollination is possible after flower opening.

Pollination requirements

Soybean flowers usually auto pollinate before flowers completely open. Especially contribute to self- or cross pollination of open flowers. This increases the number of



bees

pods per plant and seed set per pod especially of flowers which failed to selfpollinate.

Flowering and harvest periods

Flowering: October to July

Harvest: January to October

Main varieties grown in Brazil

AS 3570, As 3575, AS 3590, AS 3610, AS 3680, AS 3797, AS 3810, AS 3820, AS 3850, BRS 217 (Flora), BRS 232, BRS 252 (Serena), BRS 257, BRS 279RR, BRS 283, BRS 284, BRS 314 (Gabriela), BRS 315RR (Lívia), BRS 317, BRS 333RR, BRS 359RR, BRS 360RR, BRS 361, BRS 378RR, BRS 388RR, BRS 399RR, BRS 413RR.

List continues at p. 119

Other information

The combination of honeybee colonies and presence of wild bee species in soybean plantations increases the number of pods plant and seeds per pod. The per attractiveness of flowers for bees is highly dependent on the soybean variety. Similar as for canola, soybean is pollinated by many wild bee species but fields provide no nesting resources, farmers should provide set aside patches with flowering herbs adjacent to and within large fields.





Flowers are hermaphrodite and laterally symmetrical with a tubular calyx of five hairy green sepals. The corolla is white or pale lilac to purple, with five petals, the posterior petal is outermost, the two lateral petals with long wings and the two anterior united petals form a keel. \bigcirc organ of one carpel with a superior elongated, green and unilocular ovary, the long style terminates with the stigma. \bigcirc organ with nine stamens fused into a tube, one posterior stamen is free.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Tetragonisca sp.	Eusocial
Stingless bee	Trigona fuscipennis	Eusocial
Stingless bee	Trigona sp.	Eusocial
Bumblebee	Bombus sp.	Social
Sweat bee	Dialictus sp.	Solitary/Social
Sweat bee	Halictus sp.	Solitary/Social
Oil-collecting bee	Centris analis	Solitary/Social
Other bee	Psaenythia sp.	Solitary
Other bee	Ancyloscelis sp.	Solitary
Other bee	Exomalopsis analis	Solitary
Other bee	Exomalopsis subtilis	Solitary
Other bee	Exomalopsis tomentosa	Solitary
Other bee	Exomalopsis ypirangensis	Solitary
Other bee	Florilegus sp.	Solitary
Other bees	Melitomella grisescens	Solitary
Other bee	Augochlora sp.	Solitary
Other bees	Augochlorella sp.	Solitary
Other bees	Augochloropsis sp.	Solitary
Leafcutter bee	Megachile sp.	Solitary

References

Abrams et al. (1978); Barella (2009); Chiang & Kiang (1987); Chiari et al. (2005a,b); Chiari et al. (2011); Erickson (1975a,b); Erickson (1984); Erickson et al. (1978); Fávero & Couto (2000); Gazzoni (2017); Issa et al. (1984); Juliano (1977); Kettle & Taylor (1979); Koelling et al. (1981); Mason (1979); Masuda & Goldsmith (2009); Milfont (2012); Milfont et al. (2013); Ortiz-Perez et al. (2008); Pinzauti & Frediani (1980); Rust et al. (1980); Sheppard et al. (1979); Toledo et al. (2011)

91

Strawberry/Morango Fragaria × ananassa Duchesne

Status: Cultivated, originates from a cross of a Chilean, a European and a Northamerican species

Growing areas: BA, GO, ES, MG, RJ, SP, PR, RS, SC

Cultivation: Open and tunnel-covered fields

Attractiveness score: 4

Pollinator dependency: Modest

Plant-mating system: Self pollination (autogamy, geitogamy)

Pollination requirements

Pollination in commercial fields likely results from a combination of wind and insect pollination and both self- and cross pollination. Wind alone produces few and underdeveloped fruits. Therefy





produces few and underdeveloped fruits. Therefore, pollination by stingless bees and honeybees is essential for both fruit production and quality.

Flowering and harvest periods

Flower: Year round

Harvest: Year round

Main varieties grown in Brazil

Albion, Aromas, Camarosa, Camino Real, Cristal, Diamante, Festival, Monterrey, Oso Grande, Palomar, Portola, San Andreas, Ventana

Other information

Open pollination of flowers by stingless bees and honeybees increases fruit set, yield and quality of fruits. The combination of cross pollination and flower visits by small and large bees at the same flower results in best quality fruits with long shelf life and high market value.

In Germany it was shown that woody structures adjacent to strawberry fields enhance bee diversity and strawberry yield and fruit quality.





Flowers are hermaphrodite and radially symmetrical with five sepals and five white round petals. \bigcirc organ with many free carpels on an elevated, cylindrical receptacle. \eth organ with many stamens with yellow anthers.

Flower visitors and pollinators

Common name

Honeybee Stingless bee Stingless bee Stingless bee Stingless bee Bumblebee Ladybird

Scientific name

Apis mellifera Nannotrigona testaceicornis Paratrigona lineata Scaptotrigona depilis Tetragonisca angustula Trigona recursa Bombus sp. Coccinella septempunctata Sociality Eusocial Eusocial Eusocial Eusocial Eusocial Social Solitary



Strawberry fruit development after A) insect pollination, B) self pollination and C) wind pollination



Compared to the large *Apis mellifera* in the picture above the small stingless bee *Scaptotrigona depilis* only touches the flower parts close to the petals where the nectar is produced and hence unlikely pollinates the center of the flower

References

Abrol et al. (2017); Antonelli et al. (1988a,b); Antunes et al. (2007); Chagnon (1993); Chang et al. (2001); Castle et al. (2019); Chen et al. (2011); Connor (1975); Goodman & Oldroyd (1988); Jacobs et al. (1988); Kakutani et al. (1993); Klatt et al. (2013); Malagodi-Braga & Kleinert (2004, 2007); Matsuka & Sakai (1989); McGregor (1976); Nye & Anderson (1974); Oliveira et al. (1991); Pion & Oliveira (1980); Roselino et al. (2009)

Sunflower/Girassol Helianthus annuus L.

Status: Cultivated, native to the Neotropics **Growing areas:** AL, BA, CE, PE, PI, RN, SE, DF, GO, MS, MT, ES, MG, SP, PR, SC, PB, TO

Cultivation: Open plantations

Attractiveness score: 5

Pollinator dependency: Modest

Plant-mating system: Most varieties with self pollination (geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Sunflower flowers mature from the margin to the centre of the flower head. Individual flowers are protandrous, with the male phase of pollen release

preceding the female phase in which the stigma is receptive to pollination. Thus, the sticky





pollen needs to be transferred between flowers within the same or of different flower heads usually by pollinators.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

AG 920, AG 930, Agrobel 910, Aguará 04, Aguará 3, BRS 321, BRS 323, BRS 324, BRS 387, BRS 390, BRS H250, Cargill 11, Catissol 01, Dow M734, Dow MG2, Embrapa-122, H251, H358, H360, IAC Iarama, IAC Uruguai, Multissol, Rumbosol 90

Other information

Honeybees are the most important insect pollinators. Pollen-gathering honeybees visit only flowers in the male phase, but nectar is produced in both male and female phases. Hence, nectar-collecting bees visit all flowers of the inflorescence, pollinating those flowers that are in the female phase. The introduction of honeybee colonies into sunflower plantations increases seed set, weight, germination rate and oil content and reduces the number of mal-formed seeds.





Flower heads are composed of up to 15000 individual hermaphrodite radially symmetrical disk flowers and mostly sterile bilaterally symmetrical yellow to reddish orange ray flowers with one long petal. Bracts surround the composed flower head. Disk flowers are small and pentamerous. \bigcirc organ with inferior ovary and a long, single style and a two-parted stigma. \eth organ with five stamens, these are absent in ray flowers.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Frieseomelitta doederleini	Eusocial
Stingless bee	<i>Friesomellita</i> sp.	Eusocial
Stingless bee	Plebeia aff. flavocincta	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Trigona fuscipennis	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Trigona hyalinata	Eusocial
Stingless bee	Melipona subnitida	Eusocial
Stingless bee	Melipona asilvai	Eusocial
Stingless bee	Melipona quadrifasciata	Eusocial
Stingless bee	Melipona scutellaris	Eusocial
Stingless bee	Nannotrigona testaceicornis	Eusocial
Stingless bee	Geotrigona sp.	Eusocial
Stingless bee	Partamona helleri	Eusocial
Stingless bee	Paratrigona sp.	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Stingless bee	Scaptotrigona sp.	Eusocial
Stingless bee	Tetragona sp.	Eusocial
Stingless bee	Tetragonisca sp.	Eusocial
Bumblebee	Bombus atratus	Social
Bumblebee	Bombus bellicosus	Social
List continued at p 121		

References

Basualdo et al. (2000); Bhowmik & Bhadra (2015); Calmasur &, Ozbek (1999); Carvalheiro et al. (2011); Castro & Leite (2018); Chambó et al. (2011); Cockerell (1914); Cruz & Freitas (2013); DeGrandi-Hoffman & Chambers (2006); DeGrandi-Hoffman & Martin (1993); DeGrandi-Hoffman & Watkins (2000); Delaude et al. (1979); Fell (1986); Free (1964); Furgala et al. (1979); Greenleaf & Kremen (2006); Krause & Wilson (1981); Langridge & Goodman (1981); Machado & Carvalho (2006)

List continues at p. 119
Sweet pepper/Pimentão Capsicum annuum L.

Status: Cultivated, originates from the northern Neotropics

Growing areas: All states of Brazil

Cultivation: Open and tunnelled plantations

Attractiveness score: 4

Pollinator dependency: Little

Plant-mating system: Self pollination (autogamy and geitonogamy), cross pollination (xenogamy) can also occur

Pollination requirements

Sweet pepper flowers are mostly autogamous selfpollinated. Cross pollination is mostly conducted by bees capable of buzz pollination. The resulting fruit quality is higher than of self pollinated fruits.

fruit quality is higher than of self-pollinated fruits. The flower of the sweet pepper is best

pollinated in the morning of the day when the flower opens, when the pollen is released and the stigma becomes receptive.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

Agronômico 10G, All Big, Amarelo SF 134, Andes Kobayashi, Ário, Athenas, Atlantis, Avelar, BEti R, Bruna R, Bruno, Camaro, Canarinho, Cascadura Ikeda, Cascadura Marina, Cida R, Commandant, Conrado, Dagma, Dahra R, Dahra RX, Derick, Elisa, Esplendor, Êxito, Fortuna Super, Gigante Ikeda, Green Belt Kobayashi, Hebron, Heloísa, Isabela, Itamara, Itapuã, Konan, Laser, Lotus, Magali, Magali R, Magda, Magna Super, Magnata, Marha R, Marli R, Marta, Melina, Mirella, Nádia, Paloma, Priscila, Proveito, Quadrado Vermelho, Rubi Giant, Rubia R, Satrapo-Sais, Stela, Sucesso, Taurus, Triunfo, Yolo Wonder

Other information

Pollination by bees increases fruit size, weight, number of seeds and reduces fruit malformation in sweet pepper.









Flower description

Flowers are hermaphrodite and radially symmetrical with five white to pale lilac, triangular, slightly cupped petals. \bigcirc organ with two carpels fused into a superior ovary. \bigcirc organ with five bluish stamens fused to the petals at the base.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Melipona subnitida	Eusocial
Stingless bee	Melipona scutellaris	Eusocial
Stingless bee	Nannotrigona testaceicornis	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Trigona recursa	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Paratrigona lineta	Eusocial
Bumblebee	Bombus atratus	Social
Sweat bee	Augochlora cf. morrae	Solitary/Social
Sweat bee	Augochlora morrae	Solitary/Social
Sweat bee	Augochlora thalia	Solitary/Social
Sweat bee	Augochlora sp.	Solitary/Social
Sweat bee	Augochlorella acarinata	Solitary/Social
Sweat bee	Augochlorella theia	Solitary/Social
Sweat bee	Augochloropsis aurifluens	Solitary/Social
Sweat bee	Augochloropsis cleopatra	Solitary/Social
Sweat bee	Augochloropsis caerulior	Solitary/Social
Sweat bee	Augochloropsis cupreola	Solitary/Social
Sweat bee	Augochloropsis heterochroa	Solitary/Social
Sweat bee	Augochloropsis laeta	Solitary/Social
Sweat bee	Augochloropsis wallacei	Solitary/Social
Sweat bee	Pereirapis rizophila	Solitary/Social
Sweat bee	Halictus lanei	Solitary/Social
Sweat bee	Lasioglossum picadense	Solitary/Social
Sweat bee	Lasioglossum ypirangense	Solitary/Social
List continuos at n 121		

List continues at p. 121

References

Cruz et al. (2004, 2005); Dag & Kammer (2001); Dag et al. (2007); De Ruijter et al (1991); Faria Jr et al. (2008); Reifschneider (2000); Jarlan et al. (1997); Pereira et al. (2015); Raw (2000) Roselino (2005); Roselino et al. (2010); Silva et al. (2005)

Tomato/Tomate Solanum lycopersicum L.

Status: Cultivated, originates from the northern Neotropics

Growing areas: All states of Brazil

Cultivation: Open fields and tunnels

Attractiveness score: 3

Pollinator dependency: Little

Plant-mating system: Self pollination (autogamy, geitonogamy) and cross pollination (xenogamy) can also occurs

Pollination requirements

Pollen grain release of the anthers and transfer to the stigma is strongly facilitated through wind, rain and other impacts vibrating the flower. Especially

buzz pollination by specific bees of the genera Melipona, Xylocopa and Bombus increases



fruit set, yield and quality.

Flowering and harvest periods

Flowering: Year round

Harvest: Year round

Main varieties grown in Brazil

Aliança, Andrea, Andrea Vistory, BRS Kiara, BRS Nagai, BRS Portinar, Carina, Débora Max, Débora Plus, Débora Victory, Delta, Gisele, Giuliana, Grande HT, IPA 6, Ivete, Júpiter, Kombat, Lumi, Pleno F1, Red Petit, Red sugar, Renata, Samambaia, San Vito, Santa Clara, Santa Clara VF 5600, Santa Cruz Kada, Sheila, Sindy, Sweet gold, Tyler, Tyna, Zamir

Other information

Wild bees that buzz pollinate increase the pollen load on the stigma and consequently fruit production and quality. Some bees that do not buzz-pollinate also contribute to tomato pollination by drumming the tips of the poricidal anthers with their forelegs or milking the anthers to release pollen. In open cultivation, wind shaking the flowers also contributes to self pollination.





Flower description

Flowers are hermaphrodite and radially symmetrical with five to ten green sepals and five bright yellow triangular, slightly cupped petals. \bigcirc organ with two carpels fused into a superior ovary with a green style and stigma. \bigcirc organ with five yellow stamens fused into a tube.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Paratrigona lineata	Eusocial
Stingless bee	Melipona quinquefasciata	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Frieseomelitta sp.	Eusocial
Stingless bee	Frieseomelita flavicornis	Eusocial
Stingless bee	Melipona quadrifasciata	Eusocial
Stingless bee	Geotrigona subterranea	Eusocial
Stingless bee	Tetragonisca angustula	Eusocial
Stingless bee	Tetragona sp.	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Stingless bee	Frieseomelitta sp.	Eusocial
Bumblebee	Bombus morio	Social
Bumblebee	Bombus pauloensis	Social
Bumblebee	Bombus sp.	Social
Carpenter bee	Xylocopa suspecta	Solitary
Carpenter bee	Xylocopa frontalis	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Oil-collecting bee	Čentris aenea	Solitary
Oil-collecting bee	Centris tarsata	Solitary
Oil-collecting bee	Centris tarsata	Solitary
Oil-collecting bee	Centris fuscata	Solitary
Oil-collecting bee	Centris varia	Solitary
List continues at p. 122		,

References

Banda & Paxton (1991); Bezerra & Machado (2003); Bin & Soressi (1973); Bohart & Todd (1961); Cauich et al. (2004); Cribb (1990); Del Sarto et al. (2004,2005); Deprá et al. (2014); Eijnde & Ruijter (1989); Franceschinelli et al. (2013); Gaglianone & Campos (2015); Higo et al. (2004); Sabara et al. (2004); Sabara & Winston (2003); Santos et al. (2009, 2014); Silva-Neto et al. (2013, 2017); Vinícius-Silva et al. (2017)

Watermelon/Melancia *Citrullus lanatus* (Thunb.) Matsum, & Nakai

Status: Cultivated, originates from West Africa

Growing areas: AC, AM, BA, CE, MA, PB, PE, RJ, SC, GO, RS, SP, PI, RN, TO, MG, ES, PR

Cultivation: Creeping vine-like plants in open plantations Attractiveness score: 4 Pollinator dependency: Essential Plant-mating system: Self pollination (geitonogamy) and cross pollination (xenogamy) Pollination requirements Watermelon flowers, including seedless varieties, need pollen transfer from male to female flowers



Flowering and harvest periods

Flower & Harvest: Year round

Main varieties grown in Brazil

Blackstone, Charleston Gray, Companion List continues at p. 119

Other information

A minimum number of viable pollen grains is needed to set a fruit, which usually, results from multiple pollinator visits. Fruit set in seeded watermelon can be maximized with more than 1,000 pollen grains per flower, which is accomplished by 6 to 8 honeybee or one bumblebee visitation. Pistillate flowers of seedless varieties need more than the double number of visits to set fruit. A unique situation exists for pollination in seedless (triploid) watermelons. They are not parthenocarpic and require a pollination stimulus for fruit set. Since triploid watermelon produce mostly nonviable pollen, the pollination stimulus must be provided by viable pollen grains from seeded diploid pollinizers that are planted in close proximity.







Flower description

Plants are usually monoecious, sometimes dioecious or andromonoecious, with radially symmetrical male and female flowers. The calyx is bell shaped with five lobes and a light yellow five-parted tubular corolla. \bigcirc flower with inferior ovoid unilocular ovary, with a short style and three stigmas. \bigcirc flower with three nearly free stamens and a gland resembling an ovary.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Stingless bee	Trigona fulviventris	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Stingless bee	Scaptotrigona sp.	Eusocial
Stingless bee	Melipona sp.	Eusocial
Bumblebee	Bombus sp.	Social
Cucumber bee	Peponapis sp.	Solitary
Long-horned bee	Melissodes sp.	Solitary
Carder bee	Hypanthidium sp.	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Small carpenter bee	Čeratina sp.	Solitary
Sweat bee	Lasioglossum sp.	Solitary/Social
Sweat bee	Agapostemon sp.	Solitary
Sweat bee	Augochloropsis sp.	Solitary/Social
Sweat bee	Halictus sp.	Solitary/Social
Sweat bee	Augochlorella sp.	Solitary/Social
Sweat bee	Augochlora sp.	Solitary/Social
Masked bee	Hylaeus sp.	Solitary
Leafcutter bee	Megachile sp.	Solitary
Other bee	Exomalopsis sp.	Solitary
Other bee	Heriades sp.	Solitary

References

Adlerz (1966); Alencar (2013); Azo'o et al. (2010); Bomfim, et al. (2015); Brewer (1974); Bussman et al. (2003); Goff (1937); Kaziev & Seidova (1965); Kremen et al. (2004); Kremen et al. (2002); Meléndez-Ramirez et al. (2002); Mohamed & El-Hafez (1974); Njoroge et al. (2004, 2010); Pinkus-Rendon et al. (2005); Pisanty (2014); Smith et al. (2013); Spangler & Moffett (1979); Stanghellini et al. (1998a,b & 2002a,b); Walters (2005); Walters & Schultheis (2009); Wilson et al. (2016); Winfree et al. (2008)

Zucchini/Abobrinha Cucurbita pepo L.

Status: Cultivated, originates from the Neotropics Growing areas: TO, BA, PB, DF, GO, MS, MT, MG, RJ, SP, PR, SC, ES

Cultivation: Open plantations

Attractiveness score: 4

Pollinator dependency: Essential

Plant-mating system: Self pollination (geitonogamy) and cross pollination (xenogamy)

Pollination requirements

Although zucchini/squash plants selfare compatible, they need pollinators to set fruits, because they have separate staminate (male) and pistillate (female) flowers at the same plant and the

pollen grains are too large and sticky to be carried by wind. Pollination is most successful

plant individuals.

in the morning and requires insect activity of for example honeybees to transfer pollen between flowers within one or different

Flowering and harvest periods

Flowering period: August to May

Harvest period: 30 to 45 after pollination

Main varieties grown in Brazil

Caserta SH-202, Caserta CAC, Caserta BR, Clarinda AG-135, Clarita, Cocozelle

Other information

Fruit set and seed production are strongly influenced by bee visitation. The percentage of fruit set, fruit weight, shape, and seed production are all affected by the number of bee visits and type of bee species. A minimum number of viable pollen grains (optimum 1,200 pollen grains per flower), which usually requires multiple pollinator visits (optimum e.g. 12 honeybee visits) is needed for fruit set.

The cucumber bee Peponapis sp. 15 specialized on zucchini and cucumber and other plants of this plant family. It is a ground-nesting bee and can be promoted by open, sandy grounds without herbicide applications adjacent or in the plantations.







Flower description

Flowers are separated in male and female flowers and radially symmetrical with five yellow petals. Male flowers are smaller, with a longer pedicel. \mathcal{Q} flower with two to three lobed stigma on a short style and an inferior ellipsoid ovary. \mathcal{J} flower with five stamens (tubular stamens that join from the fillet, forming a column) and no ovary.

Flower visitors and pollinators

Common name	Scientific name	Sociality
Honeybee	Apis mellifera	Eusocial
Bumblebee	Bombus morio	Social
Bumblebee	Bombus sp.	Social
Bumblebee	Bombus atratus	Social
Stingless bee	Frieseomelitta sp.	Eusocial
Stingless bee	Geotrigona mombuca	Eusocial
Stingless bee	Melipona seminigra	Eusocial
Stingless bee	Melipona quadrifasciata	Eusocial
Stingless bee	Melipona quinquefasciata	Eusocial
Stingless bee	Partamona combinata	Eusocial
Stingless bee	Paratrigona lineata	Eusocial
Stingless bee	Trigona hyalinata	Eusocial
Stingless bee	Trigona spinipes	Eusocial
Stingless bee	Trigona fulviventris	Eusocial
Stingless bee	Tetragona sp.	Eusocial
Stingless bee	Schwarziana mourei	Eusocial
Stingless bee	Trigona sp.	Eusocial
Stingless bee	Plebeia sp.	Eusocial
Cucumber bee	Peponapis sp.	Solitary
Carpenter bee	Xylocopa sp.	Solitary
Small carpenter bee	<i>Čeratina</i> sp.	Solitary
Orchid bee	Eulaema cingulata	Solitary
Sweat bee	Agapostemon semimelleus	Solitary
Sweat bee	Augochloropsis cupreola	Solitary/Social
Sweat bee	Halictus sp.	Solitary/Social
Sweat bee	Lasioglossum sp.	Solitary/Social
List southers at a 100		

List continues at p. 122

References

Artz & Nault (2011); Baptista (2016); Delaplane et al. (2000); Enriquez et al. (2015); Fronk & Slater (1956); Grewal & Sidhu (1978); Julier & Roulston (2009); Krug et al. (2010); Kubisova (1974); Nepi & Pacini (1993); Petersen et al. (2014); Petersen et al. (2013); Phillips (2013); Phillips & Gardiner (2015); Tepedino (1981); Torezani (2015) Vidal et al. (2010); Walters & Taylor (2006)

Comprehensive list of crops cultivated in Brazil with recommendations for a potential pollinatorsafe application of pesticides

This list comprises the crops cultivated in Brazil as listed by the FAO for the year 2017, complemented by additional crops we identified to be important for some regions in Brazil but not listed by the FAO. We refer to crops used at least partly for human consumption.

Note: As a general best practice recommendation to protect pollinators, all crops with an attractiveness score of 2 or > 2 should not be treated during flowering with pesticides. If there are no options to do the application before or after flowering, we recommend pesticide applications only with products explicitly identified as harmless for pollinators. We highlight these crops in bold.

Table 4: Overview of Brazilian crops with their bee attractiveness and their dependency on animal pollinators.

Attractiveness score (1- low or no attractiveness for animal pollinators to 5- highly attractive for animal pollinators, NA = degree of attractiveness unknown). Pollinator dependency categories (essential, high, modest, little, no poll = no insect pollinators needed, seeds = pollination only for seed production, NA). More information on pollinator attractiveness and pollination dependency are available at page 12 in the explanations of this guide. Scores and categories are used from Klein et al. (2007), other scientific papers and our expert knowledge.

Сгор	Latin name	Attract.	Poll. Dep.
Açaí palm/Açaí	Euterpe oleraceae	5	High
Acerola/Acerola	Malpighia emarginata	3	High
Agave/Agave	Agave sp.	3	Seeds
Alfalfa/Alfafa	Medicago sativa	4	High
Almond/Amêndoa	Prunus dulcis	4	High
Anise/Erva doce, Anis	Pimpinella anisum	3	Modest
Anatto/Urucum	Bixa orellana	4	High
Apple/Maçã	Malus domestica	4	High
Apricot/Damasco	Persea armeniaca	4	High
Artichoke/Alcachofra	Cynara cardunculus	5	Seeds
Asparagus/Aspargo	Asparagus officinalis	2	Seeds
Avocado/Abacate	Persea americana	4	High
Bambara bean/Amendoim do	Vigna subterranea	2	Little
Bambara	5		
Banana, Plantain/Banana, Bda-	Musa spp.	2	No poll
terra	11		1
Barley/Cevada	Hordeum vulgare	1	No poll
Blueberry/Mirtilo	Vaccinium sp.	4	High
Brazilian grapetree/Jabuticaba	Plinia cauliflora	3	Little
Brazil nut/Castanha do Pará	Bertholletia excelsa	4	Essential
Broad bean/Fava	Vicia faba	4	Modest
Buckwheat/Trigo-sarraceno	Fagopyrum esculentum	5	High
Cabbage, Cauliflower etc.	Brassica oleraceae	4	Seeds/modest
/Repolho, Couve-flor etc.			
Canola/Canola	Brassica napus	5	Modest
Cardamom/Cardamono	Elettaria cardamomum	4	High
Carob/Alfarrobeira	Ceratonia siliqua	3	Modest
Carrot/Cenoura	Daucus carota sativus	5	Seeds/ high

Crop	Latin name	Attract.	Poll. Dep.
Cashew/Caju	Anacardium occidentale	4	High
Cassava/Mandioca	Manihot esculenta	2	Seeds/little
Castor bean/Mamona	Ricinus communis	2	Modest
Cherry (sweet/sour)/Cereja	Prunus avium	5	High
Chestnut/Castanha portuguesa	Castanea sativa	4	Modest
Chick pea/Grão-de-bico	Cicer arietinum	3	Little
Chicory/Chicória	Cichorium intybus	4	Little/seeds
Chilli pepper/Pimenta	Catsicum annuum	3	Little
Cinnamon/Canela	Cinnamomum verum	2	NA
Citrus Citrus Line Clementine	Citrus son	4	Little
Grapefuit Mandarine Orange	Curras spp.	-	Little
Tangerine/Erutas cítricas			
Clover (red) /Trevo	Trifolium pratonsis	4	High
Cocoa/Cacau	Theobroma cacao	4	Essential
Cocoput/Coco	Cocos murifera	4	Modest
Coffee/Cefé	Cottos nutifiera	4	Modest
Common been / Foiião	Dhassolus milamis	4	Little
Control Deall/ Teljao	Company himseter	4	Modeet
Contion/Algodao	Gossyptum Intsulum	4	Soods/ILiab
Comander/Coentro	Cortanarum sativum	4	Julia
Cowpea/Feijao-caupi	V igna unguiculala	3	Little
Cranberry/Cranberry	V accinium macrocarpon	4	High
Cucumber/Pepino	Cucums sativus	4	High
Cupuassu/Cupuaçu	I heobroma grandiflorum	2	High
Currant/Groselha, Cassis	Ribes nigrum, R. rubrum	3	Modest
Date palm/Tamareira	Phoenix dactylifera	2	Little
Eggplant/Beringela	Solanum melongena	3	Modest
Fennel/Erva-doce, Funcho	Foeniculum vulgare	4	High
Fig/Figo	Ficus carica	2	Essential
Flax seeds/Linho, linhaça	Linum usitatissimum	3	Little
Garlic/Alho	Allium sativum	3	Seeds/little
Ginger/Gengibre	Zingiber officinale	NA	NA
Gooseberry/Groselha	Ribes uva-crispa	3	Modest
Grape/Uva	Vitis vinifera	2	No poll
Groundnut/Amendoim	Arachis hypogea	2	Little
Gourd/Cabaça	<i>Curcubita</i> spp.	4	Essential
Guava/Goiaba	Psidium guajava	3	Modest
Hazelnut/Avelã	Coryllus avellana	4	No poll
Hop/Lúpulo	Humulus lupulus	2	No poll
Karite nuts (sheanuts)/Carité	Vitellaria paradoxa	3	Modest
Kiwi/Kiwi	Actinidia deliciosa	4	Essential
Kola/Noz-de-cola	Cola spp.	NA	NA
Lentil/Lentilha	Lens esculenta	3	No poll
Lettuce/Alface	Lactuca sativa	2	Seeds
Linseed/Linhaça	Linum usitatissimum	3	Little
Lychee/Lichia	Litchi chinensis	4	Little
Lupin/Tremoço	Lupinus alba	5	Little
Maize/Milho	Zea mays	2	No poll
Mango/Manga	Mangifera indica	4	High
Mangosteen/Mangostim,	Garcinia mangostana	NA	No poll
Mangostão	0		1
Maté/Mate, Erva-mate	Ilex paraguariensis	3	NA seeds/high
Melon/Melão	Cucumis melo	4	Essential
Millet/Painco	Panicum miliaceum	1	No poll
Mushroom/Cogumelo	Agaricus historus A hrasiliensis	Ô	No poll
indomo on ooguneio	Pleurotus ostreatus and others	Ŷ	rto pon
Mustard/Mostarda	Sinapis arvensis	5	High
Nutmeg/Noz-moscada	Myristica fragrans	4	High
Nectarine/Nectarine	Prunus persica var. nucipersica	4	High
Oil palm/Dendê	Elaeis guineensis	4	High
Okra/Quiabo	Abelmoschus esculentus	3	Modest
-			

Crop	Latin name	Attract.	Poll. Dep.
Olive/Oliveira	Olea europaea	2	Little
Onion/Cebola	Allium cepa	4	Seeds/high
Papaya/Mamão	Carica papaya	4	Little
Passion fruit/Maracujá	Passiflora edulis	4	Essential
Peach/Pêssego	Prunus persica	5	High
Pepper (sweet)/Pimentão	Capsicum annuum	4	Little
Pear/Pera	Pyrus communis	3	High
Pea/Ervilha	Pisum sativum, P. arvense	2	No poll
Peppermint/Hortelã-pimenta	Mentha × piperita	5	Higĥ
Persimmon/Caqui	Diospyros kaki	3	Little
Pigeon pea/Feijão guandu	Cajanus cajan	3	Little
Pineapple/Abacaxi	Ananas comosus	2	Seeds
Pistachio/Pistache	Pistacia vera	2	No poll
Plum/Ameixa	Prunus domestica	4	Higĥ
Potato/Batata	Solanum tuberosum	2	Seeds
Sloe/Abrunho	Prunus spinosa	5	High
Pumpkin/Abóbora	Cucurbita maxima	4	Essential
Quince/Marmelo	Cydonia oblonga	4	Modest
Quinoa/Quinoa	Čhenopodium quinoa	3	Modest
Rambutan/Rambutã	Nephelium lappaceum	3	High
Raspberry/Framboesa	Rubus idaeus	4	High
Rice/Arroz	Oryza sativa	1	No poll
Rye/Centeio	Secale cereale	1	No poll
Safflower/Açafroa	Carthamus tinctorius	4	Little
Sesame/Gergelim	Sesamum indicum	4	Modest
Sorghum/Sorgo	Sorghum guineense, S. vulgare, S. dura	2	No poll
Soursop/Graviola	Annona muricata	2	Essential
Soybean/Soja	Glycine max	4	Modest
Spinach/Espinafre	Spinacia oleracea	2	No Poll
Strawberry/Morango	Fragaria × ananassa	4	Modest
String bean/Feijão-da-espanha	Phaseolus coccineus	4	High
Sugar beet/Beterraba	Beta vulgaris	1	No poll
Sugar cane/Cana de açúcar	Saccharum officinarum	1	No poll
Sunflower/Girassol	Helianthus annuus	5	Modest
Sweet potato/Batata doce	Ipomoea batatas	4	Seeds
Squash/Abóbora	Cucubita mixta, C. moschata	4	Essential
Tarot/Taro, Inhame	Colocasia esculenta	2	Seeds
Tea/Chá	Camellia sinesis	2	Little
Tomato/Tomate	Solanum lycopersicum	3	Modest
Triticale/Triticale	×Triticosecale	1	No poll
Truffle/Trufas	Different species of fungy	1	No poll
Vanilla/Baunilha	Vanilla planifolia, V. pompona	3	Essential
Vetche/Ervilhaca	Vicia spp.,	4	High
Walnut/Noz comum	Junglans regia	4	No poll
Watermelon/Melancia	Citrullus lanatus	4	Essential
Wheat/Trigo	Triticum aestivum	1	No poll
Yautia/Taioba	Xanthosoma sagittifolium	NA	Seeds
Zucchini/Abobrinha	Cucurbita pepo	4	Essential

Managing pollinators – description of pollinators that can be managed to enhance or stabilize crop production

Western honeybee Apis mellifera



Honeybee on anatto flower



Honeybee on acerola flower



Honeybee on cashew flower



Honeybee on coffe flower

Sociality: Eusocial

Colony size: 10,000 to 80,000 workers

Size: 12 to 13mm

Food resources: Pollen and nectar from many wild and cultivated plants.

Nesting location: Large hollow spaces, open spaces, tree branches/trunks, abandoned ant and termite nests but mainly kept by humans (beekeepers) in artificial hives.

Further resources: Plant resin, honeydew, water

Honeybees as crop pollinators: Honeybees are known to visit all crops listed in this pollination guide, except cocoa but although they visit açaí palm, acerola, annato, brazil nut, tomato, passionfruit and sweet pepper, they are usually not pollinating these crops as the flower morphologies do not fit to the bee morphology or because buzz pollination is required and honeybees cannot buzz pollinate.

Natural occurrence and management for crop pollination

The Western honeybee (*Apis mellifera*) was introduced to North America around 1600, from there the species spread across both American continents by establishing feral populations. In 1957 the so called African honeybee (*Apis mellifera scutellata*) was introduced to Brazil. Some swarms escaped and established populations throughout all subtropical and tropical parts of the Americas (Winston et al. 1981, Whitfield et al. 2006). Feral honeybees in Brazil are currently mostly Africanized (a hybrid of *A. m. scutellata* and European races of *A. mellifera*) honeybees. They establish colonies in many rain sheltered places such as hollow tree branches or trunks, roofs of buildings or abandoned ant and termite nests.

Roubik (1995) mentioned that stocking densities in fields with watermelon or coconut in tropical southeast Asia are 4-8 colonies per ha. Delaplane et al. (2000) suggest different stocking rates for different crops in the US. Deciding about the best stocking rate, when to place and for how long is complicated and also depends on the availability of wild bees around the crop fields. It is also important that if pesticides are applied, honeybees honeybees should not be placed in crop fields during their applications. Exemptions might be products specifically labelled as non-toxic to bees, allowing bee-safe application.

The management of honeybees in hives is an elegant way to enhance pollination of many crop species. As honeybee keeping is time consuming and even more complex in tropical than in temperate regions, we advise farmers to read specific literature about honeybee crop pollination management, join local beekeeping courses or to work with experienced beekeepers to use honeybees in their crop fields. As we can not address this complex topic within the limited space of our guide, we suggest that recommendations are needed in a specific guide on honeybee management for Brazil, such as "Polinizadores e Pesticidas: princípios e manejo para os agroecossistemas brasileiros" by Freitas & Pinheiro (2012).



Honeybee on a guava flower



Honeybee on a melon flower



Honeybee on an orange flower



Honeybee on canola flowers



Honeybee on a sesame flower



Honeybee on sunflower flowers



Honeybee on a sweet pepper flower



Honeybee on a soybean flower



Honeybee on a watermelon flower

Bumblebees Bombus spp.

Sociality: Social **Colony size:** up to 500 workers

Size: 11 to 17mm

Food resources: Pollen, nectar and honeydew of plants

Nesting location: Beneath leaves and debris, burrows in the ground

Further resources: Water, leaf litter

Bumblebees as crop pollinators: We identified from the scientific literataure five bumblebee species as crop pollinators in Brazil (*Bombus atratus* (syn.: *B. pauloensis*), *B. bellicosus*, *B. brevivillus*, *B. morio*, *B. trasversalis*). Crops visited by bumblebees from our guide are avocado, apple, brazil nut, canola, coffee, common bean, cotton, cowpea, melon, passion fruit, peach, pear, plum, pumpkin, strawberry, sunflower, sweet pepper, tomato, watermelon, and zucchini. The authors of this guide also observed bumblebees on persimmon and sesame. In general, bumblebee species are primarily found in higher altitutes and more temperate regions of Brazil.

Natural occurrence and management:

Bumblebee nests can be found in different locations and structures above and below the ground. In Europe *Bombus terrestris* is frequently breed for crop pollination. The biology and the behaviour of this European bumblebee is well known. This is different in Brazil and the biology of Brazilian bumblebees is different to the managed bumblebees in Europe and in the US. Hence, we can currently not recommend breeding bumblebees for crop pollination management. For example, Bombus atratus can have multiple queens laying eggs or a single queen per nesting cycle. Multiple queens lead to aggressive behaviour, which is not the case for the species bread in Europe. B. atratus nests can pertain longer times than these of European bumblebees, which should be an advantage to use colonies kept in artificial nests as crop pollinators. Again, bumblebee breeding is only it is infancy in South America, for example see Almanza Fandiño (2007). Instead of keeping bumblebee colonies in artifical nests, we encourage farmers to observe which wild and crop flowers are visited by bumblebees near their farms and to promote those flowers to increase the population size and numbers of wild bumblebees. Using for example the free mobile phone App iNaturalist (https://inaturalist.org), farmers can identify the bumblebees and other bees and the flowers they are visiting. With this information, they can find information on the biology and ecology of the bees and the plants to gather information how to promote bumblebees on their land.



Trigona spinipes on acerola flowers

Stingless bees Meliponini

Sociality: Eusocial Colony size: 100 to 100.000 workers Size: 2.6 to 14mm

Food resources: Pollen, nectar, fruit juice, organic liquids Nesting location: Large holow spaces, in the open, tree

trunks, tree branches, hollows in the ground, termite nests, artificial hives

Further resources: Resin, seeds, petals, plant fibers, water

Stingless bees as crop pollinators: More than 200 species of stingless bees are known from Brazil with 89 endemic species (Pedro et al. 2014). We identified 49 species of stingless bees visiting the crop species in our guide and we could identify stingless bee flower visits for almost all the crops listed in this guide, except for okra, pear and soursop. The often small-sized stingless bees also visit large crop flowers such as flowers of brazil nut or passion fruit but they provide limited pollination as their body size does not match the flower size of these crops. The most common stingless bee species visiting crop flowers is Trigona spinipes but Melipona subnitida, Melipona quadrifasciata, Nannotrigona testaceicornis, Paratrigona lineata and Tetragonisca angustula visit also several different crop species. Stingless bees can buzz pollinate flowers and are therefore important pollinators for crop plants requiring buzz pollination. Such plants belong for example to the plant family Solanaceae including crops like tomato or sweet pepper. As stingless bees are diverse and abundant and occur throughout Brazil they are important crop pollinators and should be protected and promoted by farmers. Stingless bees have reduced stingers and can not sting humans but defend their nests with biting, which can be uncomfortable. Futhermore, some stingless bees have very small colonies, some do not survive in artifical hives, and some collect large amount of plant resin making nest boxes sticky. Finally, stingless bees are also nectar thieves and are known to sometimes destroy crop flowers (Gutierrez-Chacón et al. 2018). Because of these reasons only few species are currently kept in artifical hives.

Natural occurrence and management

Stingless bees usually nest in hollow tree trunks or branches, abandoned termite nests, crevices between rocks or in belowground cavities. Stingless bees can be kept by beekeepers in their original log hive eg. when a beekeeper is cutting the branch with the colony inside, or they take a nest to transfer it to a wooden box to make hive control or honey and propolis harvesting easier. Meliponiculture (keeping stingless bees in artifical nest boxes) has a long tradition in Brazil and tradionally beekeepers keep each colony inside a wooden box without divisions. Later boxes were developed that allow nest divisions and honey extraction without damaging the colony. Today there are several box types for keeping stingless bees available and the hives developed in Embrapa Amazonia Oriental have different



Frieseomelitta longipes on brazil nut flowers



Trigona spinipes on cashew flowers



Trigona sp. on coffee flowers



Stingless bee on a kiwi flower

diameter entrances adapted to the size of different species. Stingless bee species like Melipona flavolineata, Melipona fasciculata, Melipona quadrifasciata, Scaptotrigona depilis,



Stingless bees (*Trigona* sp.) on a passion fruit flower



Trigona sp. on a sesame flower



Trigona sp. on a sweet pepper flower

Scaptotrigona bipunctata and *Tetragonisca angustula* were shown to accept artifical boxes and visit many crop species in our guide. For example, these bees pollinate açaí palm (Contrere et al. 2011), strawberry or tomato (Imperatriz-Fonseca et al. 2006). Stingless bee hives are still not regularly used for crop pollination, despite the long tradition and the ongoing research.



Halictid bee on an anatto flower



Solitary bee on tomato flowers



Augochlora sp. on a tomato flower

Solitary wild bees

Colony size: No colony; these bees live solitary with one female provisioning five to fifty brood cells in one or few nests or semi-social bees with female bees nesting in nest aggregates and sometimes sharing tasks such as protecting nest entrances.

Size: 4 to 30 mm

Food resources: Pollen, nectar, honeydew

Nesting location: Diverse, above- and belowground

Further resources: Plant resin, oil, leaves and fiber, water

Solitary wild bees as crop pollinators: The majority of bee species are solitary. We still do not know the exact species numbers but at a minimum of 1000 solitary to semi-social bee species occur in Brazil (estimated from Freitas et al. 2009). Many solitary bees are food specialists and do not visit many different crops in contrast to the generalistic honeybees, bumblebees and stingless bees. For example the solitary squash bee Peponapis sp. collects only pollen of the plant family Cucurbitaceae. A squash bee that occurs in Brazil is Peponapis fervens (Giannini et al. 2010). Crops of the family Cucurbitaceae include melon, watermelon, pumpkin and zucchini (although we did not find flower visits of this bee to zucchini flowers). Peponapis species nest like many other solitary wild bees below ground. Almost all crops are visited by solitary wild bees. Solitary wild bees visiting a large number of crops in Brazil are carpenter bees, especially Xylocopa frontalis and Xylocopa grisescens, but also the communal nesting bee Exomalopsis analis and the oil-collecting bee species Centris aenea, C. flavifrons and C. tarsata. Below descriptions of the carpenter bees and the oil-collecting bees are given.

Natural occurance and management of solitary

bees

Every solitary wild bee species has its own biology/ecology and hence different species live in different ecosystems, ranging from arid to humid and from lowland to mountain habitats. Bees nest in many different locations and substrates and require many different nest building materials. All in common is their need for the offspring to be undisturbed for the whole developmental period, which can be up to an entire year, and the favour for dry places for placing or rearing their

offspring. As bees collect pollen from plants as larval food, which is commonly contaminated with bacteria and moulds high humidity in the nests decreases larval survival. Most solitary bees have a short flight period of only several weeks, the flight

season can be up to 8 times a year or only once a year depending on the harschness of the environment and the therefore predictability of the flower resources the bees exploit.

To promote solitary wild bees, many different native flower types should be grown near the crop fields to provide year-around foraging habitat. Above-ground nesting solitary bees can be promoted with nesting aids. For this bamboo or reed can be cut in 15 to 20 cm pieces, placed in a plastic tube to hang in trees adjacent to or in the crop fields. Different bamboo or reed diameters should be used to allow many different bee species to use the nesting aid. Drilled hardwood can also be used as nesting aids. Nesting aids should be placed at a sunny spot that is not exposed to strong wind and weather, and should be covered by a roof.



Trap nest for solitary bees and wasps in a coffee plantation in Indonesia



Aggregations of ground-nesting bees establish in well drained soils, near to the food resources of the species



Below-ground solitary bees are difficult or nearly impossible to manage with nesting aids and should be promoted by habitat management. To encourage ground-nesting bees to nest in or around crop fields, open (with sparse vegetation), preferably sandy soils can be offered.

Nesting aids and open grounds are also used by many solitary waps species. Solitary bees and wasps are not aggressive and are both beneficial to farmers. Bees provide pollination services and wasps provide pest control by preying on caterpillars or aphids.

Large insect house for cavity-nesting solitary wild bee species. 10.000s of reed/bamboo sticks and holes drilled in the wood of different diameters provide a nesting space for many different species and generations. Such insect houses work in temperate and tropical regions



Xylocopa frontalis on a brazil nut flower



Xylocopa sp. on sesame flowers



Xylocopa grisescens on a watermelon flower



Xylocopa sp. on an anatto flower



Xylocopa grisescens male on a passion fruit flower



Xylocopa grisescens female on a passion fruit flower

Carpenter bees Xylocopa spp.

Sociality: Solitary

Colony size: No colonies are formed, but nesting of many females within the same location is facilitated through the high return potential of female bee offspring to their place of birth. Nest establishment can be increased by the provisioning of suitable nesting aids or natural nesting sites.

Size: 12 to 25 mm

Food resources: Pollen, nectar

Nesting location: Deadwood

Further resources: Clear space for perching and mating

Carpenter bees as crop pollinators: Carpenter bees visit many crop flowers and are known to be effective pollinators on anatto, brazil nut, common bean, cowpea, passion fruit, pumpkin, sesame, tomato, persimmon, watermelon, zucchini, melon, cotton, coffee, guava, sunflower

Natural occurrence and management of carpenter bees:

Carpenter bees build their nests, as their name suggests, in dead wood. Naturally, mainly in standing or fallen dead wood, especially tree trunks but poles and construction wood can also be attractive.

To promote carpenter bees, standing dead wood and dead branches should be conserved or actively established adjacent to the orchards where those bees are needed for pollination e.g. close to passion fruit fields. Nesting can also be artifically promoted through the provision of soft wood blocks of at least 30 x 30x 30 cm. The wood should be dry and constantly sheltered from rain and other sources of humidity. The use of sun absorbent material as rain shelter should be avoided as overheating of the nesting aids decreases occupation rates and larval survival. Occupation of the nesting aids can be facilitated

through the provision of dummy entrances, holes of 30mm depth and 8 to 12mm diameter, that are drilled in an angle of about 30° pointing upwards into the lower 1/3 of the front of the nesting aid. The wood of the nesting aid should be horizontal with the grain simulating the orientation of a fallen tree trunk, as carpenter bees construct their

nests along the grain which is facilitated by horizontal nest construction potential. However, vertical nests are built as well (Roubik 1995, Silva & Freitas 2018).

Carpenter bees also require a clear space with sky view in front of the nesting aids. To attract male bees, perching opportunities like poles should be located close to the nest. Further methods of carpenter bee nesting aid constructions are given in Freitas & Oliveira (2001) and Silva & Freitas (2018).

Oil-collecting bees Centris spp.

Sociality: Solitary

Colony size: *Centris* bees do not form colonies, but nesting aggregations can be promoted.

Size: 9 to 32mm

Food resources: pollen, nectar, floral oil

Nesting location: Deadwood, sandy soil

Further resources: Resins, sand, mud



Centris analis on acerola flowers

Oil-collecting bees as crop pollinators: Crop species commonly visited: Acerola, brazil nut, cashew, tomato, coffee, cotton, guava, passionfruit, persimmon

Natural occurrence and management of oil-collecting bees

Some oil-collecting species nest in pre-existing holes usually of wooden structures, others in the soil. Open, dry and sandy soils which are not disturbed heavily or deadwood in which other insects, like beetle and lepidopteran larvae drill their holes and eventually emerge from are the natural nesting sites. To promote *Centris* bees, holes of 5 to 9 mm and 8 to 12 cm depth, can be drilled in dry wooden blocks. The blocks need to be shelterd from rain or other sources of moisture. The wood should be drilled against the grain and hundreds of holes can be drilled with a minimum distance of 1 cm to each other into one or several wooden pieces. Only blind holes will be colonized by the bees, like *Centris analis* and *Centris tarsata*. The use of 4 to 6 cm wide, 50 cm high and 10 cm deep wooden beams equipped with 2 to 3 rows of holes is advisable as short dry wood pieces rarely form cracks. Cracks reaching the holes and hence the nests facilitate parasitism or nest destruction by mould. Another advantage of small wooden beams is that completed nests, indicated by a plug of resin or mud closing the holes, can carefully be transported to new locations.

References:

- Almanza Fandiño M.T. (2007). Management of *Bombus atratus* bumblebees to pollinate lulo (*Solanum quitoense* L), a native fruit from the andes of Colombia. pp. 50–63., Cuvillier Verlag, Göttingen, Germany
- Contrera F.A.L., Menezes, C., Venturieri, G. C. (2011) New horizons on stingless beekeeping (Apidae, Meliponini). Revista Brasileira de Zootecnia 40: 48-51
- Delaplane K.S., Mayer D.R., Mayer DF (2000) Crop pollination by bees. Cabi
- Freitas B.M., Imperatriz-Fonseca V.L., Medina L.M., et al. (2009) Diversity, threats and conservation of native bees in the Neotropics. Apidoligie 40: 332-346
- Freitas B.M., Pinheiro J.N. (2012) Polinizadores e Pesticidas: princípios e manejo para os agroecossistemas brasileiros. Brasília: Ministério do Meio Ambiente. Brazil

- Freitas B.M., Oliveira-Filho J.H. (2001) Criação racional de mamangavas: para polinização em áreas agrícolas. Fortaleza: Banco do Nordeste. Brazil
- Giannini T.C., Saraiva A.M, Alves dos-Santos I. (2010) Ecological niche modeling and geographical distribution of pollinator and plants: A case study of *Peponapis fervens* (Smith, 1879) (Eucerini: Apidae) and *Cucurbita* species (Cucurbitaceae). Ecological Informatics 5: 59-66
- Gutiérrez-Chacón C., Pantoja-Santacruz J., Klein A.M. (2018) Research on florivorous bees needed: The stingless bee *Trigona* cf. *amalthea* counteracts the contribution of other bees to the pollination of granadilla (*Passiflora ligularis* Juss). Journal of Pollination Ecology 22: 75-81
- Imperatriz-Fonseca V.L., Saraiva A.M., De Jong D. (2006) Bees as pollinators in Brazil: assessing the status and suggesting best practices. Holos, Ribirao Preto, Brazil
- Pedro S.R. (2014) The stingless bee fauna in Brazil (Hymenoptera: Apidae). Sociobiology, 61: 348-354
- Roubik D.W. (Ed.). (1995). Pollination of cultivated plants in the tropics. Food and Agriculture Organization of the United Nations
- Silva C.I., Freitas B.M. (2018) Rearing carpenter bees (*Xylocopa* spp.) for crop pollination: a case study with passionfruit (*Passiflora edulis*) In: Roubik (ed.) The pollination of cultivated plants: a compendium for practitioners. Vol. 2. Rome: FAO. Italy

Continued lists for crops and pollinators

Continued list of citrus varieties grown in Brazil:

Lime fruits: Fino, Lima da Pérsia, Lima Roque, Lima Sorocaba, Lima Tardia, Lima Verde; Mandarin oranges: Afourer, Dancy, Ellenor, Fina, Fina Sodea, Imperial, Lee Marisol, Minneola, Monreal, Murcott, Nules, Orlando, Osceola Page, Pummelo (Pomelo), Robinson, Satsuma, SRA63, Tango, Temple, Washington Navel (Bahia)

Continued list of soybean main varieties grown in Brazil:

BRS 8160 RR, BRS 8381, BRS 8480, BRS 8590, BRS 8890RR, BRS 9090RR, BRS 9280RR, BRS Aurora, BRS Barreiras, BRS Corisco, BRS FavoritaRR,BRS Gralha, BRS Jiripoca, BRS Pétala, BRS Pétala, BRS Raimunda, BRS Tracajá, BRS ValiosaRR, BRS-Carnaúba, BRSGO 7460RR, BRSGO 7950RR, BRSGO 7960, BRSGO 8151RR, BRSGO 8360, BRSGO Chapadões, BRSGO Chapadões, BRSGO Luziânia, BRSMG 68 (Vencedora), BRSMG 740SRR, BRSMG 750SRR, BRSMG 7525, BRSMG 760SRR, BRSMG 790A, BRSMG 800A, BRSMG 810C, BRSMG 811CRR, BRSMG 850RR, BRS-Sambaíba, M 9144 RR, M6009 RR, M6707 RR, M7211 RR, M7639 RR, M7908 RR, M8221 RR, M8230 RR, M8527 RR, M8766 RR, M8849 RR, M8867 RR, M9056 RR, MG/BR46 (Conquista)

Continued list of watermelon main varieties grown in Brazil:

Crimson Sweet, Fiesta, Giza I, Intruder, Kaho, Leeby, Leopard, Malali, Mickylee, Millionaire, Minipol, Polimore, Quetzali, Royal Jubilee, Royal Sweet, Sakata, Samara, Shadow, Shamhor, Shipper, SP-1, Sugar Baby, Sunshade, Tri-X 313, Yellow Elongate, Yellow Skin

Continued list of canola references:

Oliveira, 2017; Oz et al. (2008); Picard-Nizou et al. (1995); Pierre (2001); Pierre et al. (2010); Sabbahi et al. (2006); Shakeel et al. (2015); Steffan-Dewenter (2003); Tomm (2013); Williams (1985); Williams& Simpkins (1989); Witter et al. (2014)

Continued list of sunflower references:

Morgado et al. (2002); Nderitu et al. (2009); Neiva (2015); Oz et al. (2009); Parker (1981a,b); Parker & Frohlich. (1983); Pisanty et al. (2014); Saez et al. (2012); Skinner (1987); Tepedino & Parker (1982); Toledo et al. (2011); Wilson et al. (2016)

Continued list of cotton flower visitors and pollinators

Scientific name	Sociality
Centris sp.	Solitary
Eufriesea auriceps	Solitary
Eulaema nigrita	Solitary
Augochlora dolichocephala	Solitary/Social
Augochlora esox	Solitary/Social
Augochlora thalia	Solitary/Social
Augochloropsis patens	Solitary/Social
Pseudaugochlora graminea	Solitary/Social
Augochlorella acarinata	Solitary/Social
Augochlora sp.	Solitary/Social
Augochloropsis sp.	Solitary/Social
Pereirapis sp.	Solitary
Ceratalictus sp.	Solitary
Lithurgus huberi	Solitary
Alepidosceles imitatrix	Solitary
Exomalopsis analis	Solitary
Diadasina riparia	Solitary
Exomalopsis fulvofasciata	Solitary/Social
Exomalopsis auropilosa	Solitary/Social
Ptilothrix plumata	Solitary
Acamptopoeum prinii	Solitary
Tapinotaspoides serraticornis	Solitary
Exomalopsis sp.	Solitary/Social
Rhophitulus sp.	Solitary
	Scientific name Centris sp. Eufriesea auriceps Eulaema nigrita Augochlora dolichocephala Augochlora esox Augochlora thalia Augochloropsis patens Pseudaugochlora graminea Augochlorella acarinata Augochloropsis sp. Psereirapis sp. Ceratalictus sp. Lithurgus huberi Alepidosceles imitatrix Exomalopsis analis Diadasina riparia Exomalopsis fulvofasciata Exomalopsis auropilosa Ptilothrix plumata Acamptopoeum prinii Tapinotaspoides serraticornis Exomalopsis sp.

Continued list of canola flower visitors and pollinators

Common name	Scientific name	Sociality
Sweat bee	Neocorynura sp.	Solitary/Social
Sweat bee	Halictus sp.	Solitary/Social
Sweat bee	Lasioglossum sp.	Solitary/Social
Sweat bee	Augochloropsis cf. cupreola	Solitary/Social
Sweat bee	Augochloropsis melanochaeta	Solitary/Social
Sweat bee	Augochlorella	Solitary/Social
Sweat bee	Augochlora amphitrite	Solitary/Social
Sweat bee	Augochlora cf. francisca	Solitary/Social
Sweat bee	Caenohalictus tessellatus	Solitary/Social
Sweat bee	Paroxystoglossa brachycera	Solitary/Social
Sweat bee	Pseudagapostemon olivaceosplendens	Solitary/Social
Sweat bee	Pseudagapostemon pruinosus	Solitary/Social
Sweat bee	Pseudagapostemon tesselatus	Solitary/Social
Sweat bees	Augochloropsis sp.	Solitary/Social
Sweat bees	Augochlora sp.	Solitary/Social
Sweat bees	Caenohalictus sp.	Solitary/Social
Other bee	Thygater mourei	Solitary
Other bee	Thygater sp.	Solitary
Other bee	Anthrenoides elioi	Solitary
Other bee	Anthrenoides petuniae	Solitary
Other bee	Anthrenoides ornatus	Solitary
Other bee	Psaenythia sp.	Solitary
Other bee	Rhophitulus reticulatus	Solitary
Other bee	Rhophitulus sp.	Solitary
Other bee	Exomalopsis perikalles	Solitary
Other bee	Hexantheda missionica	Solitary
Other bees	Exomalopsis sp.	Solitary
Wasp	Protonectarina sylveirae	Social
Wasp	Brachygastra lecheguana	Social
Wasps	Eucyrtothynnus sp.	Solitary
Wasps	<i>Tiphia</i> sp.	Solitary
Wasps	Campsomeris sp.	Solitary
Beetle	Astylus variegatus	Solitary

Continued list of passion fruit flower visitors and pollinators

Common name	Scientific name	Sociality
Other bee	Acanthopus excellens	Solitary
Other bee	Oxaea austera	Solitary
Other bee	Oxaea flavescens	Solitary
Stingless bee	Scaptotrigona sp.	Eusocial
Stingless bee	Trigona sp.	Eusocial
Sweat bee	Augochlora sp.	Solitary/Social
Sweat bee	Augochloropsis sp.	Solitary/Social

Continued list of sunflower flower visitors and pollinators

Scientific name	Sociality
Bombus sp.	Social
Augochlora amphitrite	Solitary/Social
Augochloropsis callichroa	Solitary/Social
Augochlorella ephyra	Solitary/Social
Augochlora aff. Semiramis	Solitary/Social
Pseudaugochlora sp.	Solitary/Social
Megachile angularis	Solitary
Megachile paulistana	Solitary
Eumegachile sp.	Solitary
Megachile sp.	Solitary
Melissodes nigroaenea	Solitary
Melissoptila tandilensis	Solitary
Thygater analis	Solitary
Melissodes sp.	Solitary
Xylocopa augusti	Solitary
Xylocopa carbonaria	Solitary
Xylocopa grisescens	Solitary
Xylocopa suspecta	Solitary
Xylocopa sp.	Solitary
Čeratina sp.	Solitary
Agapostemon sp.	Solitary/Social
Pseudagapostemon sp.	Solitary/Social
Halictus sp.	Solitary/Social
Dialictus sp.	Solitary/Social
Paroxystoglossa sp.	Solitary/Social
Lasioglossum sp.	Solitary/Social
Megaloptina sp.	Solitary/Social
Euglossa cordata	Solitary
Eulaema nigrita	Solitary
Hypanthidium sp.	Solitary
Melissoptila sp.	Solitary
Oragapostemon sp.	Solitary
Oxaea sp.	Solitary
<i>Diadasia</i> sp.	Solitary
Doeringiella	Solitary
Nomada sp.	Solitary
Exomalopsis sp.	Solitary
Exomalopsis analis	Solitary
Exomalopsis auropilosa	Solitary
Nezara viridula	Solitary
Astylus atromaculatus	Solitary
Chlorostilbon aureoventris	Solitary
	Scientific name Bombus sp. Augochlora amphitrite Augochloropsis callicbroa Augochlora aff. Semiramis Pseudaugochlora sp. Megachile angularis Megachile angularis Megachile sp. Megachile sp. Melissodes nigroaenea Melissoptila tandilensis Thygater analis Melissodes sp. Xylocopa augusti Xylocopa augusti Xylocopa augusti Xylocopa grisescens Xylocopa grisescens Xylocopa grisescens Xylocopa sp. Ceratina sp. Agapostemon sp. Pseudagapostemon sp. Halictus sp. Dialictus sp. Paroxystoglossa sp. Lasioglossum sp. Megaloptina sp. Englossa cordata Eulaema nigrita Hypanthidium sp. Melissoptila sp. Oragapostemon sp. Oxaea sp. Diadista sp. Diadista sp. Diadista sp. Diadista sp. Melissoptila sp. Coragapostemon sp. Oxaea sp. Diadasia sp. Doeringiella Nomada sp. Exomalopsis sp. Exomalopsis analis Exomalopsis analis Exomalopsis analis Exomalopsis analis Exomalopsis auropilosa Nezara viridula Astylus atromaculatus Chlorostilbon aureoventris

Continued list of sweet pepper flower visitors and pollinators

Common name	Scientific name	Sociality
Masked bee	Hylaeus tricolor	Solitary
Other bee	Ēxomalopsis analis	Solitary
Other bee	Exomalopsis auropilosa	Solitary
Other bee	Exomalopsis fulvofasciata	Solitary
Wasp	Polybia sp.	Social
Fly	Toxomerus sp.	Solitary
Cucurbit Beetle	Diabrotica speciosa	Solitary

Continued list of tomato flower visitors and pollinators

Common name	Scientific name	Sociality
Oil-collecting bee	Epicharis flava	Solitary
Oil-collecting bee	<i>Épicharis</i> sp.	Solitary
Oil-collecting bee	Centris sp.	Solitary
Oil-collecting bee	<i>Epicharis</i> sp.	Solitary
Orchid bee	Eulaema nigrita	Solitary
Orchid bee	<i>Euglossa</i> sp.	Solitary
Long-horned bee	Thygater analis	Solitary
Sweat bee	Augochloropsis electra	Solitary/Social
Sweat bee	Pseudaugochlora erythrogaster	Solitary/Social
Sweat bee	Augochloropsis callichroa	Solitary/Social
Sweat bee	Pseudaugochlora graminea	Solitary/Social
Sweat bee	Augochloropsis smithiana	Solitary/Social
Sweat bee	Acamptopoeum prinii	Solitary/Social
Sweat bee	Augochloropsis cupreola	Solitary/Social
Sweat bee	Exomalopsis fulvofasciata	Solitary/Social
Sweat bee	Pseudaugochlora indistincta	Solitary/Social
Sweat bee	Augochloropsis sp.	Solitary/Social
Sweat bee	Lasioglossum sp.	Solitary/Social
Sweat bee	Pseudaugochlora sp.	Solitary/Social
Sweat bee	Halictus sp.	Solitary
Sweat bee	Anthophora sp.	Solitary/Social
Sweat bee	Augochlora sp.	Solitary/Social
Other bee	Exomalopsis analis	Solitary
Other bee	Exomalopsis auropilosa	Solitary
Other bee	Oxaea flavescens	Solitary
Other bee	Exomalopsis minor	Solitary
Other bee	Exomalopsis sp.	Solitary

Continued list of zucchini flower visitors and pollinators

Common name Scientific name Sociali	ty
Sweat bee Agapostemon sp. Solitary	7
Sweat bee Pseudaugochlora sp. Solitary	/Social
Sweat bee Augochlora sp. Solitary	/Social
Sweat bee Ceratalictus sp. Solitary	/Social
Sweat bee Dialictus sp. Solitary	/Social
Oil-collecting bee <i>Epicharis</i> sp. Solitary	7
Other bee Thygater analis Solitary	7
Other bee Exomalopsis analis Solitary	7
Other bee Exomalopsis auropilosa Solitary	7
Other bee Neocorynura Solitary	7
Other bee Pseudagapostemon sp. Solitary	7
Orchid bee Eulaema sp. Solitary	7

Scientific literature

Açaí palm/Açaí

Mainly pollinators and flower visitors:

- Campbell A.J., Carvalheiro L.G., Maués M.M. et al. (2018) Anthropogenic disturbance of tropical forests threatens pollination services to açaí palm in the Amazon river delta. Journal of Applied Ecology 55: 1725-1736
- Lamarão C.V., Nogueira, R.S.F., Yamaguchi, K.K. et al. (2018) Biotechnological potential of açaí (*Euterpe oleracea* and *Euterpe precatoria*). In: Tropical fruits -from cultivation to consumption and health benefits, fruits from the amazon. p.73-94, ISBN: 978-1-53612-839-0
- Nascimento W.N.O. (2008) Açaí (*Enterpe oleracea* Mart.) Informativo técnico rede de sementes da amazônia, n.18. Versão on-line ISSN 1679-8058
- Oliveira M.S.P., Carvalho J.E.U., Nascimento W.M.O. (2000) Açaí (*Euterpe oleracea* Mart.) série frutas nativas. Funep. ISBN 85-87632-17-5

Mainly crop pollination and production:

Venturieri G.C. (2008) Floral biology and management of stingless bees to pollinate assai palm (*Euterpe oleracea* Mart., Arecaceae) in eastern Amazon. In: Pollinators Management in Brazil. Ministry of Environment, Brazil

Acerola/Acerola

Mainly pollinators and flower visitors:

- Freitas B.M., Alves J.E., Brandao G.F., Araujo Z.B. (1999) Pollination requirements of West Indian Cherry (*Malpighia emarginata*) and its putative pollinators, Centris bees, In NE Brazil. The Journal of Agricultural Science 133: 303-311
- Magalhães C.B., Freitas B.M. (2013) Introducing nests of the oil-collecting bee *Centris analis* (Hymenoptera: Apidae: Centridini) for pollination of Acerola (*Malpighia emarginata*) increases yield. Apidologie 44: 234-239
- Oliveira G.A., Aguiar C.M.L., Silva M., Gimenes M. (2013) *Centris aenea* (Hymenoptera, Apidae): A groundnesting bee with high pollination efficiency in *Malpighia emarginata* DC (Malpighiaceae) Sociobiology 60: 317-322
- Vilhena A.M.G.F., Rabelo L.S., Bastos E.M.A.F., Augusto S.C. (2012) Acerola pollinators in the savanna of central Brazil: Temporal variations in oil-collecting bee richness and a mutualistic network. Apidologie 43: 51-62

Mainly crop pollination and production:

Calgaro M., Braga M.B. (2012) A cultura da acerola. Embrapa. ISBN 978-85-7035-130-2

Guedes R.S., Zanella F.C.V., Martins C.F., Schlindwein C. (2011) Déficit de polinização da aceroleira no período seco no semiárido paraibano. Revista Brasileira de Fruticultura 33: 465-471

Annato/Urucum

Mainly pollinators and flower visitors:

- Caro A., Moo-Valle H., Alfaro R., Quezada-Euán J.J.G. (2017) Pollination services of Africanized honey bees and native *Melipona beecheii* to buzz-pollinated annatto (*Bixa orellana* L.) in the neotropics. Agricultural and Forest Entomology 19: 274-280
- Castro C.B. (2009) A cultura do urucum/Embrapa Amazônia Oriental. Embrapa Informação Tecnológica, Belem/Brazil. ISBN 85-7383-451-X
- Costa A.J.C. et al. (2008) Abelhas (Hymenoptera: Apoidea) visitantes das flores de urucum em Vitória da Conquista, BA. Ciência Rural 38: 534-537
- Fabri E.G., Iatesta C.A., Rós-Gola A. et al. (2008) Ocorrência de Urucum no estado de São Paulo. Horticultura Brasileira 26: 4800-4803
- Rocha A.N., Polatto L.P. (2017) *Bixa orellana* L. (Bixaceae): dependência de polinizadores e estratégias de forrageio dos visitantes florais. Biota Amazônia 7: 1-7

Apple/Maçã

Mainly pollinators and flower visitors:

- Benedek P, Nyeki J. (1997) Yield of selected apple cultivars as affected by the duration of bee pollination. Acta Horticulturae 437: 207-212
- Blitzer E.J., Gibbs J., Park M.G., Danforth B.N. (2016) Pollination services for apple are dependent on diverse wild bee communities. Agriculture, Ecosystem and Environment 221: 1-7
- Boyle R.M.D., Philogène B.J.R. (1983) The native pollinators of an apple orchard: Variations and significance. Journal of Horticultural Science 58: 355-363
- Free J.B. (1962) The effect of distance from pollinizer varieties on the fruit set on trees in plum and apple orchards. Journal of Horticultural Science 37: 262-271
- Free J.B. (1966a) The pollinating efficiency of honey-bee visits to apple flowers. Journal of Horticultural Science 42: 91-94
- Free J.B. (1966b) The foraging areas of honeybees in an orchard of standart apple trees. Journal of Applied Ecology 3: 261-268
- Freitas B.M. (1995) The pollination efficiency of foraging bees on apple (*Malus domestica* Borkh) and cashew (*Anacardium occidentale* L.) PhD Thesis. University of Wales College of Cardiff. 228p., UK
- Nunes-Silva P., Rosa J.M., Witter S. et al. (2016) Visitants florais e potenciais polinizadores da cultura da macieira. Comunica Técnico 184. Bento Gonçalves: Embrapa. 16p., Brazil
- Viana B.F., Diakos A.C., Silva E.A. et al. (2015) Plano de manejo para polinização de macieiras da variedade Eva: conservação e manejo de polinizadores para a agricultura sustentável através de uma abordagem ecossistêmica. Rio de Janeiro: Funbio. 56p., Brazil

Mainly crop pollination and production:

- DeGrandi-Hoffman G., Hoopingarner R., Baker K.K. (1985) The influence of honey bee "Sideworking" behaviour on cross-pollination and fruit set in apples. Journal of Horticultural Science 20: 397-399
- Free J.B., Spencer-Booth Y. (1964) The foraging behaviour of honey-bees in an orchard of dwarf apple trees. Journal of Horticultural Science 39: 78-83
- Goodell K., Thomson J.D. (1997) Comparison of pollen removal and deposition by honeybees and bumblebees visiting apple. Acta Horticulturae 437: 103-108
- Hem R., Mattu V.K. (2014) Diversity and distribution of insect pollinators on various temperate fruit crops in Himachal Himalaya, India. International Journal of Science and Nature 5: 626-631
- Kendall D.A., Solomon M.E. (1973) Quantities of pollen on the bodies of insects visiting apple blossom. Journal of Applied Ecology 10: 627-634
- Kuhn E.D., Ambrose J.T. (1984) Pollination of "Delicious" apple by Megachilid bees of the genus Osmia (Hynenoptera: Megachilidae) Journal of the Kansas Entomological Society 57: 169-180
- Lewis T., Smith B.D. (1969) The insect faunas of pear and apple orchards and the effect of windbreaks on their distribution. Annals of Applied Biology 64: 11-20
- Mayer D.F. (1984) Behavior of pollinators on Malus. Les Colloques de l'INRA 21, France
- Park M.G., Raguso R.A., Losey J.E., Danforth B.N. (2016) Per-visit pollinator performance and regional importance of wild *Bombus* and *Andrena (Melandrena*) compared to the managed honey bee in New York apple orchards. Apidologie 47: 145-160
- Salomé J. (2014) Polinização dirigida em pomares de macieiras (*Malus x doméstica* Borkh) com o uso de colmeias de Apis mellifera L. PhD Thesis Universidade Federal de Santa Catarina, Centro de Ciências Agrárias, Programa de Pós-Graduação em Recursos Genéticos Vegetais, Florianópolis, Brazil
- Schneider D., Stern R.A., Eisikowitch D., Goldway M. (2002) The relationship between floral structure and honeybee pollination efficiency in 'Jonathan' and 'Topred' apple cultivars. Journal of Horticultural Science and Biotechnology 77: 48-51
- Smith B.D., Lewis T. (1972) The effect of windbreaks on the blossom-visiting fauna of apple orchards and on yield. Annals of Applied Biology 72: 229-238
- Stern R.A., Eisikowitch D., Dag A. (2001) Sequential introduction of honeybee colonies and doubling their density increases cross-pollination, fruit set and yield in "Red Delicious" apple. Journal of Horticultural Science and Biotechnology 76: 17-23

- Storhaug Y.G. (2014) Pollination as an ecosystem service in Lofthus, Norway: a study on the distribution of wild and managed pollinators on apple crops and how they are affected by the surrounding landscape. Master thesis, University of Bergen, Norway
- Torchio P.F. (1985) Osmia lignaria propinqua Cresson, in apple orchards: (1979-1980), methods of introducing bees, nesting success, seed counts, fruit yields (Hymenoptera: Megachilidae) Journal of the Kansas Entomological Society 58: 448-464
- Vicens N., Bosh J. (2000) Weather-dependent pollinator activity in an apple orchard, with special reference to Osmia cornuta and Apis mellifera (Hymenoptera: Megachilidae and Apidae) Environmental Entomology 29: 413-420

Avocado/Abacate

Mainly pollinators and flower visitors:

- Clark O.I. (1923) Avocado pollination and bees. California Avocado Association Annual Report 1922-1923 8: 57-62
- Davenport T.L. (1986) Avocado flowering. Horticultural Reviews 8: 257-289
- Ish-Am G., Barrientos-Priego F., Castaneda-Vildozola A., Gazit S. (1999) Avocado (Persea americana Mill.) pollinators in its region of origin. Revista Chapingo Serie Horticultura 5: 137-143
- Ish-Am G., Eisikowitch D. (1993) The behaviour of honey bees (*Apis mellifera*) visiting Avocado (*Persea americana*) flowers and their contribution to its pollination. Journal of Apicultural Research 32: 175-186
- Ish-Am G., Eisikowitch D. (1998) Low attractiveness of avocado (*Persea americana* Mill.) flowers to honeybees (*Apis mellifera* L.) limits fruit set in Israel. Journal of Horticultural Science and Biotechnology 73: 195-204
- Ish-Am G., Regev Y., Peterman Y. et al. (1998) Improving avocado pollination with bumble bees: 3 seasons summary. California Avocado Society Yearbook 82: 119-135
- Papademetriou M.K. (1976) Some aspects of the flower behaviour, pollination and fruit set of avocado (Persea americana Mill.) in Trinidad. California Avocado Society Yearbook 60: 106-152
- Perez-Balam J., Quezada-Euan J.J.G., Alfaro-Bates R. et al. (2012) The contribution of honeybees, flies, and wasps to avocado (*Persea americana*) pollination in southern Mexico. Journal of Pollination Ecology 8: 42-47
- Read S.F.J., Howlett B.G., Jesson L.K., Pattemore D.E. (2017) Insect visitors to avocado flowers in the Bay of Plenty, New Zealand. Forest Pest and Beneficial Insects 70: 38-44
- Vithanage V. (1990) The role of the European honeybee (*Apis mellifera* L.) in avocado pollination. Journal of Horticultural Science 65: 81-86

Brazil nut/Castanha do Pará

Mainly pollinators and flower visitors:

- Maués M.M. (2002) Reproductive phenology and pollination of the brazil nut tree (*Bertholletia excelsa* Humb. and Bonpl. Lecythidaceae) in eastern Amazonia. Pollinating Bees The Conservation Link Between Agriculture and Nature Ministry of Environment, Brasilia: p.245-254
- Nelson B.W., Absy M.I., Barbosa E.M., Prance G.T. (1985) Observation on flower visitors to *Bertholletia excelsa* H.B.K and *Couratari tenuicarpa* A.C.Sm. (Lecythidaceae) Acta Amazonica 15: 225-234
- Prance G. (1976) The pollination and androphore structure of some Amazonian Lecythidaceae. Biotropica 8: 235-241
- Santos C.F., Absy M.L. (2010) Polinizadores de *Bertholletia excelsa* (Lecythidales: Lecythidaceae): Interacoes com abelhas sem ferrao (Apidae: Meliponini) enicho trofico. Neotropical Entomology 39: 854-861
- Santos C.F., Absy M.L. (2012) Interactions between carpenter bees and orchid bees (Hymenoptera: Apidae) in flowers of *Bertholletia excelsa* Bonpl. (Lecythidaceae) Acta Amazonica 42: 89-94

Mainly crop pollination and production:

- Cavalcante M.C. (2008) Visitantes florais e polinização da castanha-do-brasil (*Bertholletia excelsa* H. & B.) em cultivo na amazônia central. Dissertação (mestrado) Universidade Federal do Ceará, Centro de Ciências Agrárias, Departamento de Zootecnia, Fortaleza-CE, Brazil
- Cavalcante M.C., Oliveira F.F., Maués M.M., Freitas B.M. (2012) Pollination requirements and the foraging behavior of potential pollinators of cultivated brazil nut (*Bertholletia excelsa* Bonpl.) trees in Central Amazon rainforest. Psyche Volume 2012, Article ID 978019

- Cavalcante M.C., Galetto L., Maués M.M. et al. (2018) Nectar production dynamics and daily pattern of pollinator visits in brazil nut (*Bertholletia excelsa* Bonpl.) plantations in Central Amazon: implications for fruit production. Apidologie 49: 505–516
- Maués M.M., Krug C., Wadt, L.H.O. et al. (2015) A castanheira-do-brasil: avanços no conhecimento das práticas amigáveis à polinização. Rio de Janeiro: Funbio, 84p., Brazil
- Ministério Da Agricultura, Pecuária e Abastecimento (2014) Castanha-do-pará, castanha, castanha-do-brasil, Bertholletia excelsa H.B.K. Secretaria de Desenvolvimento Agropecuário e Cooperativismo. - Brasília: MAPA/ACS, 41p.

Brazilian Grapetree/Jabuticaba

Mainly pollinators and flower visitors:

- Gobato M.R.R., Gobato R., Heidari A. (2018) Planting of Jaboticaba trees for landscape repair of degraded area. Landscape Architecture and Regional Planning 3: 1-9
- Gressler E., Pizo M.A., Morellato L.P.C. (2006) Polinização e dispersão de sementes em Myrtaceae do Brasil. Brazilian Journal of Botany 29: 509-530
- Malerbo D.T.S., Toledo, V.A.A., Couto, R.H.N. (1991) Polinização entomófila em jabuticabeira (Myrciaria cauliflora Berg.) Ciência Zootécnica Jaboticabal 6: 3-5
- Malerbo-Souza, Nogueira-Couto R.H., Toledo V. de A.A. de T. (2004) Abelhas visitantes nas flores da jabuticabeira (*Myrciaria cauliflora* Berg.) e produção de frutos. Maringá, 26: 1-4
- Morton J. (1987) Jaboticabas. p.371-374. In: Fruits of warm climates. Miami, FL, USA

Canola/Canola

Mainly pollinators and flower visitors:

- Adegas J.E.B., Nogueira Couto R.H. (1992) Entomophilous pollination in rape (Brassica napus L var oleifera) in Brazil. Apidologie 23: 203-209
- Arthur A.D., Li J., Henry S., Cunningham S.A. (2010) Influence of woody vegetation on pollinator densities in oilseed Brassica fields in an Australian temperate landscape. Basic and Applied Ecology 11: 406-414
- Benedek P., Prenner J. (1972) Effect of temperature on the behaviour and pollinating efficiency of honeybees on winter rape flowers. Zeitschrift für Angewandte Entomology 71: 120-124
- Blochtein B., Witter S., Halinski R. (2015) Plano de manejo para polinização da cultura da canola: conservação e manejo de polinizadores para agricultura sustentável, através de uma abordagem ecossistêmica. Rio de Janeiro: Funbio. 40p., Brazil
- Bommarco R., Marini L., Vaissiere B.E. (2012) Insect pollination enhances seed yield, quality, and market value in oilseed rape. Oecologia 169: 1025-1032
- Brunel E., Mesquida J., Renard M., Tanguy X. (1994) Repartition de l'entomofaune pollinisatrice sur des fleurs de colza (*Brassica napus* L) et de navette (*Brassica campestris* L): incidence du caractere apetale de la Navette. Apidologie 25: 12-16
- Chifflet R., Klein E.K., Lavigne C. et al. (2011) Spatial scale of insect-mediated pollen dispersal in oilseed rape in an open agricultural landscape. Journal of Applied Ecology 48: 689-696
- Eisikowitch D. (1981) Some aspects of pollination of oil-seed rape (Brassica napus L.) Journal of Agricultural Science 96: 321-326
- Free J.B., Ferguson A.W. (1980) Foraging of bees on oil-seed rape (*Brassica napus* L.) in relation to the stage of flowering of the crop and pest control. Journal of Agricultural Science 94: 151-154
- Free J.B., Ferguson A.W. (1983) Foraging behavious of honeybees on oilseed rape. Bee World 64: 22-24
- Free J.B., Nuttall P.M. (1968) The pollination of oilseed rape (*Brassica napus*) and the behaviour of bees on the crop. Journal of Agricultural Science 71: 91-94
- Hayter K.E., Cresswell J.E. (2006) The influence of pollinator abundance on the dynamics and efficiency of pollination in agricultural *Brassica napus*: Implications for landscape-scale gene dispersal. Journal of Applied Ecology 43: 1196-1202
- Holzschuh A., Dormann C.F., Tscharntke T., Steffan-Dewenter I. (2011) Expansion of mass-flowering crops leads to transient pollinator dilution and reduced wild plant pollination. Proceedings of the Royal Society

B, Series B 278: 3444-3451

- Jauker F., Bondarenko B., Becker H.C., Steffan-Dewenter I. (2012) Pollination efficiency of wild bees and hoverflies provided to oilseed rape. Agricultural and Forest Entomology 14: 81-87
- Jenkinson J.G., Glynne Jones G.D. (1953) Observations on the pollination of oil rape and broccoli. Bee World 34: 173-177
- Kamel S.M., Mahfouz H.M., Blal A.E.H. et al. (2015) Diversity of insect pollinators with reference to their impact on yield oroduction of canola (*Brassica napus* L.) in Ismailia, Egypt. Pesticidi I Fitomedicina 30: 161-168
- Kevan P.G., Eisikowitch D. (1990) The effects of insect pollination on canola (Brassica napus L. cv. OAC Triton) seed Germination. Euphytica 45: 39-41
- Koltowski Z. (2001a) Beekeeping value and pollination requirements of double-improved cultivars of spring rapeseed (*Brassica napus* L. var. *oleifera* Metzger F. Annua Thell.) Journal of Apicultural Science 45: 69-84
- Koltowski Z. (2001b) Results of the investigations into nectar secretion and pollen production of new cultivars of rape (*Brassica napus* L.) Acta Horticulturae 561: 127-129
- Koltowski Z. (2002) Beekeeping value of recently cultivated winter rapeseed cultivars. Journal of Apicultural Science 46: 23-33
- Koltowski Z. (2005) The effect of pollinating insects on the yeld of winter rapeseed (Brassica napus L. var. napus F. biennis) cultivars. Journal of Apicultural Science 49: 29-41
- Koltowski Z. (2007) Degree of utilization of potential sugar yield of a rapeseed plantation by insects in respect of rapeseed honey yield in an apiary. Journal of Apicultural Science 51: 67-79
- Langridge D.F., Goodman R.D. (1982) Honeybee pollination of oilseed rape, cultivar Midas. Australian Journal of Experimental Agriculture and Animal Husbandry 22: 124-126
- Marsaro-Júnior A.L., Halinski R., Blochtein B. et al. (2017) Diversidade de abelhas na cultura da canola no Rio Grande do Sul. Documentos online n. 168, Embrapa Trigo. Passo Fundo, RS: Embrapa Trigo, 22p., Brazil
- Mesquida J., Renard M. (1979) Entomophilous pollination of male-sterile strains of winter rapeseed (*Brassica napus* L. Metzger var. *oleifera*) and a preliminary study of alternating devices. Proceedings of the IVth International Symposium on Pollination, Md. Agric. Exp. Sta. Spec. Misc. Publ. 1: 49-57
- Mesquida J., Renard M. (1981) Pollinisation du colza d'hiver male fertile et male sterile (*Brassica napus* L var *oleifera* Metzger) par l'abeille domestique (*Apis mellifera* L) Effets sur la phenologie et le rendement. Apidologie 12: 345-362
- Mesquida J., Renard M., Pierre J.S. (1988) Rapeseed (*Brassica napus* L.) productivity: The effect of honeybees (*Apis mellifera* L.) and different pollination conditions in cage and field tests. Apidologie 19: 51-72
- Oliveira, R.H. (2017) Polinizadores de canola: perspectivas para o manejo sustentável de insetos, produtividade de grãos e mudanças climáticas. Thesis (Doctorate in Zoology) Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, 149p., Brazil
- Oz M., Karasu A., Cakmak I. et al. (2008) Effect of honeybees pollination on seed setting, yield and quality characteristics of rapeseed (*Brassica napus oleifera*) Indian Journal of Agricultural Sciences 78: 680-683
- Picard-Nizou A.L., Pham-Delegue M.H., Kerguelen V. et al. (1995) Foraging behaviour of honey bees (Apis mellifera L.) on transgenic oilseed rape (Brassica napus L. var. oleifera) Transgenic Research 4: 270-276
- Pierre J. (2001) The role of honeybees (*Apis mellifera*) and other insect pollinators in gene flow between oilseed rape (*Brassica napus*) and wild radish (*Raphanus raphanistrum*) Acta Horticulturae 561: 47-51.
- Pierre J., Valssiere B., Vallee P., Renard M. (2010) Efficiency of airborne pollen released by honeybee foraging on pollination in oilseed rape: a wind insect-assisted pollination. Apidologie 41: 109-115
- Sabbahi R., Oliveira D., Marceau J. (2006) Does the honeybee (Hymenoptera: Apidae) reduce the blooming period of canola? Journal of Agronomy and Crop Sciences 192: 233-237
- Shakeel M., Inayatullah M., Ali H. (2015) Checklist of insect pollinators and their relative abundance on two canola (*Brassica napus*) cultivars In Peshawar, Pakistan. Journal of Entomology and Zoology Studies 3: 326-330
- Steffan-Dewenter I. (2003) Seed set of male-sterile and male-fertile oilseed rape (Brassica napus) in relation to pollinator density. Apidologie 34: 227-235
- Williams I.H. (1985) The Pollination of swede rape (Brassica napus L.) Bee World 66: 16-22

- Williams I.H., Simpkins J.R. (1989) Honeybee pollination of the double low oilseed rape cultivar ariana. Aspects of Applied Biology 23: 343-346
- Witter S., Nunes-Silva P., Blochtein B. (2014) Abelhas na polinização da canola: benefícios ambientais e econômicos. Porto Alegre: EDIPUCRS, p.71, Brazil

Mainly crop pollination and production:

- Abrol D.P., Shankar U. (2012) Pollination in oil crops: recent advances and future strategies. In: Technological innovations in major world oil crops, vol. 2, New York: Springer, New York, p.221-267, USA
- Chambó E.D., Oliveira N.T.E., Garcia R.C. et al. (2014) Pollination of rapeseed (*Brassica napus*) by africanized honeybees (Hymenoptera: Apidae) on two sowing dates. Anais da Academia Brasileira de Ciências 86: 2087-2100
- Hoyle M., Hayter K., Cresswell J.E. (2007) Effect of pollinator abundance on self-fertilization and gene flow: application to GM canola. Ecological Application 17: 2123-2135
- Mesquida J., Renard M., Pierre J. (1988) Rapeseed (*Brassica napus* L.) productivity: the effect of honeybees (*Apis mellifera* L.) and different pollination conditions in cage and field tests. Apidologie 19: 51-72
- Tomm G.O. (2013) Situação atual e perspectivas da canola no Brasil. Empraba Trigo 58: 105-113

Cashew/Caju

Mainly pollinators and flower visitors:

- Bhattacharya A. (2004) Flower visitors and fruit set of *Anacardium occidentale*. Annales Botanici Fennici 41: 385-392
- Eradasappa E., Mohana G.S. (2016) Role of pollination in improving productivity of cashew A review. Agricultural Reviews 37: 61-65
- Freitas, B.M. (2018) Cashew pollination: answering practical questions. In: The pollination of cultivated plants: a compendium for practitioners. Vol. 1, 2nd edition. FAO: Rome. p.280-287, Italy
- Freitas B.M., Pacheco Filho A.J.S., Andrade P.B. et al. (2014a) Forest remnants enhance wild pollinator visits to cashew flowers and mitigate pollination deficit in NE Brazil. Journal of Pollination Ecology 12: 22-30
- Freitas B.M., Paxton R.J. (1996) The role of wind and insects in cashew (*Anacardium occidentale*) pollination in NE Brazil. The Journal of Agricultural Science 126: 319-326
- Freitas B.M., Paxton R.J. (1998) The comparison of two pollinators: the introduced honey bee *Apis mellifera* and an indegenous bee *Centris tarsata* on cashew *Anacardium occidentale* in its native range of NE Brasil. Journal of Applied Ecology 35: 109-121
- Freitas B.M., Paxton R.J., Holanda Neto J.P. (2002) Identifying pollinators among an array of flower visitors, and the case of inadequate cashew pollination in NE Brazil. Pollinating Bees The Conservation Link Between Agriculture and Nature Ministry of Environment, Brasilia: p.229-244
- Freitas B.M., Pereira J.O.P. (2004) Crop consortium to improve pollination: can West Indian cherry (Malpighia emarginata) attract Centris bees to pollinate cashew (Anacardium occidentale)?. In: Solitary bees: conservation, rearing and management for pollination. 1ed. Fortaleza: Imprensa Universitária da UFC. p.193-201
- Freitas B.M., Silva C.I., Lemos C.Q., Rocha E.E.M., Mendonça K.S., Pereira N.O. (2014b) Plano de manejo para polinização da cultura do cajueiro: conservação e manejo de polinizadores para agricultura sustentável, através de uma abordagem ecossistêmica. Rio de Janeiro: Funbio. 52p.
- Heard T.A., Vithanage V., Chacko E.K. (1990) Pollination biology of cashew in the Northern Territory of Australia. Crop and Pasture Science 41: 1101-1114
- Holanda Neto J.P., Freitas B.M., Bueno D.M., de Araujo Z.B. (2002) Low seed/nut productivity in cashew (Anacardium occidentale): Effects of self-incompatibility and honey bee (Apis mellifera) foraging behaviour. Journal of Horticultural Science and Biotechnology 77: 226-231

Castor bean/Mamona

Mainly pollinators and flower visitors:

Rizzardo R.A.G. (2007) O papel de *Apis mellifera* L. como polinizador da mamoneira (*Ricinus communis* L.): avaliação de eficiência de polinização das abelhas e incremento de produtividade da cultura. Dissertation (Master) – Universidade Federal do Ceará, Fortaleza, 78p., Brazil

Mainly crop pollination and production:

- Giannini T.C., Cordeiro G.D., Freitas B.M. et al. (2015) The dependence of crops for pollinators and the economic value of pollination in Brazil. Journal of Economic Entomology 108: 849–857
- Rizzardo R.A., Milfont, M.O., Silva E.M.S., Freitas B.M. (2012) Apis mellifera pollination improves agronomic productivity of anemophilous castor bean (*Ricinus communis*) Anais da Academia Brasileira de Ciências 84: 1137-1145

Citrus/Frutas cítricas

Mainly pollinators and flower visitors:

- Grajales-Conesa J., Meléndez-Ramírez V., Leopoldo C.-L., Sánchez D. (2013) Native bees in blooming orange (*Citrus sinensis*) and lemon (*C. limon*) orchards in Yucatán, Mexico. Acta Zoológica Mexicana (nueva serie) 29: 437–440
- Malerbo-Souza D.T., Nogueira-Couto R.H., Couto L.A. (2004) Honey bee attractants and pollination in sweet orange, *Citrus sinensis* (L.) Osbeck, var. Pera-Rio. Journal of Venomous Animals and Toxins including Tropical Diseases 10: 144-153
- Ribeiro G.S., Alves E.M., Carvalho C.A.L. (2017) Biology of pollination of *Citrus sinensis* variety 'Pera Rio'. Revista Brasileira de Fruticultura, 39: e-033
- Toledo V.A.A., Halak A.L., Chambó E.D. et al. (2013) Polinização por abelhas (*Apis mellifera* L.) em laranjeira (*Citrus sinensis* L. Osbeck) Scientia Agraria Paranaensis 12: 236-246

Cocoa/Cacau

Mainly pollinators and flower visitors:

- Chumacero de Schawe C., Kessler M., Hensen I., Tscharntke T. (2016) Abundance and diversity of flower visitors on wild and cultivated cacao (*Theobroma cacao* L.) in Bolivia. Agroforestry Systems 92: 117-125
- Erickson E.H., Young A.M., Erickson B.J. (1988) Pollen collection by honeybees (Hymenoptera: Apidae) in a Costa Rican cacao (*Theobroma cacao*) plantation. Journal of Apicultural Research 27: 190-196
- Soria S.J. (1981) Insetos polinizadores: Forcipomya metodos para aumentar a polinizacao e seus efeitors sobre a producao. Proceedings 7th International Cacao Conference, Brazil
- Young A.M. (1982) Effects of shade cover and availability of midge breeding sites on pollinating midge populations and fruit set in two cocoa farms. Journal of Applied Ecology 19: 47-63
- Young A.M. (1983) Seasonal differences in abundance and distribution of cocoa-pollinating midges in relation to flowering and fruit set between shaded and sunny habitats of the La Lola Cocoa Farm in Costa Rica. Journal of Applied Ecology 20: 801-823

Mainly crop pollination and production:

- Adjaloo M.K., Oduro W. (2013) Insect assemblage and the pollination system in cocoa ecosystems. Journal of Applied Biosciences 62: 4582-4594
- Frimpong E.A. Gordon I., Kwapong P.K., Gemmill-Herren B. (2009) Dynamics of cocoa pollination: tools and applications for surveying and monitoring cocoa pollinators. International Journal of Tropical Insect Science 29: 62–69
- Groeneveld J.H., Tscharntke T., Moser G., Clough Y. (2010) Experimental evidence for stronger cacao yield limitation by pollination than by plant resources. Perspectives in Plant Ecology, Evolution and Systematics. 12: 183-191
- Lemos C. Q. (2014) Abelha *Plebeia* cf. *flavocincta* como potencial polinizador do cacaueiro (*Theobroma cacao* L.) no semiárido brasileiro. Master Dissertation. Universidade Federal do Ceará.71p., Brazil
- Lopes U.V., Monteiro W.S., Pires J.L. et al. (2011) Cacao breeding in Bahia, Brazil: strategies and results. Crop Breeding and Applied Biotechnology 11: 73-81
- Santarém M.C.A., Felippe-Bauer M.L. (2016) Brazilian species of biting midges espécies de maruins do Brasil (Diptera: Ceratopogonidae) Fiocruz: Rio de Janeiro. 67p., Brazil
- Silva C.R.S., Albuquerque P.S.B., Ervedosa F.R. et al. (2011) Understanding the genetic diversity, spatial genetic structure and mating system at the hierarchical levels of fruits and individuals of a continuous *Theobroma cacao* population from the Brazilian Amazon. Heredity 106: 973-985

Coconut/Coco

Mainly pollinators and flower visitors:

- Castro S.M. (2002) Bee fauna of some tropical and exotic fruits: potencial pollinators and their conservation. In: Pollinating bees - the conservation link between agriculture and nature - Ministry of Environment, p.275-288, Brazil
- Free J.B., Raw A. Williams I. H. (1975) Pollination of coconut (*Cocos nucifera* L.) In Jamaica by honeybees and wasps. Applied Animal Ethology 1: 213-223
- Hedström, I. (1986) Pollen carriers of *Cocos nucifera* L. (Palmae) in Costa Rica and Ecuador (Neo tropical region) Revista de Biología Tropica 34: 297-301

Mainly crop pollination and production:

- Benassi A.C., Fanton C.S., Santana E.N. (2013) O cultivo do coqueiro-anão-verde: tecnologias de produção. Vitória ; Incaper. 120p., Brazil
- Conceição E.S., Delabie J.H.C., Neto A. (2004) A entomofilia do coqueiro em questão: Avaliação do transporte de pólen por formigas e abelhas nas inflorescências. Neotropical Entomology 33: 679-683
- Melendez-Ramirez V., Parra-Tabla V., Kevan P.G. et al. (2004) Mixed mating strategies and pollination by insects and wind in coconut palm (*Cocos nucifera* L. (Arecaceae)): importance in production and selection. Agricultural and Forest Entomology 6: 155-163
- Regi J.T., Josephrajkumar A. (2013) Flowering and pollination biology in coconut. Journal of Plantation Crops, 41: 109-117

Coffee/Café

Mainly pollinators and flower visitors:

- Badano E.I., Vergara C.H. (2011) Potential negative effects of exotic honey bees on the diversity of native pollinators and yield of highland coffee plantations. Agricultural and Forest Entomology 13: 365-372
- Brokaw J. (2013) Pollinator habitat availability and diversity in various tropical agroforestry management systems of *Coffea arabica* in Santa Clara, Chiriqui. Independent Study Project (ISP) Collection, Panama
- Hipolito J., Boscolo D., Viana B.F. (2018) Landscape and crop management strategies to conserve pollination services and increase yields in tropical coffee farms. Agriculture, Ecosystem and Environment 256: 218-225
- Hutchinson J.L. (2012) Impact of honeybees on coffee pollination in Jamaica, West Indies. Master thesis, Humboldt State University, CA, USA
- Philpott S.M., Uno S., Maldonado J. (2006) The importance of ants and high-shade management to coffee pollination and fruit weight in Chiapas, Mexico. Biodiversity and Conservation 15: 473-487
- Roubik D.W. (2002) Feral african bees augment neotropical coffee yield. In: Pollinating Bees The Conservation Link Between Agriculture and Nature Ministry of Environment p.255-266, Brazil
- Tarno H., Wicaksono K.P., Begliomini E. (2018) Floral stimulation and behavior of insect pollinators affected by pyraclostrobin on Arabica coffee. Journal of Agricultural Science 40: 161-167
- Veddeler D., Olschewski R., Tscharntke T., Klein A.M. (2008) The contribution of non-managed social bees to coffee production: new economic insights based on farm-scale yield data. Agroforestry Systems 73: 109-114
- Vergara C.H., Badano E.I. (2009) Pollinator diversity increases fruit production in Mexican coffee plantations: The importance of rustic management systems. Agriculture, Ecosystem and Environment 129: 117-123

Mainly crop pollination and production:

- Klein A.M., Steffan-Dewenter I., Tscharntke T. (2003a) Fruit set of highland coffee increases with the diversity of pollinating bees. Proceedings of the Royal Society of London. Series B: Biological Sciences 270: 955–961
- Klein A.-M., Steffan-Dewenter I., Tscharntke T. (2003b) Bee pollination and fruit set of *Coffea arabica* and *C. canephora* (Rubiaceae) American Journal of Botany 90: 153-157
- Mesquita, C.M. et al. (2016) Manual do café: colheita e preparo (*Coffea arabica* L.) Belo Horizonte: EMATER-MG, 52p., Brazil

Common bean/Feijão

Mainly pollinators and flower visitors:

- Carpentieri-Pípolo V., Vizoni E., Giroto J.C.M. (2001) Determinação do melhor período para realização de cruzamento artificial em feijão-vagem, *Phaseolus vulgaris* L., em Londrina, Estado do Paraná. Acta Scientiarum 23: 1191-1193
- Free J.B. (1966) The pollination of the beans *Phaseolus multiflorus* and *Phaseolus vulgaris* by honeybees. Journal of Apicultural Research 5: 8791
- IbarraPerez FJ., Barnhart D., Ehdaie B. et al. (1999) Effects of insect tripping on seed yield of common bean. Crop Science 39: 428433
- Shree P. S, Gepts P., Debouck D.G. (1991) Races of common bean (*Phaseolus vulgaris*, Fabaceae) Economic Botany 45: 379-396

Cotton/Algodão

Mainly pollinators and flower visitors:

- Bozbek T., Ozbek N., Sezener V. et al. (2008) Natural crossing and isolation distance between cotton genotypes in Turkey. Scientia Agricola 65: 314
- Cusser S., Neff J.L., Jha S. (2016) Natural land cover drives pollinator abundance and richness, leading to reductions in pollen limitation in cotton agroecosystems. Agriculture, Ecosystem and Environment 226: 33-42
- Eisikowitch D., Loper G.M. (1984) Some aspects of flower biology and bee activity on hybrid cotton in Arizona, USA. Journal of Apicultural Research 23: 243-248
- Heuberger S., Ellers-Kirk C., Tabashnik B.E., Carrière Y. (2010) Pollen- and seed-mediated transgene flow in commercial cotton seed production fields. PLoS ONE 5: e14128
- McGregor S.E. (1959) Cotton-flower visitation and pollen distribution by honey bees. Science 129: 97-98
- Moffett J.O. (1977) Producing hybrid cotton seed on the high plains of Texas. Proceedings 1977 Beltwide Cotton Production Research Conferences, USA
- Moffett J.O., Stith L.S., Burkhart C.C., Shipman C.W. (1975) Honey bee visits to cotton flowers. Environmental Entomology 4: 203-206
- Moffett J.O., Stith L.S., Morton H.L., Shipman C.W. (1980) Effect of 2,4-d on cotton yield, floral nectar, seed germination, and honeybee visits. Crop Science 20: 747-750
- Pires C. S. S., Pires V.C., Rodrigues W. et al. (2015) Plano de manejo para polinizadores em áreas de algodoeiro consorciado no Nordeste do Brasil. Rio de Janeiro: Funbio. 40p., Brazil
- Pires C.S.S., Silveira F.A., Cardoso C.F. et al. (2014) Selection of bee species for environmental risk assessment of gm cotton in the Brazilian Cerrado. Pesquisa Agropecuária Brasileira
- Rhodes J. (2002) Cotton pollination by honey bees. Australian Journal of Experimental Agriculture 42: 513-518
- Waller G.D., Moffett J.O., Loper G.M., Martin J.H. (1985a) An evaluation of honey bee foraging activity and pollination efficacy for male-sterile cotton. Crop Science 25: 211-214
- Waller G.D., Vaissiere B.E., Moffett J.O., Martin J.H. (1985b) Comparison of carpenter bees (*Xylocopa varipuncta* Patton) (Hymenoptera: Anthophoridae) and honey bees (*Apis mellifera* L.) (Hymenoptera: Apidae) as pollinators of male-sterile cotton in cages. Journal of Economic Entomology 78: 558-561

Mainly crop pollination and production:

FAO (2018) The pollination of cultivated plants a compendium for practitioners Volume 1. ISBN 978-92-5-130512-6

Cowpea/Feijão-caupi

Mainly pollinators and flower visitors:

Araújo, F.W.S. (2012) Abelhas mamangavas (*Xylocopa cearensis* e *Xylocopa griscescens*) como potenciais polinizadores do feijão caupi (*Vigna unguiculata*) M.Sc. Dissertation - Universidade Federal do Ceará, Centro de Ciências Agrárias, Programa de Pós-Graduação em Zootecnia, Fortaleza, Brazil

Asiwe J.A.N. (2009) Insect mediated outcrossing and geneflow in cowpea (Vigna unguiculata (L:) Walp):
Implication for seed prodiction and provision of containment structures for genetically transformed cowpea. African Journal of Biotechnology 8: 226-230

- Fohouo F.T., Ngakou A., Kengni B.S. (2009) Pollination and yield responses of cowpea (*Vigna unguiculata* L. Walp.) to the foraging activity of *Apis mellifera adansonii* (Hymenoptera:Apidae) at Ngaoundere (Cameroon) African Journal of Biotechnology 8: 1988-1996
- Ige O.E., Olotuah O.F., Akerele V. (2011) Floral biology and pollination ecology of cowpea (*Vigna unguiculata* L. Walp) Modern Applied Science 5: 74-82
- Sheahan, C.M. (2012) Plant guide for cowpea (*Vigna unguiculata*) USDA-Natural Resources Conservation Service, Cape May Plant Materials Center, Cape May, NJ.
- Venter, H.M. (1996) Difficulties with cross-pollinating five cowpea lines: technique development. In: The Biodiversity of African Plants. p.656-660, Springer, Dordrecht, Netherlands

Mainly crop pollination and production:

D'Andrea A.C., Kahlheber S., Logan A.L., Watson D.J. (2007) Early domesticated cowpea (*Vigna unguiculata*) from Central Ghana. Antiquity 81: 686-698

Cupuaçu/Cupuaçu

Mainly pollinators and flower visitors:

- Calzavara B.B.G., Muller C.H., Kahwage O.N.N. (1984) Fruticultura tropical: o cupuaçuzeiro cultivo, beneficiamento e utilização do fruto. EMBRAPA-CPATU: Belém, 101p., Brazil
- Falcao M.A., Lleras E. (1983) Aspectos fenológicos, ecológicos e de produtividade do Cupuaçu *Theobroma* grandiflorum (Willd ex Spreng) Schum. Acta Amazonica 13: 725-735
- Maués M.M., Souza L.A. de, Miyanaga R. (2000) Insetos polinizadores do cupuaçuzeiro (*Theobrama grandiflorum* Willd. Ex . Sprengel) no Estado da Pará, Brasil. Belém: Embrapa Amazonia Oriental, 19p. (Embrapa Amazonia Oriental. Circular Técnica, 12), Brazil
- Venturieri G.A., Ribeiro Filho A.A. (1995) A polinização manual do Cupuaçuzeiro (*Theobroma grandiflorum*) Acta Amazonica 25: 181-192

Mainly crop pollination and production:

- Alves RM, Artero AS, Sebbenn AM, Figueira A (2003) Mating system in a natural population of *Theobroma grandiflorum* (Willd. ex Spreng.) Schum., by microsatellite markers. Genetics and Molecular Biology 26: 373-379.
- Venturieri, G.A. (1993) Cupuaçu: a espécie, sua cultura, usos e processamentos. Belém: Clube do Cupu, 108p., Brazil
- Venturieri, G.A. (1994) Floral biology of cupuassu [*Theobroma grandiflorum* (Willdenow ex. Sprengel Schumann)], Ph.D Thesis. University of Reading, Reading, 211p., UK
- Venturieri, G.A. (2011) Flowering levels, harvest season and yields of cupuassu (Theobroma grandiflorum), Acta Amazonica 41: 143-152

Guava/Goiaba

- Alves J.E., Freitas B.M. (2006) Comportamento de pastejo e eficiência de polinização de cinco espécies de abelhas em flores de goiabeira (*Psidium guajava* L.) Revista Ciência Agronômica 37: 216-220.
- Alves J.E., Freitas B.M. (2007) Requerimentos de polinização da goiabeira. Ciência Rural 37: 1281-1286.
- Hedström I. (1988) Pollen carriers and fruit development of *Psidium guaiava* L. (Myrtaceae) in the neotropical region. Revista de Biologia Tropical 36: 551-553
- Siqueira K.M.M., Kiill L.H.P., Martins C.F., Silva L.T. (2012) Pollination ecology of *Psidium guajava* L. (Myrtaceae): richness, frequency and time of activities of floral visitors in an agricultural system. Semana Entomológica da Bahia (SINSECTA)
- Tchuenguem Fohouo F.N., Djonwangwe D., Messi J., Brückner D. (2007) Exploitation des fleurs de entada africana, Eucalyptus camaldulensis, *Psidium guajava* et *Tricbillia emetica* Par *Apis mellifera adansonii* a Dang (Ngaoundere, Cameroun) Cameroon Journal of Experimental Biology 3: 50-60
- Viana B.F. (2008) Management plans for fruit crop pollinators in the states of Bahia and Pernambuco, northeastern, Brazil. Pollinators Management in Brazil, p.38-40, Brazil

- Freitas B.M., Alves J.E. (2008) Efeito do número de visitas florais da abelha melífera (*Apis mellifera* L.) na polinização da goiabeira (*Psidium guajava* L.) cv. Paluma. Revista Ciência Agronômica 39: 149-154
- Hamilton R.A., Seagrave-Smith H. (1959) Growing guava for processing extension. Bulletin 63, University of Hawaii, USA

Kiwifruit/Kiwi

Mainly pollinators and flower visitors:

- Costa G., Testolin R., Vizzotto G. (1993) Kiwifruit pollination: an unbiased estimate of wind and bee contribution. New Zealand Journal of Crop and Horticultural Science 21: 189-195
- MacFarlane R.P., Ferguson A.M. (1984) Kiwifruit pollination: a survey of the insect pollinators in New Zealand. Les Colloques de l'INRA 21, France
- Miñarro M., Twizell K.W. (2015) Pollination services provided by wild insects to kiwifruit (Actinidia deliciosa) Apidologie 46: 276-285
- Palmer-Jones T., Clinch P.G. (1974) Observations on the pollination of Chinese gooseberries variety 'Hayward'. New Zealand Journal of Experimental Agriculture 2: 455-458
- Palmer-Jones T., Clinch P.G. (1975) Honey bees essential for kiwifruit pollination. The Orchardist of New Zealand 347p., New Zealand
- Sharma M., Mattu V.K., Thakur M.L. (2013) Pollination studies on kiwi crop (*Actinidia deliciosa* Chev.) in Himachal Pradesh, India. International Journal of Advanced Biological Research 3: 545-548
- Testolin R., Vizzotto G., Costa G. (1991) Kiwifruit pollination by wind and insects in Italy. New Zealand Journal of Crop and Horticultural Science 19: 381-384
- Vassiere B.E., Rodet G., Cousin M. et al. (1996) Pollination effectiveness of honey bees (Hymenoptera: Apidae) in a kiwifruit orchard. Journal of Economic Entomology 89: 453-461

Mainly crop pollination and production:

Anonymous (1984) Kiwifruit pollination. The New Zealand Beekeeper, New Zealand

- Blanchet P., Douault P.H., Pouvreau A. (1991) Kiwifruit (*Actinidia deliciosa* Chev.) pollination: honey-bee behaviour and its influence on the fruit. Acta Horticulturae 282: 105-110
- Clinch P.G. (1984) Kiwifruit pollination by honey bees 1. Tauranga observations, 1978-81. New Zealand Journal of Experimental Agriculture 12: 29-38
- Clinch P.G., Heath A. (1985) Wind and bee pollination research. New Zealand Kiwifruit Journal 15, New Zealand
- Clinch P.G., Houten A.T. (1985) Pollination of pastures and crops kiwifruit. Wallaceville Animal Research Centre Annual Report, New Zealand
- Donovan B.J., Read P.E.C. (1990) Efficacy of honey bees as pollinators of kiwifruit. Acta Horticulturae 288: 220-224
- Gonzalez M.V., Coque M., Herrero M. (1998) Influence of pollination systems on fruit set and fruit quality in kiwifruit (*Actinidia deliciosa*) Annals of Applied Biology 132: 349-355
- Goodwin M., Haine H., Wise G. (1997) How strong should honeybee colonies be for kiwifruit pollination. New Zealand Kiwifruit Journal 20-21
- Howpage D., Spooner-Hart R.N., Vithanage V. (2001) Influence of honey bee (*Apis mellifera*) on kiwifruit pollination and fruit quality under Australian conditions. New Zealand Journal of Crop and Horticultural Science 29: 51-59
- Jay D., Jay C. (1983) Some observations of honeybees in kiwifruit orchards. New Zealand Beekeeper p.21-22
- Manino A., Marletto F., Patetta A., Porporato M. (1996) Researches on the role of honeybees in the fructification of kiwi-fruit in Piedmont. La Selezione Veterinaria 11: 747-754
- Palmer-Jones T., Clinch P.G. (1976) Effect of honey bee saturation on the pollination of Chinese Gooseberries Variety 'Hayward'. New Zealand Journal of Experimental Agriculture 4: 255-256
- Simonetto P.R., Grellmann E.O. (1998) Cultivares de kiwi com potencial de produção na região da Serra do Nordeste do Rio Grande do Sul. Porto Alegre: Fepagro, 19p. (Boletim Técnico, n.7), Brazil

Lychee/Lichia

Mainly pollinators and flower visitors:

- Abrol D.P. (2006) Diversity of pollinating insects visiting litchi flowers (*Litchi chinensis* Sonn.) and path analysis of environmental factors influencing foraging behaviour of four honeybee species. Journal of Apicultural Research 45: 180-187
- Ali S., Shehzad A., Rafi M.A., Zia A. (2013) Insect pollinators of litchi (*Litchi chinensis*) from district Haripur, Pakistan Journal of Agricultural Research 26: 220-229
- Mandal B.K., Galib A.J., Sultana N., Das A. (2016) Relationship of urban dust precipitation on pollination and fruit falling of *Mangifera indica* and *Litchi chinensis* in Dhaka District, Bangladesh. Journal of Entomology and Zoology Studies 4: 1185-1191
- Poonam S., Khan M. S., Shiwani B. (2010) Role of insect pollinators for enhancing the litchi fruit production. Haryana Journal of Horticultural Sciences 39: 226-227
- Rai V.L., Srivastava P., Bisht K., Mishra V.K. (2017) Diversity and relative abundance of pollinating insects visiting litchi (*Litchi chinensis* Sonn.) inflorescence under Tarai Agro-Climatic condition. Journal of Experiment Zoology, India 20: 233-239
- Singh J., Kumar N.R., Devi A. (2017) Insect pollinators, their diversity, foraging behaviour and relative abundance on litchi, okra and sarson. Entomon 42: 275-282
- Somnuk B., Suavansri T. (2005) Study on pollinating cultivars for fruit setting of 'Khom' lychee. Acta Horticulturae 665: 111-116
- Srivastava K., Sharma D., Pandey S.D. et al. (2017) Dynamics of climate and pollinator species influencing litchi (*Litchi chinensis*) in India. Indian Journal of Agricultural Sciences 87: 266-269
- Stern R.A., Gazit S. (1996) Lychee pollination by the honeybee. Journal of the American Society for Horticultural Science 121: 152-157

Mainly crop pollination and production:

- Batten D. J., McConchie C. A. (1995) Floral induction in growing buds of lychee (*Litchi chinensis*) and mango (*Manifera indica*) Australian Journal of Plant Physiology 22: 783-91
- Batten D.J. (1986) Towards an understanding of reproductive failure in lychee (*Litchi chinensis*) Acta Horticultura 175: 79-84
- Kuman R. (2014) Planned honey bee pollination in litchi (*Litchi chinensis* Sonn.) a new production strategy for enhancing fruit yield and quality in india. Acta Horticultura 1029: 281-286
- Matos E.H.S. F. (2012) Cultivo de Lichia Centro de Apoio ao Desenvolvimento Tecnológico CDT/UnB, Brazil
- Morton J. (1987) Lychee. In: Fruits of warm climates. Miami, FL. p.249-259, USA

Mango/Manga

- Amin M.R., Namni S., Miah M.R.U. et al. (2015) Insect inventories in a mango-based agroforestry area in Bangladesh: Foraging behavior and performance of pollinators on fruit set. Entomological Research 45: 217-224
- Carvalheiro L.G., Seymour C.L., Veldtman L., Nicolson S.W. (2010) Pollination services decline with distance from natural habitat even in biodiversity-rich areas. Journal of Applied Ecology 47: 810-820
- Fajardo J.R.A.C., Medina J.R., Opina O.S., Cervancia C.R. (2008) Insect pollinators and floral visitors of mango (*Mangifera indica* L. cv. Carabao) The Philippine Agricultural Scientist 91: 372-382
- Kiill L.H.P. (2008) Assessment of mango (Mangifera indica L., Anarcadiaceae) and passion fruit (Passiflora edulis F. flavicarpa Deg., Passifloraceae) pollinators in the San Francisco Valley, northeastern Brazil. Pollinators Management in Brazil, Brazil
- Kumari D.A., Madhavi J., Bhagwan A., Kumar M.R. (2014) Surveillance of pollinators and their behaviour in mango flowers. Plant Archives 14: 727-729
- Singh G. (1984) Pollination, pollinators and fruit setting in mango. Acta Horticulturae 455: 116-123
- Sung I., Lin M., Chang C. et al. (2006) Pollinators and their behaviors on mango flowers in souther Taiwan. Formosan Entomologist 26: 161-170

Tayeng M., Gogoi H. (2016) Insect pollinators of crops and fruits in Arunachal Pradesh, Eastern Himalaya: rich diversity in flowers with yellow anther. Proceedings of The Zoological Society 71: 56-62

Mainly crop pollination and production:

- Bally I.S.E. (2006) Mangifera indica (Mango) In: Species profiles for pacific island agroforestry. Permanent Agriculture Resources (PAR), Holualoa, Hawaii, USA
- Gajendra S.D.R. (1989) Insect pollinators of mango and their role in fruit setting. Acta Horticulturae 231: 629-632
- Huda, N. et al. (2015) Pollination services of mango flower pollinators. Journal of Insect Science 15: 113
- Ramírez F, Davenport T.L. (2016) Mango (Mangifera indica L.) pollination: A review. Scientia Horticulturae 203: 158-168

Melon/Melão

Mainly pollinators and flower visitors:

- Goodell K., Thomson J.D. (2007) Influence of bee species (Hymenoptera: Apiformes) with contrasting behaviours on pollen movement in a mustard, *Brassica rapa* (Brassicaceae) and the muskmelon *Cucumis melo* (Cucurbitaceae) Entomologia Generalis 29: 237-251
- Kiill L.H.P., Feitoza E.D.A., Siqueira K.M.M. et al. (2016) Evaluation of floral characteristics of melon hybrids (*Cucumis melo* L.) in pollinator attractiveness. Revista Brasileira de Fruticultura 38: e-531
- Kiill L.H.P., Ribeiro M.F., Siqueira K.M.M., Silva E.M.S. (2015) Plano de manejo de polinizadores do meloeiro. Petrolina: Embrapa Semiárido. 55p., Brazil
- Kiill L.H.P., Siqueira K.M.M., Marcia S.C. et al. (2014) Frequency and foraging behavior of *Apis mellifera* in two melon hybrids in Juazeiro, State of Bahia, Brazil. Anais da Academia Brasileira de Ciências 86: 2049-2055
- Lemasson M. (1987) Interet de l'abeille mellifere (*Apis mellifica*) dans la pollinisation de cultures en serre de cornichon (*Cucumis sativus*), de melon (*Cucumis melo*) et de tomate (*Lycopersicum esculentum*) Revue de L'Agriculture. 40: 915-924
- Meléndez-Ramirez V., Magaña-Rueda S., Parra-Tabla V. et al. (2002) Diversity of native bee visitors of cucurbit crops (Cucurbitaceae) in Yucatán, México. Journal of Insect Conservation 6: 135-147
- Mouzin T.E., Reed D.K., Chaney W.E. (1980) Influence of honey bees on cantaloupe production in Indiana. Proceedings Indiana Academy of Sciences 89: 215-217
- Ribeiro M.F., Silva E.M.S., Oliveira L. et al. (2015) Honey bees (*Apis mellifera*) visiting flowers of yellow melon (*Cucumis melo*) using different number of hives. Ciência Rural, Brazil
- Stanghellini M.S., Schultheis J.R., Ambrose J.T. (2002) Pollen mobilization in selected Cucurbitaceae and the putative effects of pollinator abundance on pollen depletion rates. Journal of the American Society for Horticultural Science 127: 729-736
- Taylor E.A. (1955) Cantaloup production increased with honey bees. Journal of Economic Entomology 48: 327
- Tschoekea P.H., Oliveira E.E., Dalcin S.M. et al. (2015) Diversity and flower-visiting rates of bee species as potential pollinators of melon (*Cucumis melo* L.) in the Brazilian Cerrado Santos. Scientia Horticulturae 186: 207-216
- Williams P. (1987) Pollination of melons. Australian Bee Journal 68: 18-21

- Bomfim I.G.A., Freitas B.M., Aragão, F.A.S., Walters S.A. (2016) Pollination in cucurbit crops. In: Handbook of Cucurbits: Growth, Cultural Practices, and Physiology. Editora UFV: Viçosa MG. p.221-245, Brazil
- Bomfim I.G.A., Bezerra A.D.M., Freitas B.M., Aragão F.A.S. (2019) A polinização do meloeiro. In: Produção de melão. CRC Press. p.181-200, Brazil
- Fernandes N.S., Silva F.A.N., Aragão F.A.S. et al. (2019) Volatile organic compounds role in selective pollinator visits to commercial melon types. Journal of Agricultural Science 11: 93-108
- Gomez S.R., Ornosa C., Selfa J. et al. (2016) Small sweat bees (Hymenoptera: Halictidae) as potential major pollinators of melon (*Cucumis melo*) in the Mediterranean. Entomological Science 19: 55-66
- Grewal G.S., Sidhu A.S. (1978) Insect-pollinators of some cucurbits in Punjab. Indian Journal of Agricultural

Sciences 48: 79-83

- Kaziev T.I., Seidova (1965) La production de nectar des fleurs de certaines especes de cucurbitacees dans les conditions de l'Azerbaidjan. Proceedings 20th International Beekeeping Congress Apimondia, Bukarest p.388-391, Romania
- Mann L.K. (1954) Fruit set in melon breeding: Hand pollination found to be less effective than pollination by honeybees in experiments at Davis. California Agriculture 8: 3.
- McGregor S.E., Levin M.D., Foster R.E. (1965) Honey bee visitors and fruit set of cantaloups. Journal of Economic Entomology 58: 968-970
- McGregor S.E., Todd F.E. (1952) Cantaloup production with honey bees. Journal of Economic Entomology 45: 43-47
- Nerson H. (2009) Effects of pollen-load on fruit yield, seed production and germination in melons, cucumbers and squash. Journal of Horticultural Science and Biotechnology 84: 560-566
- Wilson M.E., Skinner J.A., Wszelaki A.L., Drummond F. (2016) Using nonmetric multidimensional scaling to analyze bee visitation in East Tennessee crops as an indicator of pollination services provided by honey bees (*Apis mellifera* L.) and native bees. Environmental Entomology 45: 390-396

Oil palm/Dendê

Mainly pollinators and flower visitors:

- Barcelos E., Rios S.A., Cunha R.N.V. et al. (2015) Oil palm natural diversity and the potential for yield improvement. Frontiers in Plant Science 6: 1-16
- Meléndez M.R., Ponce W.P. (2016) Pollination in the oil palms *Elaeis guineensis, E. oleifera* and their hybrids (OxG), in tropical America. Pesquisa Agropecuária Tropical 46: 102-110
- Moura J.I.L., Cividanes F.J., Santos Filho L.P., Valle R.R. (2008) Polinização do dendezeiro por besouros no sul da Bahia. Pesquisa Agropecuária Brasileira 43: 289-294.

Mainly crop pollination and production:

- Mayfield M. (2005) The importance of nearby forest to known and potential pollinators of oil palm (*Elaeis guineënsis* Jacq.; Areceaceae) in Southern Costa Rica. Economic Botany 59: 190-196
- Siregar E.H., Atmowidi T., Kahono S. (2016) Diversity and abundance of insect pollinators in different agricultural lands in Jambi, Sumatera. HAYATI Journal of Biosciences 23: 13-17

Okra/Quiabo

Mainly pollinators and flower visitors:

- Azo'o Ela,M., Fohouo F.N.T., Messi J. (2011) Influence of the foraging activity of the entomofauna on okra (*Abelmoschus esculentus*) seed yield. International Journal of Agriculture and Biology 13: 761-765
- Njoya M.T., Wittmann D., Schindler M. (2005) Effect of bee pollination on seed set and nutrition on okra (*Abelmoschus esculentus*) in Cameroon. Deutscher Tropentag, Oct.11-13, Stuttgart-Hohenheim, Germany
- Purewal S.S., Randhawa G.S. (1947) Studies in *Hibiscus esculentus*. Indian Journal of Agricultural Science 17: 129-136
- Singh J., Kumar N.R., Devi A. (2017) Insect pollinators, their diversity, foraging behaviour and relative abundance on litchi, okra and sarson. Entomon 42: 275-282

- Al-Ghzawi A.A.M., Zaittoun S.T., Makadehm I., Al-Tawaha A.R.M. (2003) The impact of wild bees on the pollination of eight okra genotypes under semi-arid mediterranean conditions. International Journal of Agriculture & Biology 5: 408-410
- Ige O.E., Eludire M.O. (2014) Floral biology and pollination ecology of okra (*Abelmoschus esculentus* L. Moench) American International Journal of Biology 2: 1-9
- Malerbo-Souza D.T., Toledo V.A.A., Stuchi A.C. et al. (2001) Estudo sobre a polinização do quiabeiro, *Abelmoschus esculentus* (L.) Moench. Acta Scientiarum Maringá, 23: 1281-1285
- Shalaby G.J. (1998) Natural cross-pollination in okra. Journal on Agriculture Science 3: 381-386

Papaya/Papaia

Mainly pollinators and flower visitors:

Allan P. (1963) Pollination of pawpaws. Farming In South Africa 8: 13-15

- Dey K., Mondal S., Mandal S. (2016) Flower-visitor diversity with reference to pollen dispersal and pollination of *Carica papaya* L. International Journal of Advanced Research 3: 65-71
- Garrett A. (1995) The pollination biology of papaw (Carica papaya L.) in Central Queensland. PhD Thesis, Central Queensland University, USA
- Marin-Acosta J.C. (1969) Insects living on the papaya tree (*Carica papaya*) in Venezuela. Agronomia Tropical (Maracay), Venezuela

Mainly crop pollination and production:

- Avila Jr. R.S., Oliveira R., Pinto C.E. et al. (2012) Relação entre esfingídeos (Lepidoptera, Sphingidae) e flores no Brasil - Panorama e perspectivas de uso de polinizadores. In: Polinizadores no Brasil. São Paulo, Edusp, São Paulo. p.143-152, Brazil
- Damasceno Junior P.C., Pereira T.N.S., Pereira M.G. et al. (2009) Preferential reproduction mode of hermaphrodite papaya plant (*Carica papaya* L; Caricaceae) Revista Brasileira de Fruticultura 31: 182-189
- Martins D.J., Johnson S.D. (2009) Distance and quality of natural habitat influence hawkmoth pollination of cultivated papaya. International Journal of Tropical Insect Science 29: 114-123
- Morrisen, A. (1995) The pollination biology of papaw (*Carica papaya* L.) in central Queensland. PhD Thesis. Central Queensland University, Australia

Passionfruit/Maracujá

Mainly pollinators and flower visitors:

- Baran T.B.M., Mouga D.M.D.S., Pinheiro P.C. et al. (2017) Determination of the diversity and abundance of pollinators (Hymenoptera, Apidae) of yellow passion fruit (*Passiflora edulis* F. Flavicarpa) in southern Brazil. International Journal of Current Research 9: 49126-49134
- Bezerra A.D.M., Pacheco Filho A.S.J., Bomfim I.G.A. et al. (2019) Agricultural area losses and pollinator mismatch due to climate changes endanger passion fruit production in the Neotropics. Agricultural Systems 169: 49-57
- Gaglianone M.C. (2008) Pollinators of yellow passion fruit (*Passiflora edulis* Sims, Passifloraceae) and management of carpenter bees in southeastern Brazil. In: Pollinators Management in Brazil, p.34-35, Ministério do Meio Ambiente, Brazil
- Kiill L.H.P. (2008) Assessment of mango (Mangifera indica L., Anarcadiaceae) and passion fruit (Passiflora edulis F. flavicarpa Deg., Passifloraceae) pollinators in the San Francisco Valley, northeastern Brazil. In: Pollinators Management in Brazil, p.36-37, Ministério do Meio Ambiente, Brazil
- Oliveira P.E. (2008) Sustainable management of pollinators for passion fruit (*Passiflora edulis* F. *flavicarpa* Deg., Passifloraceae) production in central Brazil. In: Pollinators Management in Brazil, p.30-33, Ministério do Meio Ambiente, Brazil
- Siqueira, K.M.M., Kiill, L.H.P., Martins, C.F., et al. (2009) Ecologia da polinização do maracujá-amarelo, na região do Vale do Submédio São Francisco [Ecology of pollination of yellow passion fruit in the region of the Submédio São Francisco Valley]. Revista Brasileira de Fruticultura, 31: 1-12.
- Yamamoto M., Silva C.I., Augusto S.C. et al. (2012) The role of bee diversity in pollination and fruit set of yellow passion fruit (*Passiflora edulis* Forma *flavicarpa*, Passifloraceae) crop in central Brazil. Apidologie 43: 515-526

- Freitas B.M., Imperatriz-Fonseca V.L., Medina L.M. et al. (2009) Diversity, threats and conservation of native bees in the Neotropics. Apidologie 40: 332-346
- Freitas B.M., Oliveira-Filho J.H. (2001) Criação racional de mamangavas: para polinização em áreas agrícolas. Fortaleza: Banco do Nordeste. 96p., Brazil
- Freitas B.M., Oliveira Filho J.H. (2003) Ninhos racionais para mamangava (*Xylocopa frontalis*) na polinização do maracujá-amarelo (*Passiflora edulis*) Ciência Rural 33: 1135-1139
- Hoffmann M., Pereira T.N.S., Mercadante M.B., Gomes A.R. (2000) Polinização de Passiflora edulis f. flavicarpa (Passiflorales, Passifloraceae), por abelhas (Hymenoptera, Anthophoridae) em Campos dos Goytacazes,

Rio de Janeiro. Iheringia. Série Zoologia, p.149-152, Brazil

- Silva C.I., Freitas B.M. (2018) Rearing carpenter bees (*Xylocopa* spp.) for crop pollination: a case study with passionfruit (*Passiflora edulis*) In: The pollination of cultivated plants: a compendium for practitioners. Vol. 2. Rome: FAO. p.89-100, Italy
- Silveira M.V., Abot A.R., Nascimento J.N. et al. (2012) Is manual pollination of yellow passion fruit completely dispensable? Scientia Horticulturae 146: 99-103

Peach/Nectarina

Mainly pollinators and flower visitors:

- Mota M.O.S., Nogueira-Couto R.H. (2002) Polinização entomófila em pessegueiro (*Prunus persica* L.) Brazilian Journal of Veterinary Research and Animal Science 39: 124-128
- Raj H., Mattu V.K. (2014) Diversity and distribution of insect pollinators on various temperate fruit crops in Himachal Himalaya, India. International Journal of Science and Nature 5: 626-631

Mainly crop pollination and production:

Free J.B. (1993) Insect pollinaiton of crops. 2ed. Academic Press: Cardiff, UK

- Gariglio N.F., Mendow M., Weber M.E. et al. (2009) Fenologia e comportamento reprodutivo de pêssego no centro-leste da Argentina. Scientia Agricola 66: 757-763
- Mayer N.A., Bianchi V.J., Feldberg N.P. et al. (2017) Advances in peach, nectarine and plum propagation. Revista Brasileira de Fruticultura 39: e-355
- Weinbaum S.A., Shaw D.V., Muraoka T.T. (1989) Independence of self-compatibility and potentiality for self-pollination in peach x almond hybrids. Euphytica 41: 53-58
- Zhang H., Huang J., Williams P.H. et al. (2015) Managed bumblebees outperform honeybees in increasing peach fruit set in China: different limiting processes with different pollinators. PLoS ONE 10: e0121143

Pear/Pera

Mainly pollinators and flower visitors:

- Hsieh F.K., Chen C.T., Chang C.P., Chang S.Y. (2002) Foraging activities and numerical changes of honeybees on buckwheat, rape and pear. Plant Protection Bulletin 44: 1-13
- Langridge D.F., Jenkins P.T. (1972) A study on pollination of Packham'S Triumph pears. Australian Journal Experimental Agriculture and Animal Husbandry 12: 328-330
- Lewis T., Smith B.D. (1969) The insect faunas of pear and apple orchards and the effect of windbreaks on their distribution. Annals of Applied Biology 64: 11-20
- Maccagnani B., Ladurner E., Santi F., Burgio G. (2003) Osmia cornuta (Hymenoptera, Megachilidae) as a pollinator of pear (*Pyrus communis*): Fruit- and seed-set. Apidologie 34: 207-216
- Mayer D.F., Lunden J.D. (1997) A comparison of commercially managed bumblebees and honey bees (Hymenoptera: Apidae) for pollination of pears. Acta Horticulturae 437: 283-288
- Monzon V.H., Bosch J., Retana J. (2004) Foraging behavior and pollinating effectiveness of *Osmia cornuta* (Hymenoptera: Megachilidae) and *Apis mellifera* (Hymenoptera: Apidae) on "Comice" pear. Apidologie 35: 575-585
- Raj H., Mattu V.K. (2014) Diversity and distribution of insect pollinators on various temperate fruit crops in Himachal Himalaya, India. International Journal of Science and Nature 5: 626-631
- Stern R.A., Goldway M., Zisovich A.H. et al. (2004) Sequential introduction of honeybee colonies increases cross-pollination, fruit-set and yield of 'Spadona' pear (*Pyrus communis* L.) Journal of Horticultural Science and Biotechnology 79: 652-658
- Van den Eijnde J. (1996) Pollination of pear by bumblebees (*Bombus terrestris* L.) and honeybees (*Apis mellifera* L.) Acta Horticulturae 423: 73-78

Mainly crop pollination and production:

Free J.B. (1993) Insect pollination of crops. 2ed. Academic Press: Cardiff, UK

Jacquemart A.L., Michotte-Van der A., Raspe O. (2006) Compatibility and pollinator efficiency tests on *Pyrus communis* L. cv. 'Conference'. Journal of Horticultural Science and Biotechnology 81: 827-830

Onarska A., Masierowska M., Weryszko-Chmielewska E. (2005) The structure of nectaries and nectar

secretion in common pear (Pyrus communis L.) Journal of Apicultural Science 49: 85-92

Quinet M., Warzée M., Vanderplanck M. et al. (2016) Do floral resources influence pollination rates and subsequent fruit set in pear (*Pyrus communis* L.) and apple (*Malus x domestica* Borkh) cultivars? European Journal of Agronomy 77: 59-69

Persimmon/Caqui

Mainly pollinators and flower visitors:

Agustí M., Fonfría M.A. (2010) Fruticultura. Mundi-Prensa Libros, Madrid, Spain

- Giannini T.C., Boff S., Cordeiro, G.D. et al. (2015) Crop pollinators in Brazil: a review of reported interactions. Apidologie 46: 209-223
- Martins F.P., Pereira F.M. (1989) Cultura do caquizeiro. Jaboticabal: Funep, 71p., Brazil

Mainly crop pollination and production:

- Campos S.S., Wittmann M.T.S., Schwarz S.F., Veit P.A. (2015) Biologia floral e viabilidade de pólen em cultivares de caquizeiro (*Diospyros kaki* L.) e *Diospyros virginiana* L.. Revista Brasileira de Fruticultura. Jaboticabal 37: 685-691
- Chauhan N., Thakur B., Sharma G. et al. (2017) Pollination studies in relation to fruit drop in persimmon (*Diospyros kakiv* L.) cv. Hachiya. International Journal of Current Micromiology and Applied Sciences 6: 673-680
- Free J.B. (1993) Insect pollination of crops. Chapter 55, Academic Press: Cardiff, UK
- Gould H.P. (1940) Oriental Persimmons. Leaflet, United States Department of Agriculture 194p., USA
- Hodgson R.W. (1938) Girdling to reduce fruit drop in the Hachiya persimmons. American Society of Horticultural Science Proceedings 36: 405-409
- Hodgson R.W. (1939) Floral situation, sex condition and parthenocarpy in the oriental persimmons. American Society of Horticultural Science Proceedings 37: 250-252
- McGregor S.E. (1976) Insect pollination of cultivated crop plants. Agricultural Research Service, US Department of Agriculture Washington, USA
- Neuwald D.A., Saquet A.A., Sestari I., Sautter C.K. (2009) Persimmon production and commercialization in Brazil: An overview. Acta Horticulturae 833: 51-56
- Popenoe W. (1924) Manual of tropical and sub tropical fruits. New York: Macmillan, USA
- Ryerson K.A. (1927) Culture of the oriental persimmon in California. California Agricultural Experiment Station Bulletin 416: 63
- Silva M.J.R. et al. (2016) Phenology, yield and fruit quality of four persimmon (*Diospyros kaki* L.) cultivars in São Paulo's Midwest countryside, Brazil. African Journal of Agricultural Research 11: 5171-5177
- Tessmer M.A., Kluge R.A., Appezzato-da-Glória B. (2014) The accumulation of tannins during the development of 'Giombo' and Fuyu' persimmon fruits. Scientia Horticulturae 172: 292-299

Plum/Ameixeira

Mainly pollinators and flower visitors:

- Raj H., Mattu V.K. (2014) Diversity and distribution of insect pollinators on various temperate fruit crops in Himachal Himalaya, India. International Journal of Science and Nature 5: 626-631
- Wadhwa N., Sihag R.C. (2015) Melittophilous mode of pollination predominates in European plum (*Prunus domestica* L.) in the semi-arid environment of northwest India. Asian Journal of Agricultural Research 5: 189-207

Mainly crop pollination and production:

- Benedek P, Nyeki J. (1996) Fruit set of selected self-sterile and self-fertile fruit cultivars as affected by the duration of insect pollination. Acta Horticulturae 423: 57-63
- Dordević M., Cerović R., Nikolić D. et al. (2016) Influence of pollination mode on fruit set in plum (*Prunus domestica*) Acta Horticulturae 139: 347-352

Free, J.B. (1993) Insect pollinaiton of crops. 2ed. Academic Press: Cardiff, UK

Hassan H.S.A., Mostafa E.A.M., Enas A.M.A. (2007) Effect of self, open, and cross pollination on fruit characteristics of some plum cultivars. American-Eurasian Journal of Agriculture and Environmental

Science 2: 118-122

- Jun J.H., Chung, K.H. (2007) Interspecific cross compatibility among plum, apricot and plumcot. Korean Journal of Horticultural Science and Technology 25: 217-222
- Sapir G., Goldway M., Shafir S., Stern R.A. (2007) Multiple introduction of honey bee colonies increases cross-pollination, fruit set, and yield of 'Black Diamond' Japanese plum (*Prunus salicina* Lindl.) Journal of Horticultural Science and Biotechnology 82: 590-596

Pumpkin/Abobora

Mainly pollinators and flower visitors:

- Fronk W.D., Slater J.A. (1956) Insect fauna of cucurbits flowers. Journal of the Kansas Entomological Society 29: 141-145
- Matsumoto T., Yamazaki K. (2013) Distance from migratory honey bee apiary effects on community of insects visiting flowers of pumpkin. Bulletin of Insectology 66: 103-108
- Nicodemo D., Couto R.H.N., Malheiros E.B., De Jong D. (2009) Honey bee as an effective pollinator of pumpkins. Scientia Agricola 66: 476-480
- Pfister S.C., Eckerter P.W., Schirmel J. et al. (2017) Sensitivity of commercial pumpkin yield to potential decline among different groups of pollinating bees. Royal Society Open Science 4: 170102
- Shuler R.E., Roulston T.A.H., Farris G.E. (2005) Farming practices influence wild pollinator populations on squash and pumpkin. Journal of Economic Entomology 98: 790-795
- Walters S.A., Taylor B.H. (2006) Effects of honey bee pollination on pumpkin fruit and seed yield. Horticultural Science 41: 370-373

Mainly crop pollination and production:

Amarante C.V.T., Macedo A.F. (2000) Fruit set and fruit growth of 'Tetsukabuto' squash treated with the sodium salt of alfa-naphthaleneacetic acid. Horticultura Brasileira 18: 212-214

Robinson R.W., Decker-Walters D.S. (1997) Cucurbits. CABI International, Wallingford, UK.

Ramos S.R.R., Lima N.S., Carvalho H. de. et al. (2010) Aspectos técnicos do cultivo da abóbora na região Nordeste do Brasil. Embrapa Tabuleiros Costeiros-Documentos (INFOTECA-E), Brazil

Rambutan/Rambutã

Mainly pollinators and flower visitors:

Leão K.S. (2014) Manejo de Scaptotrigona sp. (Hymenoptera, Apidae, Meliponini) para polinização da rambuteira (*Nephelium lappaceum*) Dissertação Mestrado, Universidade Federal do Pará, Belém, Brasil, 73p., Brazil

Mainly crop pollination and production:

Andrade, R.A. (2012) Rambuteira. Revista Brasileira de Fruticultura 34: 1-2

- Lim A.L. (1984) The reproductive biology of rambutan (Nephelium lappaceum L. Sapindaceae), Garden Bulletin, Singapore 37: 181-192
- Muhamed S., Kurien S. (2018) Phenophases of rambutan (*Nephelium lappaceum* L.) based on extended BBCHscale for Kerala, India. Current Plant Biology 13: 37-44
- Rincón-Rabanales M., Roubik D.W., Guzmán M.A. et al. (2015) High yields and bee pollination of hermaphroditic rambutan (*Nephelium lappaceum* L.) in Chiapas, Mexico. Fruits 70: 23-27
- Sacramento C.K., Andrade R.A. (2014) Cultivo do Rambotã (Farming rambutan) Revista Brasileira de Fruticultura 36: 79-85

Sesame/Gergelim

Mainly pollinators and flower visitors:

- Andrade P.B. (2008) Potenciais polinizadores e requerimentos de polinização do gergelim (*Sesamum indicum*) Dissertação (Mestrado), Universidade Federal do Ceará, Fortaleza. 75p., Brazil
- Kamel S.M., Blal A.E.H., Mahfouz H.M., Said M. (2013) Pollinator fauna of sesame crop (*Sesamum indicum* L.) in Ismailia governorate, Egypt. Cercetari Agronomice In Moldova 46: 53-64

Mahfouz H.M., Kamel S.M., Belal A.H., Said H. (2012) Pollinators visiting sesame (Sesamum indicum L.) seed

crop whit reference to foraging activity of some bee species. Cercetari Agronomice In Moldova 45: 49-55

- Mahmoud F. (2012) Insects associated with sesame (Sesamun indicum L.) and the impact of insect pollinators on crop production. Pesticidi I Fitomedicina 27: 117-129
- Napoletano K. (2008) Impollinazione guidata su sesamo (*Sesamum indicum* L.) nem Nordeste del Brasile. Monografia (Graduação em Scienze Agrarie Tropicali e Subtropicali)) – Universitá Degli Studi di Firenzi, Firenzi. 100p., Italy
- Ngongolo K., Mtoka S., Rubanza C.D. (2015) Floral visitors and pollinators of sesame (*Sesamum indicum* L) from Kichi forest to the adjacent local communities' farms. Entomology and Applied Science Letters 2: 32-39
- Pashte V., Shylesha A.N. (2013) Pollinators diversity and their abundance on sesamum. Indian Journal of Entomology 57: 260-262

Mainly crop pollination and production:

- Andrade P.B., Freitas B.M., Macêdo Rocha E.E. et al. (2014) Floral biology and pollination requirements of sesame (*Sesamum indicum* L.) Acta Scientiarum. Animal Sciences Maringá 36: 93-99
- Ashri A. (2007) Sesame (Sesamum indicum L.) In: Genetic Ressources, Chromosome Engineering and Crop Improvement. Vol. 4. Oilseed crops. CRC Press, Boca Raton, FL., p.231-289, USA
- Porto V.C., Beltrão, N.E.M., Rocha M. do S. et al. (2013) O gergelim e seu cultivo no semiárido brasileiro. Natal: IFRN, 225p., Brazil
- Sarker A.M. (2004) Effect honeybee pollinisation on the yield of rapeseed, mustard and sesame. Geobros 31: 49-51

Soursop/Graviola

Mainly pollinators and flower visitors:

- Aguiar J.R., Bueno D.M., Freitas B.M. et al. (2000) Tecido nutritive em flores de gravioleira, *Annona muricata* L. Ciência Agronômica 31: 51-55
- Maia A.C.D., Carvalho A.T., Paulino-Neto H.F., Schlindwein C. (2012) Besouros (Insecta, Coleoptera) como polinizadores no Brasil Perspectivas no uso sustentado e conservação na polinização. In: Polinizadores no Brasil. São Paulo: Edusp. p.153-173, Brazil
- Vinay G.M., Sakthivel T., Priyanka H.L. (2017) Recent advances in annona breeding: a review, International Journal of Pure & Applied Bioscience 5: 1168-1181

Mainly crop pollination and production:

- Freitas A.L.G.E. (2012) Caracterização da produção e do mercado da graviola (*Annona muricata* L.) no Estado da Bahia. Vitória da Conquista BA: UESB, 108p. (Dissertação Mestrado em Agronomia, Área de Concentração em Fitotecnia), Brazil
- Jalikop S.H., Kumar, R. (2007) Pseudo-xenic effect of allied *Annona* spp. pollen in hand pollination of Cv.'Arka Sahan' [(*A.cherimola* x *A.squamosa*]. Horticultural Science 42: 1534-1538
- Silva S.E.L., Souza A.G.C. (1999) Avaliação preliminar de cinco tipos de graviola (*Annona muricata* L.) nas condições de Manaus-AM. Manaus: Embrapa Amazônia Ocidental, 10p. Boletim de Pesquisa 2, Brazil
- Worrel D.B., Carrington C.M.S., Huber D.J. (1994) Growth, maturation and pipening of (Annona muricata L.) fruit. Scientia Horticulturae 57: 7-15

Soybean/Soja

- Abrams R.I., Edwards C.R., Harris T. (1978) Yields and cross-pollination of soybeans as affected by honey bees and alfalfa leaf cutting bees. American Bee Journal 118: 555-560.
- Barella W.M., Guerreiro M. (2009) Abelhas polinizadoras na cultura da soja (*Glycine max* L.) V Congresso interno de Iniciação Científica, Anais do V CONIC, Brazil
- Chiang Y.C., Kiang Y.T. (1987) Geometric position of genotypes, honeybee foraging patterns and outcrossing in soybean. Botanical Bulletin of Academia Sinica 28: 1-11
- Chiari W.C., Toledo V.A.A., Ruvolo-Takasusuki M.C.C. et al. (2005a) floral biology and behavior of Africanized honeybees *Apis mellifera* in soybean (*Glycine max* L. Merril) Brazilian Archives of Biology and Technology 48: 367-378

- Chiari W.C., Toledo V.A.A., Ruvolo-Takasusuki M.C.C. et al. (2005b) Pollination of Soybean (*Glycine max* L. Merril) by Honeybees (*Apis mellifera* L.) Brazilian Archives of Biology and Technology 48: 31-36
- Chiari, W.C., Ruvolo-Takasusuki, M.C.C., Chambó, E.D. et al. (2011) Gene Flow Between Conventional and Transgenic Soybean Pollinated by Honeybees. In: Herbicides - Mechanisms and Mode of Action. InTech -Open Access Publisher 8: 137-152
- Erickson E.H. (1975a) Effect of honey bees on yield of three soybean cultivars. Crop Science 15: 84-86
- Erickson E.H. (1975b) Honey bees and soy bean. American Bee Journal 115: 351-372
- Erickson E.H. (1984) Soybean pollination and honey production- A research progress report. American Bee Journal 124: 775-779
- Erickson E.H., Berger G.A., Shannon J.G., Robbins J.M. (1978) Honey bee pollination increases soybean yields in the Mississippi Delta region of Arkansas and Missouri. Journal of Economic Entomology 71: 601-603
- Fávero A.C., Couto R.H.N. (2000) Polinização entomófila em soja (*Glycine max* L. var. FT2000) In: Anais do 13th Congresso Brasileiro de Apicultura. CBA, Brazil
- Issa M.R.C., Velocci M.E.P., Gonçalves L.S., Soares A. (1984) Ensaio de polinização em soja (*Glycine max*) por abelhas *Apis mellifera* L. Anais do 5 Congresso Brasileiro de Apicultura Vicosa, MG, Brazil
- Juliano J.C. (1977) Polinização entomofila na soja. Anais do 4th Congresso Brasileiro de Apicultura 1976. Curitiba, PR, Brazil
- Kettle W.D., Taylor O.R. (1979) Ecological interactions of honey bees and soybeans. Journal of the Kansas Entomological Society 52: 549
- Koelling P.D., Kenworthy W.J., Caron D.M. (1981) Pollination of male-sterile soybeans in caged plots. Crop Science 21: 559-561
- Mason C.E. (1979) Honey bees foraging activity on soybeans in Delaware. Proceedings of the IVth International Symposium on Pollination, Md. Agric. Exp. Sta. Spec. Misc. Publ. 1: 117-122, USA
- Milfont M.O. (2012) Uso da abelha melífera (*Apis mellifera* L.) na polinização e aumento de produtividade de grãos em variedade de soja [*Glycine max* (L.) Merril.] adaptada às condições climáticas do nordeste brasileiro. Thesis (Doctorate), Universidade Federal do Ceará, 129p., Brazil
- Ortiz-Perez E., Mian R.M.A., Cooper R.L. et al. (2008) Seed-set evaluation of four male-sterile, female-fertile soybean lines using alfalfa leafcutting bees and honey bees as pollinators. Journal of Agricultural Science 146: 461-469
- Pinzauti M., Frediani D. (1980) The importance of honeybee pollination in soya bean production. Apicoltore Moderno 71: 155-160
- Rust R.W., Mason C.E., Erickson E.H. (1980) Wild bees on soybeans, *Glycine max*. Environmental Entomology 9: 230-232
- Sheppard W.S., Jaycox E.R., Parise S.G. (1979) Selection and management of honey bees for pollination of soybeans. Proceedings of the IVth International Symposium on Pollination, Md. Agric. Exp. Sta. Spec. Misc. Publ. 1: 123-130, USA
- Toledo V.A.A., Malerbo-Souza D.T., Selegatto-Filho J.C. et al. (2011) Biodiversidade de agentes polinizadores e seu efeito na produção de grãos em soja var. Mon Soy 3329. Revista Varia Scientia Agrárias 2: 123-130

- Milfont M.O., Rocha E.E., Lima A.O., Freitas B.M. (2013) Higher soybean production using honeybee and wild pollinators, a sustainable alternative to pesticides and autopollination. Environmental Chemistry Letters 11: 335-341
- Gazzoni D.L. (2017) Soybean and bees. Embrapa, 147p. Brasília, DF. ISBN 978-85-7035-592-8, Brazil
- Masuda T., Goldsmith P.D. (2009) World soybean production: area harvested, yield, and long-term projections. International Food and Agribusiness Managment Review 12: 143-162

Strawberry/Morango

Mainly pollinators and flower visitors:

Abrol D.P., Gorka A.K., Ansari M.J. et al. (2017) Impacts of insect pollinators on yield and fruit quality of strawberry. Saudi Journal of Biological Sciences 26: 524-530

- Antonelli A.L., Mayer D.F., Burgett D.M., Sjulin .S. (1988) Pollinating insects and strawberry yields in the Pacific Northwest. American Bee Journal 128: 618-620
- Antunes O.T., Calvete E.O., Rocha H.C. et al. (2007) Produção de cultivares de morangueiro polinizadas pela abelha jataí em ambiente protegido. Horticultura Brasileira 25: 94-99
- Chang Y.D., Lee M.Y., Mah Y.I. (2001) Pollination on strawberry in the vinyl house by *Apis mellifera* L. and *A. cerana* Fab. Acta Horticulturae 561: 257-262
- Chen W.F., An J.D., Dong J. et al. (2011) Flower-visiting behavior and pollination ecology of different bee species on greenhouse strawberries. Chinese Journal of Ecology 30: 290-296
- Connor L.J. (1975) The role of cultivar in insect pollination of strawberries. 3rd International Symposium on Pollination, p.149-154, USA
- Goodman R.D., Oldroyd B.P. (1988) Honeybee pollination of strawberries (*Fragaria* x *ananassa* Duchesne) Australian Journal of Experimental Agriculture 28: 435-438
- Jacobs F.J., Houbaert D., De Rycke P.H. (1988) Pollinisation des fraises (*Fragaria* x ananassa) sous verre par des abeilles (*Apis mellifera*) Agricontact 193
- Kakutani T., Inoue T., Tezuka T., Maeta Y. (1993) Pollination of strawberry by the stingless bee, *Trigona minangkabau*, and the honey Bee, *Apis mellifera*: an experimental study of fertilization efficiency. Researches on Popululation Ecology 35: 95-111
- Klatt B.K., Holzschuh A., Westphal C. et al. (2013) Bee pollination improves crop quality, shelf life and commercial value. Proceedings of the Royal Society, Series B 281: 20132440
- Matsuka M., Sakai T. (1989) Bee pollination in Japan with special reference to strawberry production in greenhouses. Bee World 70: 55-61
- McGregor S.E. (1976) Insect pollination of cultivated crop plants. Agricultural Research Service, 411p., US Department of Agriculture, USA
- Nye W.P., Anderson J.L. (1974) Insect pollinators frequenting strawberry blossoms and the effect of honey bees on yield and fruit quality. Journal of the American Society of Horticultural Science 99: 40-41
- Oliveira D., Savoie L., Vincent C. (1991) Pollinators of dultivated strawberry in Québec. Acta Horticulturae 288: 420-424
- Pion S., Oliveira D. (1980) Agents pollinisateurs et productivité du fraisier 'Redcoat', Fragaria x ananassa Duch. Phytoprotection 61: 72-78
- Roselino A.C., Santos S.B., Hrncir M., Bego L.R. (2009) Differences between the quality of strawberries (*Fragaria* x ananassa) pollinated by the stingless bees *Scaptotrigona* aff. depilis and Nannotrigona testaceicornis. Genetics and Molecular Research 8: 539-545

- Chagnon M., Gingras J, Oliveira D. (1993) Complementary aspects of strawberry pollination by honey and indigenous bees (Hymenoptera) Journal of Economic Entomology 86: 416-420
- Castle D., Grass I, Westphal C. (2019) Fruit quantity and quality of strawberries benefit from enhanced pollinator abundance at hedgerows in agricultural landscapes. Agriculture Ecosystems & Environment 275: 14-22
- Malagodi-Braga K.S., Kleinert A.M.P. (2004) Could *Tetragonisca angustula* Latreille (Apinae, Meliponini) be effective as strawberry pollinator in greenhouses? Australian Journal of Agricultural Research 55: 771-773
- Malagodi-Braga K.S., Kleinert A.M.P. (2007) How bee behavior on strawberry flower (*Fragaria ananassa* Duchesne) can influence fruit development? Bioscience Journal 23: 76-81

Sunflower/Girassol

- Basualdo M., Bedascarrasbure E., de Jong D. (2000) Africanized honey bees (Hymenoptera: Apidae) have a greater fidelity to sunflowers than European bees. Journal of Economic Entomology 93: 304-307
- Bhowmik B., Bhadra K. (2015) Insect pollinators and their role on crop yield and quality of sunflower (*Helianthus annuus*, PAC-361) from West Bengal, India. International Journal of Current Science 18: 76-87
- Calmasur O., Ozbek H. (1999) Pollinator bees (Hymenoptera, Apoidea) on sunflower (*Helianthus annuus* L.) and their effects on seed setting in the Erzurum Region. Turkish Journal of Biology 23: 73-89

- Carvalheiro L.G., Veldtman R., Shenkute A.G. et al. (2011) Natural and within-farmland biodiversity enhances crop productivity. Ecology Letters 14: 251-259
- Cockerell T.D.A. (1914) Bees visiting Helianthus. The Canadian Entomologist 46: 409-415.
- Cruz D.O., Freitas B.M. (2013) Diversidade de abelhas visitantes florais e potenciais polinizadores de culturas oleaginosas no Nordeste do Brasil. Ambiência 9: 411-418
- DeGrandi-Hoffman G., Chambers M. (2006) Effects of honey bee (Hymenoptera: Apidae) foraging on seed set in self-fertile sunflowers (*Helianthus annuus* L.) Environmental Entomology 35: 1103-1108
- DeGrandi-Hoffman G., Martin J.H. (1993) The size and distribution of the honey bee (*Apis mellifera* L.) crosspollinating population on male-sterile sunflowers (*Helianthus annuus* L.) Journal of Apicultural Research 32: 135-142
- DeGrandi-Hoffman G., Watkins J.C. (2000) The foraging activity of honey bees *Apis mellifera* and non-Apis bees on hybrid sunflowers (*Helianthus annuus*) and its influence on cross-pollination and seed set. Journal of Apicultural Research 39: 37-45
- Delaude A., Tasei J.N., Rollier M. (1979) Pollinator insects of sunflower (*Helianthus annuus* L.) in France. Pollination of sterile lines for hybrid seed production. Proceedings of the IVth International Symposium on Pollination, Md. Agric. Exp. Sta. Spec. Misc. Publ. 1: 29-40, USA
- Fell R.D. (1986) Foraging behaviors of Apis mellifera L. and *Bombus* spp. on oilseed sunflower (*Helianthus annuus* L.) Journal of the Kansas Entomological Society 59: 72-81
- Free J.B. (1964) The behaviour of honeybees on sunflowers (*Helianthus annuus* L.) Journal of Applied Ecology 1: 19-27
- Furgala B., Noetzel D.M., Robinson R.G. (1979) Observations on the pollination of hybrid sunflowers. In: Proceedings of the IVth International Symposium on Pollination, Md. Agric. Exp. Sta. Spec. Misc. Publ. 1: 45-48, USA
- Greenleaf S.S., Kremen C. (2006) Wild bees enhance honey bees' pollination of hybrid sunflower. Proceedings of the National Academy of Sciences USA 103: 13890-13895
- Krause G.L., Wilson W.T. (1981) Honey bee pollination and visitation patterns on hybrid oilseed sunflowers in Central Wyoming (Hymenoptera: Apidae) Journal of the Kansas Entomological Society 54: 75-82
- Langridge D.F., Goodman R.D. (1981) Honeybee pollination of sunflower cultivars Hysun 30 and Sunfola. Australian Journal Experimental Agriculture and Animal Husbandry 21: 435-438
- Machado C.S., Carvalho C.A.L. (2006) Abelhas (Hymenoptera: Apoidea) visitantes dos capítulos de girassol no recôncavo baiano. Ciência Rural 36: 1404-1409
- Nderitu J., Nyamasyo G., Kasina M., Oronje M.L. (2008) Diversity of sunflower pollinators and their effect on seed yield in Makueni District, eastern Kenya. Spanish Journal of Agricultural Research 6: 271-278
- Neiva I.S. (2009) Abelhas visitantes das flores do girassol *Helianthus annus* Linnaeus (Asterales: Asteraceae) em Dourados-MS, Brasil. Dissertação (Mestrado), Universidade Federal da Grande Dourados, Dourados, Brazil, 28p., Brazil
- Oz M., Karasu A., Cakmak I. et al. (2009) Effects of honeybee (*Apis mellifera*) pollination on seed set in hybrid sunflower (*Helianthus annuus* L.) African Journal of Biotechnology 8: 1037-1043
- Parker F.D. (1981a) How efficient are bees in pollinating sunflowers? Journal of the Kansas Entomological Society 54: 61-67
- Parker F.D. (1981b) Sunflower pollination: abundance, diversity, and seasonality of bees on male-sterile and male-fertile cultivars. Environmental Entomology 10: 1012-1017
- Parker F.D., Frohlich D.R. (1983) Hybrid sunflower pollination by a manageable composite specialist: the sunflower leafcutter bee (Hymenoptera: Megachilidae) Environmental Entomology 12: 576-581
- Pisanty G., Klein A.M., Mandelik Y. (2014) Do wild bees complement honeybee pollination of confection sunflowers in Israel. Apidologie 45: 235-247
- Saez A., Sabatino M., Aizen M.A. (2012) Interactive effects of large- and small-scale sources of feral honey bees for sunflower in the Argentine Pampas. PLoS ONE 7: e30968
- Skinner J.A. (1987) Abundance and spatial distribution of bees visiting male-sterile and male-fertile sunflower cultivars in California. Environmental Entomology 16: 922-927
- Tepedino V.J., Parker F.D. (1982) Interspecific differences in the relative importance of pollen and nectar to

bee species foraging on sunflowers. Environmental Entomology 11: 246-250

- Toledo V.A.A., Chambó E.D., Halak A.L. et al. (2011) Biologia floral e polinização em girassol (*Helianthus annuus* L.) por abelhas africanizadas. Scientia Agraria Paranaenis 10: 5-17
- Wilson M.E., Skinner J.A., Wszelaki A.L., Drummond F. (2016) Using nonmetric multidimensional scaling to analyze bee visitation in east Tennessee crops as an indicator of pollination services provided by honey bees (*Apis mellifera* L.) and native bees. Environmental Entomology 45: 390-396

Mainly crop pollination and production:

- Castro C., Leite R.M.V.B.C. (2018) Main aspects of sunflower production in Brazil. OCL 25: D104
- Chambó E.D., Garcia R.C., Oliveira N.T.E., Duarte-Júnio J.B. (2011) Honey bee visitation to sunflower: effects on pollination and plant genotype. Scientia Agricola 68: 647-651
- Greenleaf S.S. & Kremen C. (2006) Wild bees enhance honey bees' pollination of hybrid sunflower. Proceedings of the National Academy of Sciences 113: 13890-13895
- Morgado L.N., Carvalho C.F., Souza B., Santana M.P. (2002) Fauna of bees (Hymenoptera: Apoidea) on sunflower flowers, *Helianthus annuus* L. Ciência Agrotecnologia 26: 1167-1177

Sweet pepper/Pimentão

Mainly pollinators and flower visitors:

- Raw A. (2000) Foraging behaviour of wild bees at hot pepper flowers (*Capsicum annuum*) and its possible influence on cross pollination. Annals of Botany 85: 487-492
- Roselino A.C., Santos S.A.B., Bego L.R. (2010) Qualidade dos frutos de pimentão (*Capsicum annuum* L.) a partir de flores polinizadas por abelhas sem ferrão (*Melipona quadrifasciata anthidioides* Lepeletier 1836 e *Melipona scutellaris* Latreille 1811) sob cultivo protegido. Revista Brasileira de Biociências 8: 154-158
- Roselino A.C. (2005) Polinização em culturas de pimentão Capsicum annuum por Melipona quadrifasciata anthidioides e Melipona scutellaris e de morango – Fragaria x ananassa por Scaptotrigona aff. depilis e Nannotrigona testaceicornis (Hymenoptera, Apidae, Meliponini) Dissertação (Mestrado em Ciências), Universidade de São Paulo, Ribeirão Preto, Brazil, 95p., Brazil
- Silva E.M.S, Freitas B.M., Siva L.A. et al. (2005) Biologia floral do pimentão (*Capsicum annuum*) e a utilização da abelha jandaíra (*Melipona subnitida* Ducke) como polinizador em cultivo protegido. Revista Ciência Agronômica, Fortaleza, 36: 386-390

- Cruz D.O., Freitas B.M., Silva L.A. et al. (2004) Adaptação e comportamento de pastejo da abelha jandaíra (*Melipona subnitida* Ducke) em ambiente protegido. Acta Scientiarum -Animal Sciences 26: 293-298
- Cruz D.O., Freitas B.M., Silva L.A. et al. (2005) Pollination efficiency of the stingless bee *Melipona subnitida* on greenhouse sweet pepper. Pesquisa Agropecuaria Brasileira 40: 1197-1201
- Dag A., Kammer Y. (2001) Comparison between the effectiveness of honey bee (*Apis mellifera*) and bumble bee (*Bombus terrestris*) as pollinators of greenhouse sweet pepper (*Capsicum annuum*) American Bee Journal 141: 447-448
- Dag A., Zvieli Y., Afik O., Elkind Y. (2007) Honeybee pollination affects fruit characteristics of sweet pepper grown under net-houses. International Journal of Vegetable Science 13: 45-59
- De Ruijter A., van den Eijnde J., van der Steen J. (1991) Pollination of sweet pepper (*Capsicum annuum* L.) in greenhouses by honeybees. Acta Horticulturae 288: 270-274
- Faria-Júnior L.R.R., Bendini J.N., Barreto L.M.R.C. (2008) Eficiência polinizadora de Apis mellifera L. e polinização entomófila em pimentão 'Cascadura Ikeda'. Bragantia 67: 261-266
- Jarlan A., Oliveira D., Gingras J. (1997) Pollination of sweet pepper (*Capsicum annuum* L.) in greenhouse by the syrphid fly *Eristalis tenax* L. Acta Horticulturae 437: 335-340
- Pereira C., Cruz Taques, T., Valim, J.O.S. et al. (2015) The management of bee communities by intercropping with flowering basil (*Ocimum basilicum*) enhances pollination and yield of bell pepper (*Capsicum annuum*) Journal of Insect Conservervation 19: 479-48.
- Reifschneider, FJ.B. (2000) Capsicum pimentas e pimentões no Brasil. Brasília: Embrapa-SNT; Embrapa-CNPH, 113p., Brazil

Tomato/Tomate

Mainly pollinators and flower visitors:

Banda H.J., Paxton R.J. (1991) Pollination of greenhouse tomatoes by bees. Acta Horticulturae 288: 194-198

- Bin E, Soressi G.P. (1973) Insetti pronubi e produzione di seme ibrido di pomodoro. Genetica Agraria 27: 35-74
- Bohart G.E., Todd F.E. (1961) Pollination of seed crops by insects. In: Seeds. The yearbook of agriculture. U.S. Government Printing Office, p.240-246, Washington DC, USA
- Cauich O., Quezada-Euán J.J.G., Macias J.O. et al. (2004) Behavior and pollination efficiency of Nannotrigona perilampoides (Hymenoptera: Meliponini) on greenhouse tomatoes (Lycopersicon esculentum) in Subtropical México. Horticultural Entomology 97: 475-481
- Cribb D. (1990) Pollination of tomato crops by honeybees. Bee Craft 72: 228-231
- Del Sarto M.C.L., Peruquetti R.C., Campos L.A.O. (2005) Evaluation of the neotropical stingless bee *Melipona quadrifasciata* (Hymenoptera: Apidae) as pollinator of greenhouse tomatoes. Journal of Economic Entomology 98: 260-266
- Deprá M.S., Delaqua G.C.G., Freitas L., Gaglianone M.C. (2014) Pollination deficit in open-field tomato crops (Solanum hycropersicum L. Solanaceae) In Rio De Janeiro State, South Brazil. Journal of Pollination Ecology 12: 1-8
- Eijnde J., Ruijter A. (1989) Pollination of glasshouse tomatoes by honeybees. Apidologie 20: 492-493
- Franceschinelli E.V., Neto C.M.S., Lima F.G. et al. (2013) Native bees pollinate tomato flowers and increase fruit production. Journal of Pollination Ecology 11: 41-45
- Higo H.A., Rice N.D., Winston M.L., Lewis B. (2004) Honey bee (Hymenoptera: Apidae) distribution and potential for supplementary pollination in commercial tomato greenhouses during winter. Journal of Economic Entomology 97: 163-170
- Sabara H.A., Gillepsie D.R., Elle E., Winston M.L. (2004) Influence of brood, vent screening, and time of year on honey bee (Hymenoptera: Apidae) pollination and fruit quality of greenhouse tomatoes. Journal of Economic Entomology 97: 727-734
- Sabara H.A., Winston M.L. (2003) Managing honey bees (Hymenoptera: Apidae) for greenhouse tomato pollination. Journal of Economic Entomology 96: 547-554
- Santos S.A.B., Roselino A.C., Hrncir M., Bego L.R. (2009) Pollination of tomatoes by the stingless bee Melipona quadrifasciata and the honey bee Apis mellifera (Hymenoptera, Apidae) Genetics and Molecular Research 8: 751-757
- Santos A.O., Bartelli B.F., Nogueira-Ferreira F.H. (2014) Potential pollinators of tomato, *Lycopersicon esculentum* (Solanaceae) in open crops and the effect of a solitary bee in fruit set and quality. Journal of Economic Entomology 107: 987-94
- Silva-Neto C., Lima E.G., Gonçalves B.B. et al. (2013) Native bees pollinate tomato flowers and increase fruit production. Journal of Pollination Ecology 11: 41-45
- Silva-Neto C.M., Bergamini L.L., Elias M.A.S. et al. (2017) High species richness of native pollinators in Brazilian tomato crops. Brazilian Journal of Biology 77: 506-513
- Vinícius-Silva R., Parma D.D.F., Tostes R.B. et al. (2017) Importance of bees in pollination of Solanum lycopersicum L. (Solanaceae) in open-field of the southeast of Minas Gerais State, Brazil. Hoehnea 44: 349-360

- Bezerra E.L.S., Machado I.C. (2003) Biologia floral e sistema de polinização de *Solanum stramonifolium* Jacq. (Solanaceae) em remanescente de mata atlântica, Pernambuco. Acta Botanica Brasiliense 17: 247-257
- Del Sarto M.C.L., Peruquetti R.C., Campos L.A.O. (2004) Polinização em ambiente protegido: uso da abelhasem-ferrão mandaçaia na polinização do tomateiro em sistema orgânico de produção. In: Cultivo em Ambiente Protegido -Histórico, Biotecnologia e Perspectivas. UFV, Viçosa, Brazil
- Gaglianone M.C., Campos L.A.O. (2015) Plano de manejo para os polinizadores do tomateiro. Rio de Janeiro: Funbio. 48p., Brazil

Watermelon

- Adlerz W.C. (1966) Honey bee visit numbers and watermelon pollination. Journal of Economic Entomology 59: 28-30
- Alencar L.C., Moura S.G. (2013) Efeitos de abelhas na frutificação e qualidade de melancia (cv. Crimson Sweet) na região Central do estado do Piauí. 64p. Jaboticabal, São Paulo, Brazil
- Azo'o M.E., Messi J., Tchuenguem Fohouo F.N. et al. (2010) Foraging behaviour of *Apis mellifera adansonii* and its impact on pollination, fruit and seed yields of *Citrullus lanatus* at Nkolbisson (Yaoundé, Cameroon) Cameroon Journal of Experimental Biology 6: 41-48
- Brewer J.W. (1974) Pollination requirements for watermelon seed production. Journal of Apicultural Research 13: 207-2012
- Bussman R., Njoroge G.N., Gemmill B. et al. (2003) Some applied aspects of pollination for increased fruit and seed productivity with special reference to *Citrullus lanatus* (Watermelon). African Crop Science Congress Proceedings 6: 108-112
- Goff C.C. (1937) The importance of bees in the production of watermelons. The Florida Entomologist 20: 30-31
- Kaziev I.P., Seidova S.S. (1965) La production de nectar des fleurs de certaines especes de cucurbitacees dans les conditions de l'Azerbaidjan. Azerbaidjan
- Kremen C., Williams N.M., Bugg R.L. et al. (2004) The area requirements of an ecosystem service: crop pollination by native bee communities in California. Ecology Letters 7, 1109-1119
- Kremen C., Williams N.M., Thorp R.W. (2002) Crop pollination from native bees at risk from agricultural intensification. Proceedings of the National Academy of Science 99: 16812-16816
- Meléndez-Ramirez V., Magaña-Rueda S., Parra-Tabla V. et al. (2002) Diversity of native bee visitors of cucurbit crops (Cucurbitaceae) in Yucatán, México. Journal of Insect Conservation, 6: 135-147
- Mohamed M.I., El-Hafez A. (1974) Visitation of honey bees to some species of Cucurbitaceae. Annals of Agricultural Science 2: 269-275
- Njoroge G.N., Gemmill B., Bussmann R. et al. (2004) Pollination ecology of *Citrullus lanatus* at Yatta, Kenya. International Journal of Tropical Insect Science 24: 73-77
- Njoroge G.N., Gemmill B., Bussmann R. et al. (2010) Diversity and efficiency of wild pollinators of watermelon (*Citrullus lanatus* (Thunb.) Mansf.) at Yatta (Kenya) The Society for Advancement of Horticulture 12: 35-41
- Pinkus-Rendon M.A., Parra-Tabla V., Melendez-Ramirez V. (2005) Floral resource use and interactions between *Apis mellifera* and native bees in cucurbit crops in Yucatan, Mexico. The Canadian Entomologist 137: 441-449
- Pisanty G. (2014) Pollination services in mediterranean natural-agricultural systems: assessing wild bees' contribution to crop pollination and the effects of land use on them. PhD Thesis, Hebrew University of Jerusalem, Israel
- Smith A.A., Bentley M., Reynolds H.L. (2013) Wild bees visiting cucumber on midwestern U.S. organic farms benefit from near-farm semi-natural areas. Journal of Economic Entomology 106: 97-106
- Spangler H.G., Moffett J.O. (1979) Pollination of melons in greenhouses. Gleanings in Bee Culture 107: 17-18
- Stanghellini M.S., Ambrose J.T., Schultheis J.R. (1998a) Using commercial bumble bee colonies as backup pollinators for honey bees to produce cucumbers and watermelons. HortTechnology 8: 590-594
- Stanghellini M.S., Ambrose J.T., Schultheis J.R. (1998b) Seed production in watermelon: a comparison between two commercially available pollinators. HortScience 33: 28-30
- Stanghellini M.S., Ambrose J.T., Schultheis J.R. (2002a) Diurnal activity, floral visitation and pollen deposition by honey bees and bumble bees on field-grown cucumber and watermelon. Journal of Apicultural Research 40: 27-34
- Stanghellini M.S., Schultheis J.R., Ambrose J.T. (2002b) Pollen mobilization in selected Cucurbitaceae and the putative effects of pollinator abundance on pollen depletion rates. HortScience 127: 729-736
- Walters S.A., Schultheis J.R. (2009) Directionality of pollinator movements in wwatermelon plantings. HortScience 44: 49-52

- Wilson M.E., Skinner J.A., Wszelaki A.L., Drummond F. (2016) Using nonmetric multidimensional scaling to analyze bee visitation in east Tennessee crops as an indicator of pollination services provided by honey bees (*Apis mellifera* L.) and native bees. Environmental Entomology 45: 390-396
- Winfree R., Williams N.M., Gaines H. et al. (2008) Wild bee pollinators provide the majority of crop visitation across land-use gradients in New Jersey and Pennsylvania, USA. Journal of Applied Ecology 45: 793-802

Bomfim I.G.A., Bezerra A.D., Nunes A. et al. (2015) Requerimentos de polinização de variedades de minimelancia com e sem semente cultivadas em ambiente protegido. Pesquisa Agropecuária Brasileira 50: 44-53

Walters S.A. (2005) Honey bee pollination requirements for ttriploid watermelon. HortScience 40: 1268-1270

Zucchini

Mainly pollinators and flower visitors:

- Artz D.R., Nault B.A. (2011) Performance of *Apis mellifera*, *Bombus impatiens*, and *Peponapis pruinosa* (Hymenoptera: Apidae) as pollinators of pumpkin. Journal of Economic Entomology 94: 609-616
- Baptista C.F. (2016) Polinização de *Cucurbita pepo* (Cucurbitaceae) por *Melipona quadrifasciata* (Hymenoptera: Apidae: Meliponini) em cultivo protegido, Master Thesis, Universidade Federal de Viçosa, Brazil
- Enriquez E., Ayala R., Gonzalez V.H., Nunez-Farfan J. (2015) Alpha and beta diversity of bees and their pollination role on *Cucurbita pepo* L. (Cucurbitaceae) In: The Guatemalan Cloud Forest. The Pan-pacific Entomologist 91: 211-222
- Fronk W.D., Slater J.A. (1956) Insect fauna of cucurbits flowers. Journal of the Kansas Entomological Society 29: 141-145
- Grewal G.S., Sidhu A.S. (1978) Insect-pollinators of some cucurbits in Punjab. Indian Journal of Agricultural Sciences 48: 79-83
- Julier H.E., Roulston T.A.H. (2009) Wild bee abundance and pollination service in cultivated pumpkins: Farm management, nesting behavior and landscape effects. Journal of Economic Entomology 102: 563-573
- Krug C., Alves-dos-Santos I., Cane J (2010) Visiting bees of cucurbita flowers (Cucurbitaceae) with emphasis on the presence of *Peponapis fervens* Smith. (Eucerini-Apidae) Santa Catarina, South Brazil. Oecologia Australis 14: 128-139
- Kubisova (1974) L'abeille, pollinisatrice des cucurbitacees. Isntitut für Bienenzucht 18: 155-163
- Nepi M., Pacini E. (1993) Pollination, pollen viability and pistil receptivity in *Cucurbita pepo*. Annals of Botany 72: 527-536
- Petersen J.D., Huseth A.S., Nault B.A. (2014) Evaluating pollination deficits in pumpkin production in New York. Environmental Entomology 43: 1247-1253
- Petersen J.D., Reiners S., Nault B.A. (2013) Pollination services provided by bees in pumpkin fields supplemented with either *Apis mellifera* or Bombus impatiens or not supplemented. PLoS ONE 8: e69819
- Phillips B. (2013) The ecological impacts of non-native annual and native perennial floral insectaries on beneficial insect activity density and arthropod-mediated ecosystem services within Ohio pumpkin (*Cucurbita pepo*) agroecosystems. Master Thesis, Ohio State University, USA
- Phillips B.W., Gardiner M. (2015) Use of video surveillance to measure the influences of habitat management and landscape composition on pollinator visitation and pollen deposition in pumpkin (*Cucurbita pepo*) agroecosystems. PeerJ 3: e1342
- Tepedino V.J. (1981) The pollination efficiency of the squash bee (*Peponapis pruinosa*) and the honey bee (*Apis mellifera*) on summer squash (*Cucurbita pepo*) Journal of the Kansas Entomological Society 54: 359-377
- Torezani K.R. (2015) Polinização da aboboreira (*Cucurbita pepo* L.): um estudo sobre a comunidade de abelhas em sistemas orgânicos e convencionais de produção no Distrito Federal. Thesis, Universidade de Brasília, Brazil
- Vidal M.G., De Jong D., Wien H.C., Morse R.A. (2010) Pollination and fruit set in pumpkin (*Cucurbita pepo*) by honey bees. Brazilian Journal of Botany 33: 106-113
- Walters S.A., Taylor B.H. (2006) Effects of honey bee pollination on pumpkin fruit and seed yield. HortScience 41: 370-373

Mainly crop pollination and production:

Delaplane K.S., Mayer D.R., Mayer D.F. (2000) Crop pollination by bees. 352p. Cabi, USA

Photography credits

© Alexandra-Maria Klein: © Betina Blochtein: © Breno Freitas: © Carolina F. Cardoso: © Cristiano Menezes: © Denise Alves: © Diego Bezerra: © Felix Fornoff: © Isac Bomfim: © Jânio Angelo Felix: © João Vinicius S. Della Vecchia © Juliana Hipolito: © Kátia Sampaio Malagodi-Braga: © Kristina Krewenka: © Darclet Malerbo Souza: © Marcelo Cavalcante: © Marcelo Milfont: C Márcia Maués: © Marcos Venâncio Lima: © Mikail Oliveira:

© Patrícia Barreto:

© Rômulo Rizzardo:

© Thiago Mahlmann:

© Victor Monteiro:

Pages

vii, 115t 26, 27, 34, 35, 111ml 25br, 38ml, 39tr, 39br, 40, 41tr, 45, 64, 72, 73, 98bl, 110mbl, 110bl, 111tm, 113mtr, 114tl, 114tr, 116tm, 116tr, 117 10t iv, ix, x, 10m, 20, 21, 43b, 92, 93 111tr 65 94, 115m, 115b 100, 101, 111br, 116mbl 9b, 114bl 60tl 46ml, 47tr 102, 103 93bl 51, 50bl, 68bl, 69br, 76ml 30, 31, 113tr, 116tl 90, 91t, 91ml, 111mr 10b, 54ml, 55br, 66, 67 82,83 cover, 8, 9t, 22, 23, 24, 25tr, 28, 29, 32, 33, 36, 37, 41br, 42, 43t, 44bl, 48, 49, 50ml, 52, 53, 54bl, 54tr, 55bl, 55tr, 56, 57, 58, 59, 60bl, 61, 62, 63, 68ml, 69tr, 70, 71, 73, 75, 76bl, 77, 78, 79, 80, 81, 84, 85, 88, 89tr, 95, 96, 97, 98ml, 99, 110mtl, 110tl, 111tl, 111bl, 111bm, 112, 113br, 114mtl, 114mbl, 116mbl 86, 87, 111mm, 114tm, 116mtl 38bl, 39bl 46bl, 113mbr 89br